

Anabelian Phenomena in Geometry and Arithmetic

Florian Pop, University of Pennsylvania

PART I: Introduction and motivation

The term “anabelian” was invented by GROTHENDIECK, and a possible translation of it might be “beyond Abelian”. The corresponding mathematical notion of “anabelian Geometry” is vague as well, and roughly means that under certain “anabelian hypotheses” one has:

* * * Arithmetic and Geometry are encoded in Galois Theory * * *

It is our aim to try to explain the above assertion by presenting/explaining some results in this direction. For GROTHENDIECK’s writings concerning this the reader should have a look at [G1], [G2].

A) First examples:

a) *Absolute Galois group and real fields*

Let K be an arbitrary field, K^a an algebraic extension, K^s the separable extension of K inside K^a , and finally $G_K = \text{Aut}(K^s|K) = \text{Aut}(K^a|K)$ the absolute Galois group of K . It is a celebrated well known Theorem by ARTIN–SCHREIER from the 1920’s which asserts the following: *If G_K is a finite non-trivial group, then $G_K \cong G_{\mathbb{R}}$ and K is real closed.* In particular, $\text{char}(K) = 0$, and $K^a = K[\sqrt{-1}]$. Thus the non-triviality + finiteness of G_K imposes very strong restrictions on K . Nevertheless, the kind of restrictions imposed on K are not on the *isomorphism type of K* as a field, as there is a big variety of isomorphy types of real closed fields (and their classification up to isomorphism seems to be out of reach). The kind of restriction imposed on K is rather one concerning the algebraic behavior of K , namely that the algebraic geometry over K looks like the one over \mathbb{R} .

b) *Fundamental group and topology of complex curves*

Let X be a smooth complete curve over an algebraically closed field of characteristic zero. Then using basic results about the structure of algebraic fundamental groups, it follows that the geometric fundamental group $\pi_1(X)$ of X is isomorphic – as a profinite group – to the profinite completion $\widehat{\Gamma}_g$ of the fundamental group Γ_g of the compact orientable topological surface of genus g . Hence $\pi_1(X)$ is the profinite group on $2g$ generators σ_i, τ_i ($1 \leq i \leq g$) subject to the unique relation $\prod_i [\sigma_i, \tau_i] = 1$. In particular, the genus g of the curve X is encoded in $\pi_1(X)$. But as above, the isomorphy type of the curve X , i.e., of the

object under discussion, is not “seen” by its geometric fundamental group $\pi_1(X)$ (which in some sense corresponds to the absolute Galois group of the field K). Precisely, the restriction imposed by $\pi_1(X)$ on X is of topological nature (one on the complex points $X(\mathbb{C})$ of the curve).

B) Galois characterization of global fields

More than forty years after the result of ARTIN–SCHREIER, it was NEUKIRCH who realized (in the late 1960’s) that there must be a *p-adic* variant of the Artin–Schreier Theorem; and that such a result would have highly interesting consequences for the arithmetic of number fields (and more general, global fields). The situation is as follows: In the notations from a) above, suppose that K is a field of algebraic numbers, i.e., $K \subset \mathbb{Q}^a$. Then the Artin–Schreier Theorem asserts that if G_K is finite and non-trivial, then K is isomorphic to the field of real algebraic numbers $\mathbb{R}^{\text{abs}} = \mathbb{R} \cap \mathbb{Q}^a$. This means that the only finite non-trivial subgroups of $G_{\mathbb{Q}}$ are the ones generated by the $G_{\mathbb{Q}}$ -conjugates of the complex conjugation; in particular, all such subgroups have order 2, and their fixed fields are the conjugates of the field of real algebraic numbers. Now the idea of NEUKIRCH was to understand the fields of algebraic numbers $K \subset \mathbb{Q}^a$ having absolute Galois group G_K isomorphic (as profinite group) to the absolute Galois group $G_{\mathbb{Q}_p}$ of the *p*-adics \mathbb{Q}_p . Note that $G_{\mathbb{Q}_p}$ is much more complicated than $G_{\mathbb{R}}$. It is nevertheless a topologically finitely generated field, and its structure is relatively known, by work of JAKOVLEV, POITOU, JANNSEN–WINGBERG, ETC, see e.g. [J–W]. Finally, NEUKIRCH proved the following surprising result, which in the case of subfields $K \subset \mathbb{Q}$ is the perfect *p*-adic analog of the Theorem of Artin–Schreier:

Theorem (See e.g. NEUKIRCH [N1]).

For fields of algebraic numbers $K, K' \subset \mathbb{Q}^a$ the following hold:

(1) *Suppose that $G_K \cong G_{\mathbb{Q}_p}$. Then K is the decomposition field of some prolongation of $|\cdot|_p$ to \mathbb{Q}^a . Or equivalently, K is $G_{\mathbb{Q}}$ -conjugated to the field of algebraic *p*-adic numbers $\mathbb{Q}_p^{\text{abs}}$.*

(2) *Suppose that $G_{K'}$ is isomorphic to an open subgroup of $G_{\mathbb{Q}_p}$. Then there exists a unique $K \subset \mathbb{Q}^a$ as at (1) above such that K' is a finite extension of K .*

The Theorem above has the surprising consequence that an *isomorphism of Galois groups* of number fields gives rise functorially to an *arithmetical equivalence* of the number fields under discussion. The precise statement is as follows: For number fields K , let $\mathcal{P}(K)$ denote the set of their places. Let $\Phi : G_K \rightarrow G_L$ be an isomorphism of Galois groups of number fields. Then a consequence of the above Theorem reads: *Φ maps the decomposition groups of the places of K isomorphically onto the decomposition places of L . This bijection respects the arithmetical invariants $e(\mathfrak{p}|p), f(\mathfrak{p}|p)$ of the places $\mathfrak{p}|p$, thus defines an arithmetical*

equivalence:

$$\varphi : \mathcal{P}(K) \rightarrow \mathcal{P}(L).$$

Finally, applying basic facts concerning arithmetical equivalence of number fields, one gets: In the above context, suppose that $K|\mathbb{Q}$ is a Galois extension. Then $K \cong L$ as fields. Naturally, this isomorphism is a \mathbb{Q} -isomorphism. Since $K|\mathbb{Q}$ is a normal extension, it follows that $K = L$ when viewed as sub-extensions of fixed algebraic closure \mathbb{Q}^{a} . In particular, $G_K = G_L$ as subgroups of $G_{\mathbb{Q}}$. Thus the open normal subgroups of $G_{\mathbb{Q}}$ are *equivariant*, i.e., they are invariant under automorphisms of $G_{\mathbb{Q}}$. This lead NEUKIRCH to the following questions:

- 1) Does $G_{\mathbb{Q}}$ have inner automorphisms only?
- 2) Is every isomorphism $\Phi : G_K \rightarrow G_L$ as above defined by the conjugation by some element inside $G_{\mathbb{Q}}$?

Finally, the first peak in this development was reached at the beginning of the 1970's, with a *positive answer* to Question 1) by IKEDA [Ik] (and partial results by KOMATSU), and the break through by UCHIDA [U1], [U2], [U3] (and unpublished notes by IWASAWA) showing that the answer to Question 2) is positive. Even more, the following holds:

Theorem. *Let K and L be global fields. Then the following hold:*

- (1) *If $G_K \cong G_L$ as profinite groups, $L \cong K$ as fields.*
- (2) *More precisely, for every profinite group isomorphism $\Phi : G_K \rightarrow G_L$, there exists a unique field isomorphism $\phi : L^{\text{s}} \rightarrow K^{\text{s}}$ defining Φ , i.e., such that*

$$\Phi(g) = \phi^{-1} \circ g \circ \phi \quad \text{for all } g \in G_K.$$

In particular, $\phi(L) = K$. And therefore we have a bijection:

$$\text{Isom}_{\text{fields}}(L, K) \cong \text{Out}_{\text{prof.gr.}}(G_K, G_L)$$

This is indeed a very remarkable fact: The Galois theory of the global fields encodes the isomorphism type of such fields in a functorial way! Often this result is called the *Galois characterization of global fields*.

We recall briefly the idea of the proof, as it is very instructive for the future developments. First, recall that by results of TATE and SHAFAREVICH, we know that the virtual ℓ -cohomological dimension $\text{vcd}_{\ell}(K) := \text{vcd}(G_K)$ of a global field K is as follows, see e.g. SERRE [S1], Ch.II:

- i) If K is a number field, then $\text{vcd}_{\ell}(K) = 2$ for all ℓ .
- ii) If $\text{char}(K) = p > 0$, then $\text{vcd}_p(K) = 1$, and $\text{vcd}_{\ell}(K) = 2$ for $\ell \neq p$.

In particular, if $G_K \cong G_L$ then K and L have the same characteristic.

Case 1. $K, L \subset \mathbb{Q}^{\text{a}}$ are number fields. Then the isomorphism $\Phi : G_K \rightarrow G_L$ defines an arithmetical equivalence of K and L . Therefore, K and L have the

same normal hull M_0 over \mathbb{Q} inside \mathbb{Q}^a ; and moreover, for every finite normal sub-extension $M|\mathbb{Q}$ of \mathbb{Q}^a which contains K and L one has: Φ maps G_M isomorphically onto itself, thus defines an isomorphism

$$\overline{\Phi}_M : \text{Gal}(M|K) \rightarrow \text{Gal}(M|L)$$

In order to conclude, one shows for a properly chosen *Abelian* extension $M_1|M$, every isomorphism $\overline{\Phi}_M$ which can be extended to an isomorphism $\overline{\Phi}_{M_1}$, can also be extended to an automorphism of $\text{Gal}(M|\mathbb{Q})$. Finally, one deduces from this that Φ can be extended to an automorphism of $G_{\mathbb{Q}}$, etc.

Note that the fact that the arithmetical equivalence of normal number fields implies their isomorphism relies on the Chebotarev Density Theorem, thus *analytical methods*. Until now we do not have a purely *algebraic proof* of that fact.

Case 2. K, L are global fields over \mathbb{F}_p . First recall that the space of all the non-trivial places $\mathcal{P}(K)$ of K is in a canonical bijection with the closed points of the unique complete smooth model $X \rightarrow \mathbb{F}_p$ of K . In particular, given an isomorphism $\Phi : G_K \rightarrow G_L$, the “arithmetical equivalence” of K and L , is just a bijection $X^0 \rightarrow Y^0$ from the closed points of X to the closed points of the complete smooth model $Y \rightarrow \mathbb{F}_p$ of L . And the problem is now to show that this abstract bijection comes from geometry. The way to do it is by using the class field theory of global function fields as follows: First, one recovers the Frobenius elements at each place \mathfrak{p} of K ; and then the multiplicative group K^\times by using Artin’s reciprocity map; and finally the addition on $K = K^\times \cup \{0\}$. Since the recipe for recovering these objects is invariant under profinite group isomorphisms, it follows that $\Phi : G_K \rightarrow G_L$ defines a group isomorphism $\phi_K : K^\times \rightarrow L^\times$. Finally, one shows that ϕ_K respects the addition, by reducing it to the case $\phi_K(x+1) = \phi_K(x) + 1$. Moreover, by performing this construction for all finite sub-extensions $K_1|K$ of $K^s|K$, and the corresponding finite sub-extensions $L_1|L$ of $L^s|L$, and using the functoriality of the class field theory, one finally gets a field isomorphism $\phi : K^s \rightarrow L^s$ which defines Φ , i.e., $\Phi(g) = \phi \circ g \circ \phi^{-1}$ for all $g \in G_K$.

PART II: Grothendieck’s Anabelian Geometry

The natural context in which the above result appears as a first prominent example is *Grothendieck’s anabelian geometry*, see [G1], [G2]. We will formulate Grothendieck’s anabelian conjectures in a more general context later, after having presented the basic facts about étale fundamental groups. But it is easy and appropriate to formulate here the so called *birational anabelian Conjectures*, which involve only the usual absolute Galois group.

A) Warm-up: Birational anabelian Conjectures

The so called birational anabelian Conjectures place the Results by NEUKIRCH, IKEDA, IWASAWA, UCHIDA et al —at least conjecturally— into a bigger

picture. And in their most naive form, these conjectures assert that the isomorphism type of the absolute Galois group encodes the isomorphism type of a finitely generated infinite field up to a finite purely inseparable extension. Recall that for an arbitrary field K we denote by K^i its maximal purely inseparable extension. Thus if $\text{char}(K) = 0$, then $K^i = K$. Further, we say that two field homomorphisms $\phi, \psi : L \rightarrow K$ differ by an absolute Frobenius twist, if $\psi = \phi \circ \text{Frob}^n$ on L^i for some power Frob^n of the absolute Frobenius Frob .

Birational anabelian Conjectures.

(1) *There exists a group theoretic recipe in order to recover finitely generated fields K from their absolute Galois groups G_K . In particular, if for such fields K and L one has $G_K \cong G_L$, then $K^i \cong L^i$.*

(2) *Moreover, given such fields K and L , one has the following:*

- *Isom-form: Every isomorphism $\Phi : G_K \rightarrow G_L$ is defined by a field isomorphism $\phi : L^a \rightarrow K^a$, and ϕ is unique up to Frobenius twists. In particular, one has $\phi(L^i) = K^i$.*

- *Hom-form: Every open homomorphism $\Phi : G_K \rightarrow G_L$ is defined by a field embedding $\phi : L^a \hookrightarrow K^a$, and ϕ is unique up to Frobenius twists. In particular, one has $\phi(L^i) \subseteq K^i$.*

As in the case of global fields, the Isom-form of the Birational anabelian Conjecture is also called the *Galois characterization of the finitely generated infinite fields*. The main known facts are summarized below:

Theorem.

(1) (See POP [P2], [P3]) *There is a group theoretical recipe by which one can recover in a functorial way finitely generated infinite fields K from their absolute Galois groups G_K .*

Moreover, this recipe works in such a way that it implies the Isom-form of the birational anabelian Conjecture, i.e., every isomorphism $\Phi : G_K \rightarrow G_L$ is defined by an isomorphism $\phi : L^a \rightarrow K^a$, and ϕ is unique up to Frobenius twists.

(2) (See MOCHIZUKI [Mzk3], Theorem B) *The relative Hom-form of the birational anabelian Conjecture is true in characteristic zero, which means the following: Given function fields K and L over \mathbb{Q} , every open $G_{\mathbb{Q}}$ -homomorphism $\Phi : G_K \rightarrow G_L$ is defined by a unique field embedding $\phi : L^a \rightarrow K^a$, which in particular, maps L into K .*

We give here the sketch of the proof of the Isom-form. MOCHIZUKI's Hom-form relies on his proof of the anabelian conjectures for curves over sub- p -adic fields, and we will say some words about that later on in the Lecture.

The main steps of the proof are the following:

The first part of the proof consists in developing a higher dimensional *Local Theory* which, roughly speaking, is a direct generalization of NEUKIRCH’s result above concerning the description of the places of global fields. Nevertheless, there are some difficulties with this generalization, because in higher dimensions the finitely generated fields do not have unique normal (or smooth) complete models. Recall that a model $X \rightarrow \mathbb{Z}$ for such a field K is by definition a separated, integral scheme of finite type over \mathbb{Z} whose function field is K . We will consider only quasi-projective normal models, maybe satisfying some extra conditions, like regular, etc. In particular, if X is a model of K , then the Kronecker dimension $\dim(K)$ of K equals $\dim(X)$ as a scheme. One has:

- K is a global field if and only if every normal model X of K is an open of either $X_K := \text{Spec } \mathcal{O}_K$ if K is a number field, or of the unique complete smooth model $X_K \rightarrow \mathbb{F}_p$ of K , if K is a global function field with $\text{char}(K) = p$. Further, there exists a natural bijection between the *prime Weil divisors* of X_K and the non-archimedean places of K . The basic result by NEUKIRCH [N1] can be interpreted as follows: First let us say that a closed subgroup $Z \subset G_K$ is a *divisorial like subgroup*, if it is isomorphic to a decomposition group $Z_{\mathfrak{q}}$ over some prime \mathfrak{q} of some global field L . Note that the structure of such groups as profinite groups is known, see e.g., JANNSEN–WINGBERG [J–W]. Then the decomposition groups over the places of K are the *maximal divisorial like subgroups* of G_K .

This gives then the group theoretic recipe for describing the prime Weil divisors of X_K in a functorial way.

- In general, i.e., if K is not necessarily a global field, there is a huge variety of normal complete models $X \rightarrow \mathbb{Z}$ of K . In particular, we cannot hope to obtain much information about a single specific model X of K , as in general there is no privileged model for K as in the global field case. (Well, maybe with the exception of arithmetical surfaces, where one could choose the minimal model, but this doesn’t help much...) A way to avoid this is to consider –in a first approximation– the space of *Zariski prime divisors* \mathcal{D}_K of K . This is by definition, the set of all the discrete valuations v of K defined by the Weil prime divisors of all possible normal models $X \rightarrow \mathbb{Z}$ of K .

A Zariski prime divisor v is called *geometrical* if $\text{char}(K) = \text{char}(Kv)$, or equivalently, if v is trivial on the prime field of K , and *arithmetical* otherwise. Clearly, arithmetical Zariski prime divisors exist only if $\text{char}(K) = 0$. If so, and if v is defined by a Weil prime divisor X_1 of some normal model $X \rightarrow \mathbb{Z}$, then v is *geometric* if and only if v is a “horizontal” divisor of $X \rightarrow \mathbb{Z}$. We denote by \mathcal{D}_K^1 the space of all geometrical Zariski prime divisors of K .

For every Zariski prime divisor $v \in \mathcal{D}_K$ of K , let Z_v be the decomposition group of some prolongation v^s of v to K^s . We will call the totality of all the closed subgroups of the form Z_v the *divisorial subgroups* of G_K or of K . Finally, as above, a closed subgroup $Z \subset G_K$ is called *divisorial like subgroup*, if it is isomorphic to

a divisorial subgroup of a finitely generated field L with $\dim(L) = \dim(K)$. The main results of the Local Theory are as follows, see [P1]:

a) For a Zariski prime divisor v , the numerical data $\text{char}(K)$, $\text{char}(Kv)$, and $\dim(K)$ are group theoretically encoded in Z_v ; in particular, whether v is geometric or not. Further, the inertia group $T_v \subset Z_v$ of $v^s|v$, and the canonical projection $\pi_v : Z_v \rightarrow G_{Kv}$ are also encoded group theoretically in Z_v . In particular, the residual absolute Galois group G_{Kv} at all the Zariski prime divisors v of K is group theoretically encoded in G_K .

b) Every divisorial like subgroup $Z \subset G_K$ is contained in a unique divisorial subgroup Z_v of G_K . Thus the divisorial like subgroups of G_K are exactly the *maximal divisorial like subgroups* of G_K . And the space \mathcal{D}_K is in bijection with the conjugacy classes of divisorial subgroups of G_K .

The results from the local theory above suggest that one should try to prove the birational anabelian Conjecture by *induction on* $\dim(K)$. This is the idea for developing a *Global Theory* along the following lines:

First, the Isom-form of the birational anabelian Conjecture for global fields, i.e., $\dim(K) = 1$, is known; and we think of it as the first induction step. Now suppose that $\dim(K) = d > 1$. By the induction hypothesis, suppose that the Isom-form of the birational anabelian Conjecture is true in dimension $< d$. Then one recovers the field K^1 up to Frobenius twists from G_K along the following steps (and from this recipe it will be clear, what we do mean by a “group theoretic recipe”).

Step 1) Recover the cyclotomic character $\chi_K : G_K \rightarrow \widehat{\mathbb{Z}}^\times$ of G_K .

The recipe is as follows: Since $\dim(Kv) = \dim(K) - 1 < d$, the cyclotomic character χ_{Kv} is “known” for each geometric Zariski prime divisor $v \in \mathcal{D}_K^1$. Therefore, the cyclotomic character

$$\chi_v : Z_v \xrightarrow{\pi_v} G_{Kv} \xrightarrow{\chi_{Kv}} \widehat{\mathbb{Z}}^\times$$

is known for all $v \in \mathcal{D}_K^1$. On the other hand, using the higher dimensional Chebotarev Density Theorem, see e.g., SERRE [S3], it follows that $\ker(\chi_K)$ is the closed subgroup of G_K generated by all the $\ker(\chi_v)$, $v \in \mathcal{D}_K^1$. Thus χ_K is the unique character $\chi : G_K \rightarrow \widehat{\mathbb{Z}}^\times$ which coincides with χ_v on each Z_v .

Next let $\mathbb{T}_K = \varprojlim_m \mu_m$ be the Tate module of K . Denote by $\widehat{\mathbb{Z}}'(1)$ the adic completion of \mathbb{Z} with respect to all integers m relatively prime to $\text{char}(K)$, and fix an identification ι of these two G_K -modules. The Kummer Theory gives a functorial completion homomorphism as follows:

$$K^\times \xrightarrow{J_K} \widehat{K} \xrightarrow{\delta} \mathbf{H}^1(K, \widehat{\mathbb{Z}}'(1)),$$

where “functorial” means that performing this construction for finite extensions $M|K$ inside $K^a|K$, we get corresponding commutative “inclusion-restriction” diagrams (which we omit to write down here). An essential point to make here is that the completion morphism $j_K : K^\times \rightarrow \widehat{K}$ is injective. This follows by induction on $\dim(K)$ by using the following fact: Let $K|k$ be the function field of a geometrically irreducible complete curve $X \rightarrow k$. Then K^\times/k^\times is the group of principal divisors of X , thus a free Abelian group. And for K a number field, one knows that K^\times/μ_K is a free Abelian group.

Step 2) Recover the geometric small sets of Zariski prime divisors.

We will say that a subset $\mathcal{D} \subset \mathcal{D}_K^1$ of Zariski prime divisors is *geometric*, if there exists a quasi-projective normal model $X \rightarrow k$ of K such that $\mathcal{D} = \mathcal{D}_X$ is the set of Zariski prime divisors of K defined by the Weil prime divisors of X . Here, k is the field of constants of K . It is a quite technical point to show —by induction on $d = \text{td}(K)$, that the geometric sets of Zariski prime divisors can be recovered from G_K , see POP [P3]. Next let $\mathcal{D} = \mathcal{D}_X$ be a geometric set of Zariski prime divisors. One has a canonical exact sequence

$$1 \rightarrow U_{\mathcal{D}} \rightarrow K^\times \rightarrow \text{Div}(X) \rightarrow \mathfrak{Cl}(X) \rightarrow 0,$$

where $U_{\mathcal{D}}$ are the units in the ring of global sections on X , and the other notations are standard. Since the base field k is either finite or a number field, the Weil divisor class group $\mathfrak{Cl}(X)$ is finitely generated. Thus if X is “sufficiently small”, then $\mathfrak{Cl}(X) = 0$. A geometric set of Zariski prime divisors $\mathcal{D} = \mathcal{D}_X$ will be called a *small geometric set* of Zariski prime divisors, if the adic completion $\widehat{\mathfrak{Cl}}(X)$ is trivial. One shows that the small geometric sets of Zariski prime divisors can be recovered from G_K , see loc.cit. In this process, one shows that the adic completion of the above exact sequence can be recovered from G_K too:

$$1 \rightarrow \widehat{U}_{\mathcal{D}} \rightarrow \widehat{K}^\times \rightarrow \widehat{\text{Div}}(X) = \widehat{\bigoplus}_v \widehat{\mathbb{Z}}' \rightarrow \widehat{\mathfrak{Cl}}(X) \rightarrow 0,$$

Step 3) Recover the multiplicative group K^\times inside \widehat{K} .

Let $\mathcal{D} = \mathcal{D}_X$ be a small geometric set of Zariski prime divisors of K . The resulting exact sequence defined above becomes $1 \rightarrow \widehat{U}_{\mathcal{D}} \rightarrow \widehat{K} \rightarrow \widehat{\bigoplus}_v \widehat{\mathbb{Z}}' \rightarrow 0$, as $\widehat{\mathfrak{Cl}}(X) = 0$. Next let $v \in \mathcal{D}$ be arbitrary. Then the group of global units $U_{\mathcal{D}}$ is contained in the group of v -units \mathcal{O}_v^\times . Thus the $(\text{mod } \mathfrak{m}_v)$ reduction homomorphism $p_v : \mathcal{O}_v^\times \rightarrow K v^\times$ is defined on $U_{\mathcal{D}}$. Using some arguments involving Hilbertian fields, one shows that there exist “many” $v \in \mathcal{D}$ such that $U_{\mathcal{D}}$ as well as $\widehat{U}_{\mathcal{D}}$ are actually mapped isomorphically into $K v^\times$, respectively $\widehat{K} v$; and moreover, that inside $\widehat{K} v$ one has

$$(*) \quad p_v(U_{\mathcal{D}}) = \widehat{p}_v(\widehat{U}_{\mathcal{D}}) \cap K v^\times.$$

On the Galois theoretic side, the reduction map p_v is defined by the restriction coming from the inclusion $Z_v \hookrightarrow G_K$. And moreover, since $U_{\mathcal{D}}$ is contained in the

v -units, it follows that under the restriction map $\widehat{K} \rightarrow H^1(Z_v, \widehat{\mathbb{Z}}'(1))$, the image of $\widehat{U}_{\mathcal{D}}$ is contained in the image of the inflation map $\text{inf}(\pi_v) : \widehat{K}v \rightarrow H^1(Z_v, \widehat{\mathbb{Z}}'(1))$ defined by the canonical projection $\pi_v : Z_v \rightarrow G_{Kv}$.

Finally, the recipe to recover K^\times inside \widehat{K} is as follows. First, for each small geometric set of Zariski prime divisors \mathcal{D} and v as above, $U_{\mathcal{D}}$ is exactly the preimage of $\hat{p}_v(\widehat{U}_{\mathcal{D}}) \cap Kv^\times$, by assertion $(*)$ above. Since $K^\times = \cup_{\mathcal{D}} U_{\mathcal{D}}$, when \mathcal{D} runs over smaller and smaller (small) geometric sets of Zariski prime divisors, we finally recover K^\times inside \widehat{K} .

Step 4) Define the addition in $K = K^\times \cup \{0\}$.

This is easily done using the induction hypothesis: Let namely $x, y \in K^\times$ be given. Then $x + y = 0$ iff $x/y = -1$, and this fact is encoded in K^\times . Now suppose that $x + y \neq 0$. Then $x + y = z$ in K iff for all v such that x, y, z are all v -units one has: $p_v(x) + p_v(y) = p_v(z)$. On the other hand, this last fact is encoded in the field structure of G_{Kv} , which we already know.

Finally, in order to conclude the proof of the Isom-form of the birational anabelian Conjecture, we proceed as follows: Let $\Phi : G_K \rightarrow G_L$ be an isomorphism of absolute Galois group $\Phi : G_K \rightarrow G_L$. Then the recipe of recovering the fields K and L are “identified” via Φ , and shows that the p -divisible hulls of K and L inside $\widehat{K} \cong \widehat{L}$ must be the same, where $p = \text{char}(K)$. This finally leads to an isomorphism $\phi : L^a \rightarrow K^a$ which defines Φ . Its uniqueness up to Frobenius twists follows from the fact that given two such field isomorphisms ϕ', ϕ'' , then setting $\phi := \phi'^{-1} \circ \phi''$ we obtain an automorphism of K^a which commutes with G_K . And one checks that any such automorphism is a Frobenius twist.

B) Anabelian Conjectures for Curves

a) Étale fundamental groups

Let X be a connected scheme endowed with a geometric base point \bar{x} . Recall that the étale fundamental group $\pi_1(X, \bar{x})$ of (X, \bar{x}) is the automorphism group of the fiber functor on the category of all the étale connected covers of X . The étale fundamental group is functorial in the following sense: Let connected schemes with geometric base points (X, \bar{x}) and (Y, \bar{y}) , and a morphism $\phi : X \rightarrow Y$ be given such that $\bar{y} = \phi \circ \bar{x}$. Then ϕ gives rise to a morphism between the fiber functors $\mathcal{F}_{\bar{x}}$ and $\mathcal{F}_{\bar{y}}$, which induces a continuous morphism of profinite groups $\pi_1(\phi) : \pi_1(X, \bar{x}) \rightarrow \pi_1(Y, \bar{y})$ in the canonical way. In particular, setting $Y = X$ and \bar{y} some geometric point of X , a “path” from between \bar{x} and \bar{y} , gives rise to an inner automorphism of $\pi_1(X, \bar{x})$. In other words, $\pi_1(X, \bar{x})$ is *determined by X up to inner automorphisms*. (This means that the situation is completely parallel to the one in the case of the topological fundamental group.) It is one of the basic properties of the étale fundamental group that it is invariant under *radicial morphisms*, in particular under purely inseparable covers and Frobenius twists.

Next let \mathcal{G} be the category of all profinite groups and outer continuous homomorphisms as morphisms. The objects of \mathcal{G} are the profinite groups, and for given objects G and H , a \mathcal{G} -morphism from G to H is a set of the form $\text{Inn}(H) \circ f$, where $f : G \rightarrow H$ is a morphism of profinite groups, and $\text{Inn}(H)$ is the set of all the inner automorphisms of H . Clearly, if $f, g : G \rightarrow H$ differ by an inner automorphism of G , then $\text{Inn}(H) \circ f = \text{Inn}(H) \circ g$, thus they define the same \mathcal{G} -homomorphism from G to H . Further, $\text{Inn}(H) \circ f$ is a \mathcal{G} -isomorphism if and only if $f : G \rightarrow H$ is an isomorphism of profinite groups.

Therefore, viewing the étale fundamental group π_1 as having values in \mathcal{G} rather than in the category of profinite groups, the relevance of the geometric points \bar{x} vanishes. Therefore, we will simply write $\pi_1(X)$ for the fundamental group of a connected scheme X .

In the same way, if S is a connected base scheme, and X is a connected S -scheme, then the structure morphism $\varphi_X : X \rightarrow S$ gives rise to an *augmentation morphism* $p_X : \pi_1(X) \rightarrow \pi_1(S)$. Thus the category \mathfrak{Sch}_S of all the S -schemes is mapped by π_1 into the category \mathcal{G}_S of all the $\pi_1(S)$ -groups, i.e. the profinite groups G with an “augmentation” morphism $pr_G : G \rightarrow \pi_1(S)$.

Now let us consider the more specific situation when the base scheme S is a field, and the k -schemes X are geometrically connected. Denote $\bar{X} = X \times_k k^s$ the base to the separable closure of k (in some fixed “universal field”), and remark that by the facts above one has an exact sequence of profinite groups of the form

$$1 \rightarrow \pi_1(\bar{X}) \rightarrow \pi_1(X) \rightarrow G_k \rightarrow 1.$$

In particular, we have a representation $\rho_X : G_k \rightarrow \text{Out}(\pi_1(\bar{X})) = \text{Aut}_{\mathcal{G}}(\pi_1(\bar{X}))$ which encodes most of the information carried by the exact sequence above. The group $\pi_1(\bar{X})$ is called the *algebraic* (or *geometric*) fundamental group of X . In general, little is known about $\pi_1(\bar{X})$, and in particular, even less about $\pi_1(X)$. Nevertheless, if X is a k -variety, and $k \subset \mathbb{C}$, then the base change to \mathbb{C} gives a realization of $\pi_1(\bar{X})$ as the *profinite completion* of the topological fundamental group of $X^{\text{an}} = X(\mathbb{C})$.

In terms of function fields, if $X \rightarrow k$ is geometrically integral, one has the following: Let $k(X) \hookrightarrow k(\bar{X})$ be the function fields of $\bar{X} \rightarrow X$. Then the algebraic fundamental group $\pi_1(\bar{X})$ is (canonically) isomorphic to the Galois group of a maximal unramified Galois field extension $\mathcal{K}_{\bar{X}} | k(\bar{X})$.

Finally, we recall that $\pi_1(X)$ is a *birational invariant* in the case X is complete and regular. In other words, if X and X' are birationally equivalent complete regular k -varieties, then $\pi_1(X) \cong \pi_1(X')$ and $\pi_1(\bar{X}) \cong \pi_1(\bar{X}')$ canonically.

b) *Étale fundamental groups of curves*

Specializing even more, we turn our attention to curves, and give a short review of the basic known facts in this case. In this discussion we will suppose

that X is a *smooth connected curve*, having a *smooth completion* say X_0 over k . We denote $S = X_0 \setminus X$, and $\bar{S} = \bar{X}_0 \setminus \bar{X}$. We will say that X is a (g, r) curve, if X_0 has (geometric) genus g , and $|\bar{S}| = r$. We will say that X is a *hyperbolic curve*, if its Euler–Poincaré characteristic $2 - 2g - r$ is negative. And we will say that a curve X as above is *virtually hyperbolic*, if it has an étale connected cover $X' \rightarrow X$ such that X' is a *hyperbolic curve* in the sense above. (Note that every étale cover $f : X' \rightarrow X$ as above is smooth and has a smooth completion which is a (g', r') -curve with $g \leq g'$ and $r' \leq r \deg(f)$ over some finite $k'|k$.)

In the above notations, let $X \rightarrow k$ be a (g, r) curve. Then a short list of the known facts about the algebraic fundamental group $\pi_1(\bar{X})$ is as follows. First, let $\Gamma_{g,r}$ be the fundamental group of the orientable compact topological surface of genus g with r punctures. Thus

$$\Gamma_{g,r} = \langle a_1, b_1, \dots, a_g, b_g, c_1, \dots, c_r \mid \prod_i [a_i, b_i] \prod c_j = 1 \rangle$$

is the discrete group on $2g + r$ generators $a_1, b_1, \dots, a_g, b_g, c_1, \dots, c_r$ with the given unique relation. (The generators a_i, b_i, c_j have a precise interpretation as loops around the handles, respectively around the missing points.) In particular, if $r > 0$, then $\Gamma_{g,r}$ is the discrete free group on $2g + r - 1$ generators. It is well known that $\Gamma_{g,r}$ is *residually finite*, i.e., $\Gamma_{g,r}$ injects into its profinite completion:

$$\Gamma_{g,r} \hookrightarrow \widehat{\Gamma}_{g,r}.$$

Finally, given a fixed prime number p , respectively arbitrary prime numbers ℓ , we will denote by $\widehat{\Gamma}_{g,r} \rightarrow \widehat{\Gamma}'_{g,r}$ the maximal prime- p quotient of $\widehat{\Gamma}_{g,r}$, and by $\widehat{\Gamma}_{g,r} \rightarrow \widehat{\Gamma}^\ell_{g,r}$ the maximal pro- ℓ quotient of $\Gamma_{g,r}$.

Case 1. $\text{char}(k) = 0$.

Using the remark above, in the case $k \hookrightarrow \mathbb{C}$, it follows that $\pi_1(\bar{X}) \cong \widehat{\Gamma}_{g,r}$ via the base change $X \times_k \mathbb{C} \rightarrow \bar{X}$. If $\kappa(\bar{X})$ is the function field of \bar{X} , then $\pi_1(\bar{X})$ is the Galois group of a maximal Galois unramified field extension $\mathcal{K}_{k(\bar{X})} | k(\bar{X})$. Moreover, the loops $c_j \in \Gamma_{g,r}$ around the missing points $x_i \in \bar{X}_0 \setminus \bar{X}$ are canonical generators of inertia groups T_{x_i} over these points in $\pi_1(\bar{X})$. In particular we have:

a) X is a complete curve of genus g if and only if $\pi_1(\bar{X})$ has $2g$ generators a_i, b_i with the single relation $\prod_i [a_i, b_i] = 1$, provided X is not \mathbb{A}_k^1 .

b) X is of type (g, r) with $r > 0$ if and only if $\pi_1(\bar{X})$ is a profinite free group on $2g + r - 1$ generators, provided X is not k -isomorphic to \mathbb{A}_k^1 or \mathbb{P}_k^1 .

Clearly, the dichotomy between the above subcases a) and b) can be as well deduced from the pro- ℓ maximal quotient $\pi_1^\ell(\bar{X})$ of $\pi_1(\bar{X})$, by simply replacing “profinite” by “pro- ℓ ”.

Further, the following conditions on X are equivalent:

(i) X is hyperbolic

(ii) X is virtually hyperbolic.

(iii) $\pi_1(\overline{X})$ is non-Abelian, or equivalently, (iii)^ℓ $\pi_1^\ell(\overline{X})$ is non-Abelian.

Case 2. $\text{char}(k) > 0$.

First recall that the *tame fundamental group* $\pi_1^t(X)$ of X is the maximal quotient of $\pi_1(X)$ which classifies étale connected covers $X' \rightarrow X$ whose ramification above the missing points $x_i \in \overline{X}_0 \setminus \overline{X}$ is tame. We will denote by $\pi_1^t(\overline{X})$ the tame quotient of $\pi_1(\overline{X})$, and call it the *tame algebraic fundamental group* of X . Now the main technical tools used in understanding $\pi_1(\overline{X})$ and its tame quotient $\pi_1^t(\overline{X})$ are the following two facts:

Shafarevich’s Theorem.

In the context above, set $\text{char}(k) = p > 0$, and denote by $\pi_1^p(\overline{X})$ the maximal pro- p quotient of $\pi_1(\overline{X})$. Further let $r_{X_0} = \dim_{\mathbb{F}_p} \text{Jac}_{X_0}[p]$ denote the Hasse–Witt invariant of the complete curve \overline{X}_0 . Then one has:

- (1) If $X = X_0$, then $\pi_1^p(\overline{X})$ is a pro- p free group on $r_{X_0} \leq g$ generators.
- (2) If X is affine, then $\pi_1^p(X)$ is a pro- p free group on $|k^a|$ generators.

Let k be an arbitrary base field, and v a complete discrete valuation with valuation ring $R = R_v$ of k and residue field $kv = \kappa$. Let $X \rightarrow k$ be a smooth curve which has a smooth completion $X_0 \rightarrow k$. We will say that $X \rightarrow k$ has *good reduction* at v , if the following hold: $X_0 \rightarrow k$ has a smooth model $X_{0,R} \rightarrow R$ over R , and there exists an étale divisor $S_R \rightarrow R$ of $X_{0,R}$ such that the generic fiber of the complement $X_{0,R} \setminus S_R =: X_R \rightarrow R$ is X .

Now let $X \rightarrow k$ be a hyperbolic curve having good reduction at v . In the notations from above, let $X_s \rightarrow \kappa$ be the special fiber of $X_R \rightarrow R$. Then the canonical diagram of schemes

$$\begin{array}{ccccc} X & \hookrightarrow & X_R & \hookrightarrow & X_s \\ \downarrow & & \downarrow & & \downarrow \\ k & \hookrightarrow & R & \hookrightarrow & \kappa \end{array}$$

gives rise to a diagram of fundamental groups as follows:

$$\begin{array}{ccccc} \pi_1^t(X) & \rightarrow & \pi_1^t(X_R) & \leftarrow & \pi_1^t(X_s) \\ \downarrow & & \downarrow & & \downarrow \\ G_k & \rightarrow & G_k^t & \leftarrow & G_\kappa \end{array}$$

where $\pi_1^t(X_R)$ is the “tame fundamental group” of X_R , i.e., the maximal quotient of $\pi_1(X)$ classifying connected covers of X_R which have ramification only along S_R and the generic point of the special fiber, and this ramification is tame, and G_k^t is the Galois group of the maximal tamely ramified extension of k . The fundamental result concerning the fundamental groups in the diagram above is the following:

Grothendieck's Specialization Theorem.

In the context above, let X be a smooth curve of type (g, r) . Further let R^t be the extension of R to k^t , and $\overline{X}_R := X_R \times_R R^t$. Then one has:

(1) $\pi_1^t(\overline{X}) \rightarrow \pi_1^t(\overline{X}_R)$ is surjective, and $\pi_1^t(\overline{X}_R) \leftarrow \pi_1^t(\overline{X}_s)$ is an isomorphism. The resulting surjective homomorphism

$$\mathrm{sp}_v : \pi_1^t(\overline{X}) \rightarrow \pi_1^t(\overline{X}_s)$$

is called the specialization homomorphism of tame fundamental groups at v . In particular, $\pi_1^t(\overline{X})$ is a quotient of $\widehat{\Gamma}_{g,r}$ in such a way that the generators c_j are mapped to inertia elements at the missing points $x_i \in \overline{X}_0 \setminus \overline{X}$.

(2) Further, let $\mathrm{char}(k) = p$, and denote by $\pi_1'(\overline{X})$ the maximal prime to p quotient of $\pi_1(\overline{X})$ (which then equals the maximal prime to p quotient of $\pi_1^t(\overline{X})$ too). Then $\pi_1'(\overline{X}) \cong \widehat{\Gamma}'_{g,r}$, and sp_v maps $\pi_1'(\overline{X})$ isomorphically onto $\pi_1'(\overline{X}_s)$. In particular, $\pi_1'(\overline{X})$ depends on (g, r) only.

Combining Shafarevich's Theorem and Grothendieck's Specialization Theorem above, we immediately see that the following facts and invariants of $X \rightarrow k$ are encoded in $\pi_1(\overline{X})$:

a) If $\ell \neq p$, then $\pi_1^\ell(\overline{X}) \cong \widehat{\Gamma}_{g,r}^\ell$, and $\pi_1^p(\overline{X}) \not\cong \widehat{\Gamma}_{g,r}^p$. Therefore, $p = \mathrm{char}(k)$ can be recovered from $\pi_1(\overline{X})$, provided X is not \mathbb{P}_k^1 .

a)^t The same is true correspondingly concerning the tame fundamental group $\pi_1^t(\overline{X})$, provided X is not isomorphic to \mathbb{A}_k^1 of \mathbb{P}_k^1 .

b) $X \rightarrow k$ is complete if and only if $\pi_1^p(\overline{X})$ is finitely generated.

b)^t Correspondingly, $X \rightarrow k$ is complete if and only if $\pi_1^\ell(\overline{X})$ not pro- ℓ -free, provided X is not isomorphic to \mathbb{A}_k^1 .

c) In particular, if X is complete, then $\pi_1^\ell(\overline{X})$ has $2g$ generators, thus g can be recovered from $\pi_1^\ell(\overline{X})$.

Finally, concerning the virtual hyperbolicity of X we have the following:

d) Every affine curve X is virtually hyperbolic.

Remark.

Clearly, the applicability of Grothendieck's Specialization Theorem is limited by the fact that one would need *a priori* criterions for the good reduction of the given curve X at the (completions of k with respect to the) discrete valuations v of k . At least in the case of hyperbolic curves $X \rightarrow k$ such criteria do exist. The setting is as follows: Let $X \rightarrow k$ be a hyperbolic curve, and let v be a discrete valuation of k . Let $T_v \subseteq Z_v$ be the inertia, respectively the decomposition, groups of some prolongation of v to k^s . Recall the canonical projections $\pi_1(X) \rightarrow G_k$ and $\pi_1^t(X) \rightarrow G_k$ and the resulting Galois representations $\rho_X : G_k \rightarrow \mathrm{Out}(\pi_1(\overline{X}))$ and $\rho_X^t : G_k \rightarrow \mathrm{Out}(\pi_1^t(\overline{X}))$. Then one can characterize the fact that X has (potentially) good reduction at v as follows, see ODA [O] in the case of complete

hyperbolic curves, and by TAMAGAWA [T1] in the case of arbitrary hyperbolic curves:

In the above notations, $X \rightarrow k$ has good reduction at v if and only if the representation ρ_X^t is trivial on T_v .

The concrete picture of how to apply the above remark in studying fundamental groups of hyperbolic curves $X \rightarrow k$ over either finitely generated infinite base field k or finitely generated fields over some fixed base field k_0 is as follows: Let $X \rightarrow k$ be a smooth curve of type (g, r) . Further let S be a smooth model of k over \mathbb{Z} , if k is a finitely generated field, respectively over the base k_0 otherwise. For every closed point $s \in S$, there exists a discrete valuation v_s whose valuation ring R_s dominates the local ring $\mathcal{O}_{S,s}$, and having residue field $\kappa_{v_s} = \kappa(s)$. Let us choose such a valuation v_s . Then in the previous notations, X has good reduction at s if and only if ρ_X^t is trivial on the inertia group T_s over the point s . Note that by the uniqueness of the smooth model $X_{R_s} \rightarrow R_s$ —in the case it does exist, the existence of such a good reduction *does not depend on the concrete valuation v_s* used. One should also remark here, that in the context above, $X \rightarrow k$ has good reduction on a Zariski open subset of S . This follows e.g., from the Jacobian Criterion for smoothness.

Finally, we now come to announcing Grothendieck’s anabelian Conjectures for Curves and the Section Conjectures.

Let \mathcal{P} be a property defined for some category of schemes X . We will say that the property \mathcal{P} is an *anabelian property*, if it is encoded in $\pi_1(X)$ in a group theoretical way, or in other words, if \mathcal{P} can be recovered by a group theoretic recipe from $\pi_1(X)$. In particular, if X has the property \mathcal{P} , and $\pi_1(X) \cong \pi_1(Y)$, then Y has the property \mathcal{P} .

Examples:

a) In the category of all the fields K , the property “ K is real closed” is anabelian. This is the Theorem of Artin–Schreier from above.

b) In the category of all the smooth k -curves X which are not isomorphic to \mathbb{A}_k^1 , the property: “ X is complete and has genus g ” is anabelian. This follows from the structure theorems for the fundamental group of complete curves as discussed above.

We will say that a scheme X is *anabelian* if the isomorphy type of X up to some natural transformations, which are not encoded in Galois Theory, can be recovered *group theoretically* from $\pi_1(X)$ in a functorial way; or equivalently, if there exists a *group theoretic recipe* to recover the isomorphy type of X , up to the natural transformations in discussion, from $\pi_1(X)$. Typical examples of such “natural transformations” which are not seen by Galois Theory are the radicial

covers and the birational equivalence of complete regular schemes. Concretely, for k -varieties $X \rightarrow k$ with $\text{char}(k) = p > 0$, there are two typical radicial covers: First the maximal purely inseparable cover $X^i \rightarrow k^i$. And second, the Frobenius twists $X(n) \rightarrow k$ of $X \rightarrow k$ and/or $X^i(n) \rightarrow k^i$ of $X^i \rightarrow k^i$ obtained by acting by Frob^n “on the coefficients”: $X(n) := X \times_{\text{Frob}^n} k \rightarrow k$. In the same way, if $X \rightarrow k$ and $Y \rightarrow k$ are complete regular k varieties which are birationally equivalent, then $\pi_1(X)$ and $\pi_1(Y)$ are canonically isomorphic, but X and Y might be very different.

A good set of examples of anabelian schemes are the *finitely generated infinite fields*, as we have seen in the previous section. Given such a field K , one has $\pi_1(\text{Spec } K) = G_K$, and by the birational anabelian Conjectures we know that K can be recovered from $\pi_1(K)$ in a functorial way, up to pure inseparable extensions and Frobenius twists.

Anabelian Conjecture for Curves (absolute form)

(1) *Let $X \rightarrow k$ be a virtually hyperbolic curve over a finitely generated base field k . Then X is anabelian in the sense that the isomorphism type of X can be recovered from $\pi_1(X)$ up to pure inseparable covers and Frobenius twists.*

(2) *Moreover, given such curves $X \rightarrow k$ and $Y \rightarrow l$, one has the following:*

- *Isom-form: Every isomorphism $\Phi : \pi_1(X) \rightarrow \pi_1(Y)$ is defined by an isomorphism $\phi : X^i \rightarrow Y^i$, and ϕ is unique up to Frobenius twists.*

- *Hom-form: Every open homomorphism $\Phi : \pi_1(X) \rightarrow \pi_1(Y)$ is defined by a dominant morphism $\phi : X^i \rightarrow Y^i$, and ϕ is unique up to Frobenius twists.*

One could as well consider a relative form of the above conjecture as follows:

Anabelian Conjecture for Curves (relative form)

(1) *Let $X \rightarrow k$ be a virtually hyperbolic curve over a finitely generated base field k . Then $X \rightarrow k$ is anabelian in the sense that $X \rightarrow k$ can be recovered from $\pi_1(X) \rightarrow G_k$ up to pure inseparable covers and Frobenius twists.*

(2) *Moreover, given such curves $X \rightarrow k$ and $Y \rightarrow k$, one has the following:*

- *Isom-form: Every G_k -isomorphism $\Phi : \pi_1(X) \rightarrow \pi_1(Y)$ is defined by a unique k^i -isomorphism $\phi : X^i(n) \rightarrow Y^i$ for some n -twist.*

- *Hom-form: Every open G_k -homomorphism $\Phi : \pi_1(X) \rightarrow \pi_1(Y)$ is defined by a unique dominant k^i -morphism $\phi : X^i(n) \rightarrow Y^i$ of some n -twist.*

C) The Section Conjectures

Let $X_0 \rightarrow k$ be an arbitrary irreducible k -variety, and $X \subset X_0$ an open k -subvariety. Let $x \in X_0$ be a regular k^i -rational point of X_0 . Then choosing a system of regular parameters (t_1, \dots, t_d) at x , we can construct —by the standard procedure— a valuation v_x of the function field $k(X)$ of X with value group

$v_x(K^\times) = \mathbb{Z}^d$ ordered lexicographically, and residue field $k(X)v_x = \kappa(x)$, thus a subfield of k^i . Let v be a prolongation of v_x to $k(X)^s$, and let $T_v \subset Z_v$ be the inertia, respectively decomposition group of $v|v_x$ in G_K . By general valuation theory, see e.g., KUHLMANN–PANK–ROQUETTE [K–P–R], one has: T_v has complements G_v in Z_v . And clearly, since $k(X)v_x = \kappa(x) \subset k^i$, under the canonical exact sequence

$$1 \rightarrow T_v \rightarrow Z_v \rightarrow G_{k(X)v_x} = G_k \rightarrow 1,$$

every complement G_v is mapped isomorphically onto $G_k = G_{k^i}$. Therefore, the canonical projection

$$(*) \quad pr_{k(X)} : G_{k(X)} \rightarrow G_k$$

has sections $s_v : G_k \rightarrow G_v \subset G_{k(X)}$ constructed as shown above.

Moreover, let us recall that under the canonical projection $G_{k(X)} \rightarrow \pi_1(X)$, the decomposition group Z_v is mapped onto the decomposition group Z_x of v in $\pi_1(X)$, and T_v is mapped onto the inertia group T_x of v in $\pi_1(X)$. And finally, any complement G_v of T_v is mapped isomorphically onto a complement G_x of T_x in Z_x . Clearly $G_x \rightarrow G_k$ isomorphically, thus

$$(**) \quad pr_X : \pi_1(X) \rightarrow G_k$$

has a section $s_x : G_k \rightarrow G_x \subset \pi_1(X)$ defined via the k^i -rational point $x \in X(k^i)$. Moreover, I think it's instructive to remark that one has to distinct cases:

a) Suppose that $x \in X$. Then $T_x = \{1\}$, as the étale covers of X are not ramified over x . Therefore, $Z_x = G_x$. And in this case the sections of pr_X of the form above build a full conjugacy class of sections.

b) Next let $x \in (X_0 \setminus X)(k^i)$. Then $T_x \neq \{1\}$ and $G_x \neq Z_x$ in general. Therefore, for a given k^i -rational point x , there might exist several complements G_x of T_x in Z_x , thus sections of $pr_X : \pi_1(X) \rightarrow G_k$, which are not conjugate inside $\pi_1(X)$. Such sections are called *sections at infinity* (for the variety X). In the case of an arbitrary variety $X \rightarrow k$ it is not known how to classify the conjugacy classes of such sections. But if $X \rightarrow k$ is a curve, and $\text{char}(k) = 0$, then these sections are classified by $H^1(G_k, \widehat{\mathbb{Z}}(1))$.

Section Conjectures.

Let $X \rightarrow k$ be a hyperbolic curve over a finitely generated infinite field k , and in the case $\text{char}(k) > 0$, suppose that X has no finite covers which are defined over a finite field. In the notations from above, the following hold:

(1) Birational form: The sections of $pr_{k(X)}$ arise from k^i -rational points of X_0 as indicated above.

(2) Curve form: The sections of pr_X arise from k^i -rational points of X_0 of X as indicated above.

Concerning **Higher dimensional anabelian Conjectures**, there are only vague ideas. There are some obvious necessary conditions which higher dimensional varieties X have to satisfy in order to be anabelian (like being of general type, being $K(\pi_1)$, etc.). Also, easy counter-examples show that one cannot expect a naive Hom-form of the conjectures. See GROTHENDIECK [G2], and IHARA–NAKAMURA [I–N], MOCHIZUKI [Mzk3], [Mzk4] for more about this.

Remark (Standard reduction technique).

Before going into the details concerning the known facts about the anabelian Conjectures for curves, let us set the technical frame for a fact used several times below. Let $X \rightarrow k$ be a smooth curve over the field k . Suppose that k is either a finitely generated infinite field, or a function field over some base field k_0 . Let $S \rightarrow \mathbb{Z}$, respectively $S \rightarrow k_0$ be a smooth model of k .

Next let $X \rightarrow k$ be a hyperbolic curve, say with smooth completion X_0 . Let $\pi_1(\overline{X}) \rightarrow G_k$, respectively $\pi_1^\dagger(\overline{X}) \rightarrow G_k$ be the corresponding canonical projections. Then choosing for each closed point $s \in S$ a discrete valuation v_s which dominates the local ring of s , we have the Oda–Tamagawa Criterion (mentioned above) for deciding whether $X \rightarrow k$ has good reduction at s . We also know, that $X \rightarrow k$ has good reduction on a Zariski open subset of S . In particular, the Oda–Tamagawa Criterion is a group theoretic criterion for describing the Zariski open subset of S on which $X \rightarrow k$ has good reduction. Moreover, if s is a point of good reduction of $X \rightarrow k$, then Grothendieck’s Specialization Theorem for π_1^\dagger gives a commutative diagram of the following form:

$$\begin{array}{ccc} \pi_1^\dagger(X_{k_v}) & \xrightarrow{\text{sp}_s} & \pi_1^\dagger(X_s) \\ \downarrow & & \downarrow \\ Z_v & \xrightarrow{\text{pr}_v} & G_{\kappa(s)} \end{array}$$

where k_v is the fixed field of Z_v inside k^s , thus it is a decomposition field over v . Therefore, *from the data $\pi_1^\dagger(X) \rightarrow G_k$, one recovers in a canonical way its special fibers $\pi_1^\dagger(X_s) \rightarrow G_{\kappa(s)}$ at all the points s where X has good reduction.*

I) Tamagawa’s Results concerning affine hyperbolic curves

In this subsection we will sketch a proof of the following result by AKIO TAMAGAWA concerning affine hyperbolic curves.

Theorem. (See TAMAGAWA [T1])

(1) *There exists a group theoretic recipe by which one can recover an affine smooth connected curve X defined over a finite field from $\pi_1(X)$. Moreover, if X is hyperbolic, then this recipe recovers X from $\pi_1^\dagger(X)$.*

Further, the absolute and the relative Isom-form of the anabelian conjecture for Curves holds for affine curves over finite fields; and its tame form holds for affine hyperbolic curves over finite fields.

(2) *There exists a group theoretic recipe by which one can recover affine hyperbolic curves $X \rightarrow k$ defined over finitely generated fields k of characteristic zero from $\pi_1(X)$.*

Further, the absolute and the relative Isom-form of the anabelian conjecture for Curves holds for affine hyperbolic curves over finitely generated fields of characteristic zero.

The strategy of the proof is as follows:

First consider the case when the base field k is finitely generated and has $\text{char}(k) = 0$. We claim that the canonical exact sequence

$$(*) \quad 1 \rightarrow \pi_1(\overline{X}) \rightarrow \pi_1(X) \rightarrow G_k \rightarrow 1,$$

is encoded in $\pi_1(X)$. Indeed, recall that the algebraic fundamental group $\pi_1(\overline{X})$ is a finitely generated normal subgroup of $\pi_1(X)$. Therefore, since G_k has no proper finitely generated normal subgroups, see e.g. [F–J], Ch.16, Proposition 16.11.6, it follows that $\pi_1(\overline{X})$ is the unique maximal finitely generated normal subgroup of $\pi_1(X)$. Thus the exact sequence above can be recovered from $\pi_1(X)$. Further, by either using the characterization of the geometric inertia elements in $\pi_1(X)$ given by NAKAMURA [Na1], or by using specialization techniques, one finally recovers the projection $\pi_1(X) \rightarrow \pi_1(X_0)$, where X_0 is the completion of X .

After having recovered the exact sequence $(*)$ above, one reduces the case of hyperbolic affine curves over finitely generated fields of characteristic zero to the π_1^\dagger -case of affine hyperbolic curves over finite fields. This is done by using the “standard reduction technique” mentioned above.

TAMAGAWA also shows that the absolute form of the Isom-conjecture and the relative one are roughly speaking equivalent (using the birational anabelian Conjecture described at the beginning of Part II).

We now turn our attention to the case of affine curves over finite fields, respectively the π_1^\dagger -case of hyperbolic curves over finite fields. TAMAGAWA’s approach is a tremendous refinement of UCHIDA’s strategy to tackle the birational case, i.e., to prove the birational anabelian Conjecture for global function fields. (Naturally, since $\pi_1(X)$ seems to encode much less information than the absolute Galois group $G_{k(X)}$, the things might/should be much more intricate in the case of curves.) A rough approximation of TAMAGAWA’s proof is as follows. Let $X \rightarrow k$ be an affine smooth geometrically connected curve, where k is a finite field with $\text{char}(k) = p$. As usual let X_0 be the smooth completion of X . Thus we have surjective canonical projections

$$\pi_1(X) \rightarrow \pi_1(X_0) \rightarrow G_k \rightarrow 1$$

and correspondingly for the tame fundamental groups.

The first part of the proof consists in developing a *Local Theory*, which as in the birational case, will give a description of the closed points $x \in X_0$ in terms of

the conjugacy classes of the decomposition groups $Z_x \subset \pi_1(X)$ above each closed point $x \in X$.

The steps for doing this are as follows:

Step 1) Recovering the several arithmetical invariants.

Here TAMAGAWA shows that the canonical projections $\pi_1(X) \rightarrow G_k$, thus $\pi_1(\overline{X})$, as well as $\pi_1(\overline{X}) \rightarrow \pi_1^t(\overline{X})$ are encoded in $\pi_1(X)$. Further, by a combinatorial argument he recovers the Frobenius element $\varphi_k \in G_k$. In particular, one gets the *cyclotomic character* of $\pi_1(X)$, and so one knows the ℓ -adic cohomology of $\pi_1(X)$, as well as the Galois action of G_k on the ℓ -adic Galois cohomology groups of $\pi_1(\overline{X})$.

The next essential remark is that after replacing X by some “sufficiently general” finite étale cover $Y \rightarrow X$, the completion $Y_0 \rightarrow k$ of Y is itself hyperbolic. In particular, the ℓ -adic Galois cohomology groups $H^i(\pi_1(\overline{Y}_0), \mathbb{Z}_\ell(r))$ of $\pi_1(\overline{Y}_0)$ are the same as the ℓ -adic étale cohomology $H^i(\overline{Y}_0, \mathbb{Z}_\ell(r))$ of \overline{Y}_0 . Thus by the remarks above, one can recover the ℓ -adic cohomology of \overline{Y}_0 for every étale cover $Y \rightarrow X$ having a hyperbolic completion Y_0 .

This is a fundamental observation in TAMAGAWA’s approach, as it can be used in order to tackle the following problem:

Which sections of the canonical projection $pr_X : \pi_1(X) \rightarrow G_k$ are defined by points $x \in X_0(k)$ in the way as described in the Section Conjecture?

We remark that since $G_k \cong \widehat{\mathbb{Z}}$ is profinite free on one generator, one cannot expect that all such sections are defined by points as asked by the Section Conjecture. (Indeed, there are uncountable many such conjugacy classes of sections, thus too “many” in order to be defined by points, even if X_0 has no k -rational points.)

Here is TAMAGAWA’s answer: Let $s : G_k \rightarrow \pi_1(X)$ be a given section. For every open neighborhood $U \subset \pi_1(X)$ of $s(G_k)$, we denote by $X_U \rightarrow X$ the finite étale cover of X classified by U . First, since U projects onto G_k , the curve $X_U \rightarrow k$ is geometrically connected. Further, we have in tautological way: $\pi_1(X_U) = U$, and $\overline{U} := U \cap \pi_1(\overline{X}) = \pi_1(\overline{X}_U)$. Let $X_{U,0}$ be the smooth completion of X_U . Then by Step 1), the canonical projection $\pi_1(X_U) \rightarrow \pi_1(X_{U,0})$ can be recovered from $U = \pi_1(X_U)$, thus from $\pi_1(X)$ endowed with the section $s : G \rightarrow \pi_1(X)$.

Now we remark that for U sufficiently small, the complete curve $X_{U,0}$ is hyperbolic, both in the case X is affine, or if X was hyperbolic and we were working $\pi_1^t(X)$. We set $U_0 := \pi_1(X_{U,0})$ and view it as quotient of $\pi_1(X_U)$, and $\overline{U}_0 = \pi_1(\overline{X}_{U,0})$. Since $X_{U,0}$ is complete and hyperbolic, the ℓ -adic cohomology group $H_{\text{et}}^i(\overline{X}_{U,0}, \mathbb{Z}_\ell(1))$ equals the Galois cohomology group $H^i(\pi_1(\overline{X}_{U,0}), \mathbb{Z}_\ell(1))$, thus the cohomology group $H^i(\overline{U}_0, \mathbb{Z}_\ell(1))$ for all i .

Finally, since the Frobenius element $\varphi_k \in G_k$ is known, by applying the Lefschetz Trace Formula, we can recover the *number of k -rational points* of X_{U_0} :

$$|X_{U_0}(k)| = \sum_{i=0}^2 (-1)^i \text{Tr}(\varphi_k) |H^i(\overline{U}_0, \mathbb{Z}_\ell(1))|$$

In this way we obtain the technical input for the following:

Proposition. *Let $s : G_k \rightarrow \pi_1(X)$ be a section of $pr_X : \pi_1(X) \rightarrow G_k$. Then s is defined by a point of $X_0(k)$ if and only if for every open sufficiently small neighborhood U of $s(G_k)$ as above, one has $X_{U_0}(k) \neq \emptyset$.*

Step 2) Recover the decomposition groups Z_x over closed points.

This is done using the Proposition above. Actually, using the Artin's Reciprocity law, one shows that in the case of a *complete hyperbolic curve*, like the X_{U_0} above, the set $X_{U_0}(k)$ is in bijection with the conjugacy classes of sections s defining points. And this gives a recipe to recover the points $X_0(k)$ which come from points in $X_{U_0}(k)$ for some U as above; thus finally for recovering all the points in $X_0(k)$. By replacing k by finite extension $l|k$, one recovers in a functorial way $X(l)$ too, etc. Thus finally one recovers the closed points x of X_0 as being in bijection with the conjugacy classes of decomposition groups $Z_x \subset \pi_1(X)$. Correspondingly the same is done for π_1^\dagger -case.

The second part of the proof is to develop a *Global Theory*, as done by UCHIDA in the birational case. Naturally, $\pi_1(X)$ endowed with all the decomposition groups over the closed points of X_0 carries much less information than $G_{k(X)}$ endowed with all the decomposition groups Z_v over the places v of $k(X)$.

Step 3) Recover the multiplicative group $k(X)^\times$ together with the valuations $v_x : k(X) \rightarrow \mathbb{Z}$.

Since the Frobenius elements $\varphi_x \in Z_x$ are known for closed points $x \in X_0$, by applying global class field theory as in the birational case, one gets the multiplicative group $k(X)^\times$ together with the valuations $v_x : k(X)^\times \rightarrow \mathbb{Z}$.

Step 4) Recovering the addition on $k(X)^\times \cup \{0\}$.

This is much more difficult than that in the birational case. And here is where the hypothesis that X is affine is used. Namely, if $x \in X_0 \setminus X$ is any point "at infinity", then from a decomposition group Z_x over x , one finally can recover the evaluation map

$$\mathfrak{p}_x : k(X) \rightarrow k^a \cup \infty.$$

One proceeds by applying the following:

Proposition. *Let $X_0 \rightarrow \kappa$ be a complete smooth curve over an algebraically closed field κ . Suppose that the multiplicative group $k(X_0)^\times$ together with the valuations $v_x : k(X_0)^\times \rightarrow \mathbb{Z}$ at closed points $x \in X_0$, and the evaluation of the functions at at least three k -points x_0, x_1, x_∞ of X are known. Then from these data the structure field of $k(X_0)$ can be recovered.*

In order to conclude the proof of Tamagawa's Theorem above over finite fields, we remark that all closed points of X_0 were recovered from $\pi_1(X)$ via the decomposition groups above them. In particular, such a point lies in X if and only if the inertia group above such a point is trivial. This gives the recipe to identify X inside X_0 . Thus we finally have a group theoretic recipe for recovering the curve $X \rightarrow k$ from its fundamental group $\pi_1(X)$.

Finally, the functoriality of the recipe for recovering X shows that any isomorphism of fundamental groups $\Phi : \pi_1(X) \rightarrow \pi_1(Y)$ is defined by an isomorphism of function fields $\phi : k(X_0) \rightarrow k(Y_0)$ which induces an isomorphism of schemes $X \rightarrow Y$. The uniqueness of ϕ up to Frobenius twists follows the same pattern as in the birational case, but using the fact that the center of $\pi_1(X)$ is trivial in the cases under discussion.

II) Mochizuki's results for hyperbolic curves in characteristic zero

In this subsection we discuss briefly some of MOCHIZUKI's results concerning hyperbolic curves in characteristic zero.

The first such result was announced by MOCHIZUKI shortly after Tamagawa's Theorem discussed above. The result deals with hyperbolic curves over finitely generated fields of characteristic zero, and more or less extends the corresponding result by TAMAGAWA to *complete hyperbolic curves*. The proof relies heavily on Tamagawa's Theorem, but MOCHIZUKI's strategy for the proof goes beyond TAMAGAWA's approach.

Theorem (See MOCHIZUKI [Mzk1]).

The hyperbolic curves over finitely generated fields of characteristic zero are anabelian. Further, both the relative and the absolute Isom-form of the anabelian Conjecture for hyperbolic curves over such fields hold.

We indicate briefly the idea of the proof. Let $X \rightarrow k$ be a hyperbolic curve over a finitely generated field k of characteristic zero. Proceeding as TAMAGAWA did in the case of affine hyperbolic curves, we can recover the exact sequence

$$1 \rightarrow \pi_1(\overline{X}) \rightarrow \pi_1(X) \rightarrow G_k \rightarrow 1,$$

and also the projection $\pi_1(X) \rightarrow \pi_1(X_0)$, where X_0 is the completion of X . In this way one reduces the question to the case of *complete hyperbolic curves*.

Thus let $X \rightarrow k$ be a complete hyperbolic curve over some finitely generated field k of characteristic zero. The idea of MOCHIZUKI is to reduce the problem in this case to the π_1^{t} -case of affine hyperbolic curves over finite fields and then use Tamagawa's Theorem for affine hyperbolic curves over finite fields. In order to do that, MOCHIZUKI uses log-schemes and log-fundamental groups. In essence one does the following: In the context above, recall the setting explained in the "standard reduction technique". In the notations from there, let v_s be a discrete

valuation of k with valuation ring R_s dominating some closed point $s \in S$ such that the residue field equals $\kappa(s)$, thus finite. Let $p = \text{char}(\kappa(s))$, thus $\kappa(s)$ is finite over \mathbb{F}_p . In the case $X \rightarrow k$ has good reduction at s —and this is the case on a Zariski open subset of S , in particular if p is big enough, the special fiber $X_s \rightarrow \kappa(s)$ of $X_{R_s} \rightarrow R_s$ at s is a complete hyperbolic curve. Thus one cannot apply and use Tamagawa’s Theorem in order to recover $X_s \rightarrow \kappa(s)$ from $\pi_1^\dagger(X_s)$ (even if the projection $\pi_1^\dagger(X_s) \rightarrow G_{\kappa_v}$ is known). Nevertheless, the fact that X has good reduction at v is encoded in the canonical exact sequence $\pi_1(X) \rightarrow G_k$ endowed with a decomposition group $Z_{v_s} \subset G_k$ above v_s by Oda’s Criterion for good reduction of hyperbolic complete curves.

In the above notations, suppose that $X_{R_s} \rightarrow R_s$ is smooth. Let us consider a finite Galois étale cover $Y^{(p)} \rightarrow X$ such that its geometric part $\overline{Y}^{(p)} \rightarrow \overline{X}$ is the maximal p -elementary Abelian cover of \overline{X} . After enlarging k , we eventually can suppose that $Y^{(p)} \rightarrow k$ is geometrically connected, and that $\text{Aut}_X(Y^{(p)})$ is defined over k . Under this hypothesis one has:

$$\deg(Y^{(p)} \rightarrow X) = p^{2g_X} \quad (g_X \text{ is the genus of } X).$$

On the other hand, considering the maximal geometric p -elementary Abelian cover $Z^{(p)} \rightarrow X_s$, and recalling that $r_{X_s} \leq g_{X_s} = g_X$ denotes the Hasse–Witt invariant of X_s , we see that

$$\deg(Z^{(p)} \rightarrow X_s) = p^{r_{X_s}} \leq p^{g_X}.$$

We conclude that $Y^{(p)} \rightarrow k$ does not have potentially good reduction. Moreover, for p getting larger, the special fiber $Y_s^{(p)} \rightarrow \kappa_s$ of the stable model of $Y^{(p)} \rightarrow k$ (which is defined over some finite extension $l|k$ and corresponding extensions R_w of R_{v_s} , etc.) has “many” double points. We set $Y_s^{(p)} = \cup_i Y_i$, where Y_i are the irreducible components of Y_s . For each Y_i , let U_i be the smooth part of Y_s inside Y_i . Since Y_s is connected, it follows that each $U_i \rightarrow \kappa_w$ is an affine hyperbolic curve over the finite field κ_w .

Now it is part of the theory of log-fundamental groups, that the tame fundamental group $\pi_1^\dagger(U_i)$ can be recovered from $\pi_1(Y^{(p)}) \rightarrow G_l$. Thus applying Tamagawa’s Theorem for the π_1^\dagger -case of affine hyperbolic curves over finite fields, we can recover $U_i \rightarrow \kappa_w$ in a functorial way from $\pi_1^\dagger(U_i)$. And finally, one can recover $Y_s^{(p)} \rightarrow \kappa_w$, and $X_s \rightarrow \kappa_v$ as well.

One concludes by using the standard reduction/globalization techniques.

The result above by MOCHIZUKI is the precursor of his much stronger result concerning hyperbolic curves over sub- p -adic fields as explained below. First let us introduce Mochizuki’s notations. A *sub- p -adic field* k is any field which can be embedded into some function field over \mathbb{Q}_p . Let k be a sub- p -adic field, and let

$X \rightarrow k$ be a geometrically connected scheme over k . Consider the exact sequence of fundamental groups

$$1 \rightarrow \pi_1(\overline{X}) \rightarrow \pi_1(X) \rightarrow G_k \rightarrow 1.$$

We denote $\Delta_X := \pi^p(\overline{X})$ the maximal pro- p quotient of $\pi_1(\overline{X})$, and remark that the kernel N of the map $\pi_1(\overline{X}) \rightarrow \Delta(X)$ is a characteristic subgroup of $\pi_1(\overline{X})$, i.e., it is invariant under all automorphisms of $\pi_1(\overline{X})$. In particular, N is invariant under the conjugation in $\pi_1(X)$. Thus N is a normal subgroup in $\pi_1(X)$ too. We will set $\Pi_X = \pi_1(X)/N$. Therefore, the above exact sequence gives rise to a canonical exact sequence of fundamental groups:

$$1 \rightarrow \Delta_X = \pi_1^p(\overline{X}) \rightarrow \Pi_X = \pi_1(X)/N \rightarrow G_k \rightarrow 1.$$

With these notations, the main result by MOCHIZUKI can be stated as follows:

Theorem (See MOCHIZUKI [Mzk3]).

Let $Y \rightarrow k$ be a geometrically integral hyperbolic curve over a sub- p -adic field. Then Y can be recovered from the canonical projection $\Pi_Y \rightarrow G_k$.

Moreover, this recipe is functorial in such a way that it implies the following Hom-form of the relative anabelian Conjecture for curves: Let $X \rightarrow k$ be a geometrically integral smooth variety. Then every open G_k -homomorphism $\Pi_X \rightarrow \Pi_Y$ is defined in a functorial way by a unique dominant k -morphism $\phi : X \rightarrow Y$.

The main tools used by MOCHIZUKI are the p -adic Tate–Hodge Theory and Faltings’ Theory of almost étale morphisms. The proof is very technical and difficult to follow for non-experts (maybe even for experts!). I will nevertheless try to summarize here the main points in the proof (which are though more intricate and complex, than I might suggest here...). I should also mention that in this case we do not have a recipe to recover $X \rightarrow k$ from $\pi_X \rightarrow G_k$, which is as explicit as in the previous cases. The main difficulty in this respect lies in not having an *explicit local theory* as in the previous case. In particular and unfortunately, until now we do not have a way of describing $X(k)$, i.e., we do not have any kind of an answer to the Section Conjecture so far.

Coming back to the proof of the Theorem above, the first observation is that via more or less standard specialization techniques, the problem is reduced to the following case: $k|\mathbb{Q}_p$ is a finite extension, and $X \rightarrow k$ is a smooth hyperbolic curve, and $Y \rightarrow k$ is a complete hyperbolic curve. And finally replacing X by the étale cover classified by the image of Φ , one can suppose that Φ is surjective.

One should remark that a further reduction step to the case where X is complete is not at all trivial, and it is one of the facts which complicates things a lot. Naturally, by using the canonical projection $\Pi_{k(X)} \rightarrow \Pi_X$, one might reformulate the problem above correspondingly, and ask whether every surjective G_k -morphism $\Phi : \Pi_{k(X)} \rightarrow \Pi_Y$ is defined by a unique morphism of function

fields $k(Y) \hookrightarrow k(X)$ over k . We will nevertheless take this last reduction step for granted, and only outline the proof of the following:

Let $X \rightarrow k$ and $Y \rightarrow k$ be complete hyperbolic curves, where $k|\mathbb{Q}_p$ is finite. Then every surjective G_k -homomorphism $\Phi : \Pi_X \rightarrow \Pi_Y$ is defined by a unique k -morphism $\phi : X \rightarrow Y$ in a functorial way.

The *Local Theory* in this case is as follows: As mentioned above, one has no clue at all how to recover $X(k)$ from the given data $\Pi_X \rightarrow G_k$, and the situation is not better if we replace Π_X by the full étale fundamental group $\pi_1(X)$. Nevertheless, MOCHIZUKI develops another kind of a “Local Theory”, which fortunately does the right job for the problem. The kind of points one can recover are as follows: Let R be the valuation ring of k , and let $X_R \rightarrow R$ be a semi-stable model of $X \rightarrow k$ (such models exist after enlarging the base field k). Let $(X_i)_i$ be the irreducible components of the special fiber $X_s \rightarrow \kappa$ of $X_R \rightarrow R$. If $\eta_i \in X_R$ is the generic point of X_i , then the local ring $\mathcal{O}_{X_R, \eta_i}$ is a discrete valuation ring of $k(X)$ dominating R , and the residue field $\kappa(\eta_i)$ is the function field of $X_i \rightarrow \kappa$. Let us call such points η_i *arithmetical points* of X (arising for the several models X_R).

Next let (L, v) be a discrete complete valued field over k such that the valuation of L prolongs the p -adic valuation of k , and the residue field $Lv | kv$ is a function field in one variable. Remark that the completion of $k(X)$ with respect to the valuation $v := v_{\eta_i}$ defined by an arithmetical point is actually such a discrete complete valued field over k . We denote for short $\mathcal{H}_L^\Omega = H^1(G_{Lk^a}, \widehat{\mathcal{O}}_{L^a}(1))/(\text{torsion})$, where $\widehat{\mathcal{O}}_{L^a}$ is the completion of the valuation ring of L^a (similar to the completion \mathbb{C}_p of the algebraic closure of \mathbb{Q}_p).

We will say that a G_k -homomorphism $\Phi_X : G_L \rightarrow \Pi_X$ is *non-degenerate*, if the induced map on the p -adic cohomology

$$H^1(\Delta_X, \mathbb{Z}_p(1)) \xrightarrow{\text{infl}_{\Phi_X}} H^1(G_L, \widehat{\mathcal{O}}_{L^a}(1)) \xrightarrow{\text{can}} \mathcal{H}_L^\Omega$$

is non-trivial. Now the main technical points of the proof are as follows:

1) In the context above, let $\Phi : \Pi_X \rightarrow \Pi_Y$ be an open G_k -homomorphism. Then there exists a non-degenerate G_k -homomorphism $\phi_L : G_L \rightarrow \Pi_X$ such that the composition $\Phi_Y := \Phi \circ \phi_L$ is a non-degenerate homomorphism. Thinking of the Local Theory from the birational case, this assertion here corresponds more or less to the characterization of arithmetical Zariski prime divisors.

2) Every non-degenerate G_k -morphism $\Phi_X : G_L \rightarrow \Pi_X$ as above is of geometrical nature: Given such a Φ_X , there exists a unique L -rational point $\psi_{\Phi_X} : L \rightarrow X$ defining Φ_X in a functorial way.

3) In particular, for Φ and Φ_X as at 1) above, there exist L -rational points $\psi_{\Phi_X} : L \rightarrow X$ and $\psi_{\Phi_Y} : L \rightarrow Y$ defining the non-degenerate morphisms Φ_X and $\Phi_Y = \Phi \circ \phi_L$ in a functorial way.

Finally, MOCHIZUKI's *Global Theory* is a very nice application of the p -adic Hodge–Tate Theory and of Faltings' Theory of almost étale morphisms. The idea is as follows:

First, by the p -adic Hodge–Tate Theory, the sheaf of global differentials on X , say $D_X := H^0(X, \Omega_X)$, can be recovered from the action of G_k on the \mathbb{C}_p -twists with the p -adic cohomology $H_{\text{et}}^i(\overline{X}, \mathbb{Z}_p(j))$ of \overline{X} . On the other hand, since X is a complete hyperbolic curve over a field of characteristic zero, thus $\neq p$, the p -adic cohomology $H_{\text{et}}^i(\overline{X}, \mathbb{Z}_p(j))$ is the same as the Galois cohomology of Δ_X , thus known.

Let us denote $D_X^i = H^0(X, \Omega_X^{\otimes i})$, and let $\mathcal{R}_X^i := \ker(D_X^{\otimes i} \rightarrow D_X^i)$ be the space of i^{th} homogeneous relations in D_X^i . If X is not a hyperelliptic curve (what we can suppose after replacing X by a properly chosen étale cover whose geometric part is p -elementary Abelian), then the system of all the D_X^i completely defines X . Equivalently, the system of all the data $\mathcal{R}_X^i \subset D_X^{\otimes i}$ completely defines X .

Second, let $\Phi : \Pi_X \rightarrow \Pi_Y$ be a surjective G_k -homomorphism. Then Φ induces in a functorial way a morphism of k vector spaces $\iota_\Phi : D_Y \rightarrow D_X$; thus also morphisms of k vector space $\iota_\Phi^{\otimes i} : D_Y^{\otimes i} \rightarrow D_X^{\otimes i}$ for each $i \geq 1$. And by the general non-sense concerning the canonical embedding, if each $\iota_\Phi^{\otimes i}$ “respects the relations”, i.e., it maps \mathcal{R}_Y^i into \mathcal{R}_X^i , then ι_Φ is defined by some dominant k -morphism $\phi : X \rightarrow Y$ in the canonical way.

Finally, in order to check that ι_Φ does indeed respect the relations, one uses the Local Theory and Faltings' Theory of almost étale morphisms: Choose a non-degenerate morphism $\Phi_X : G_L \rightarrow \Pi_X$ as at 3) above. Let Ω_L denote the p -adically continuous k -differentials of L , and Ω_L^i be its powers. Since ψ_{Φ_X} is a non-degenerate point of X , the differential $d_X := d(j_{\Phi_X}) : D_X \rightarrow \Omega_L$ of $j_{\Phi_X} : L \rightarrow X$ and its powers $d_X^i : D_X^i \hookrightarrow \Omega_L^i$ are embeddings. Thus in order to check that ι_Φ respects the relations, it is sufficient to check that this is the case for the composition

$$\iota_L^{\otimes i} : D_Y^{\otimes i} \rightarrow D_X^{\otimes i} \hookrightarrow \Omega_L^{\otimes i}.$$

On the other hand, the composition of the map $\iota_L^{\otimes i}$ with $\Omega_L^{\otimes i} \rightarrow \Omega_L^i$ is exactly the canonical map $D_Y^{\otimes i} \rightarrow D_Y^i \hookrightarrow \Omega_L^i$ defined via the non-degenerate morphism $\Phi_Y = \Phi \circ \Phi_X$ and the resulting point $\psi_{\Phi_Y} : L \rightarrow \Pi_Y$. This concludes the proof.

Remarks.

1) First, Theorem A' of Mochizuki [Mzk3] shows that also *truncated*- Π_X *versions* of the assertion of the main result above are valid. One can namely replace Δ_X by its central series quotient $\Delta_X^{(n)}$, and consequently Π_X by the corresponding quotient $\Pi_X^{(n)}$ which fits into the exact sequence

$$1 \rightarrow \Delta_X^{(n)} \rightarrow \Pi_X^{(n)} \rightarrow G_k \rightarrow 1.$$

Let $n \geq 3$. Then given an open G_k -homomorphism $\Phi^{(n+2)} : \Pi_Y^{(n+2)} \rightarrow \Pi_X^{(n+2)}$ there exists a unique dominant k -homomorphism $Y \rightarrow X$ such that the canonical open morphism $\Pi_Y^{(n)} \rightarrow \Pi_X^{(n)}$ induced by ϕ coincides with $\Phi^{(n)}$ on $\Pi_Y^{(n)}$.

2) Using specializations techniques, MOCHIZUKI proves the relative Hom-form of the birational anabelian Conjecture for finitely generated fields over sub- p -adic fields k as follows:

Theorem (MOCHIZUKI [Mzk3], Theorem B).

Let $K|k, L|k$ be regular function fields. Then every open G_k -homomorphism $\Phi : G_K \rightarrow G_L$ is defined functorially by a unique k -embedding of fields $L \rightarrow K$.

I would like to remark that using techniques developed in order to prove a pro- p form of the birational conjecture, one can sharpen the above result and show the following: *Every open Π_k -homomorphism $\Phi : \Pi_K \rightarrow \Pi_L$ is defined by a unique k -embedding $L \rightarrow K$ in a functorial way.*

3) MOCHIZUKI also shows that *hyperbolically fibered surfaces* are anabelian. And moreover, the Isom-form of an anabelian Conjecture for fibered surfaces is true. Here, a hyperbolically fibered surface X is the complement of an étale divisor in a smooth proper family $\tilde{X} \rightarrow X_1$ of hyperbolic complete curves over a hyperbolic base curve X_1 . The result is:

Theorem (MOCHIZUKI [Mzk3], Theorem D).

Let $Y \rightarrow k$ and $X \rightarrow k$ be geometrically integral hyperbolically fibered surfaces over a sub- p -adic field k . Then every G_k -isomorphism $\Phi : \pi_1(Y) \rightarrow \pi_1(X)$ is defined by a unique k -isomorphism $\phi : Y \rightarrow X$ in a functorial way.

Note that in the Theorem above the *full fundamental group* π_1 is needed. It is maybe useful to remark that a naive Hom-form of the above Theorem is not true. Indeed, let k be an infinite base field. Then using general hyperplane arguments, one can show that for every smooth quasi-projective k -variety $X \subseteq \mathbb{P}^N$, there exist smooth k -curves $Y \subseteq X$ obtained from $X \rightarrow k$ by intersections with general hyperplanes such that the canonical map $\pi_1^t(Y) \rightarrow \pi_1^t(X)$ is surjective. In a second step, one can realize $\pi_1^t(Y)$ in many ways as quotients of fundamental groups $\pi_1^t(Z) \rightarrow \pi_1^t(Y)$ for several smooth k -varieties (which can be chosen to be projective, if X is complete), e.g., $Z = Y \times \dots \times Y$ finitely many times. Finally, the composition

$$\pi_1^t(Z) \rightarrow \pi_1^t(Y) \rightarrow \pi_1^t(X)$$

is a surjective G_k -morphism, but by its construction, it does not originate from a dominant k -rational map.

III) Jakob Stix's results concerning hyperbolic curves in positive characteristic

The results of STIX deal with *hyperbolic non-constant curves* over finitely generated infinite fields k of positive characteristic (but apply as well to such

fields of characteristic zero, where the results are already known). Recall that given a curve $X \rightarrow k$ with $\text{char}(k) = p > 0$, one says that X is *potentially isotrivial*, if there exists a finite étale cover $X' \rightarrow X$ such that X' is defined over a finite field. One can show that if X is hyperbolic, then X is potentially isotrivial if and only if there exists a finite field extension $k'|k$ such that the base change $X' = X \times_k k'$ is defined over a finite field. Further, recall that for a curve $X \rightarrow k$ as above, we denote by $X^i \rightarrow k^i$ the maximal purely inseparable cover of $X \rightarrow k$. And for every integer n we denote by $X^i(n) \rightarrow k^i$ the relative Frobenius n -twist.

Theorem (STIX [St1], [St2]).

Let $X \rightarrow k$ be a non potentially isotrivial hyperbolic curve over a finitely generated infinite field k with $\text{char}(k) = p > 0$. Then one can recover $X^i \rightarrow k^i$ from $\pi_1^\dagger(X) \rightarrow G_k$ in a functorial way.

Moreover, the relative Isom-form of the anabelian Conjecture for hyperbolic curves over k is true in the following sense: Let $Y \rightarrow k$ be some hyperbolic curve, and let a G_k -isomorphism $\Phi : \pi_1^\dagger(X) \rightarrow \pi_1^\dagger(Y)$ be given. Then there exists a unique n and a k^i -isomorphism $\phi : X^i(n) \rightarrow Y^i$ defining Φ .

The strategy of proof is as follows:

Let k be a finitely generated infinite field, and $X \rightarrow k$ a hyperbolic curve over k . In the notations from the “standard reduction technique”, let $X_S \rightarrow S$ be a smooth surjective family of hyperbolic curves whose generic fiber is $X \rightarrow k$. The idea is as follows:

Case 1. $X \rightarrow k$ is an affine hyperbolic curve.

By shrinking S if necessary, we can suppose that $X \rightarrow k$ has good reduction at all closed points $s \in S$. From $\pi_1^\dagger(X) \rightarrow G_k$ one recovers the local projections $\Phi_s : \pi_1^\dagger(X_s) \rightarrow G_{\kappa(s)}$ for all closed points $s \in S$. By Tamagawa’s Theorem, we can recover the isomorphy type of $X_s^i \rightarrow \kappa(s)$ up to Frobenius twists. In particular, let $\Phi : \pi_1^\dagger(X) \rightarrow \pi_1^\dagger(Y)$ be a G_k -isomorphism, where $Y \rightarrow k$ is some hyperbolic curve over k . By the “standard specialization technique”, we obtain $\kappa(s)$ -isomorphisms of some relative Frobenius twists of the special fibers, say $\phi_s : X_s^i(n_s) \rightarrow Y_s^i$ defining Φ_s . Unfortunately, the usual globalization techniques work only under the hypothesis the Frobenius twists n_s are constant, say equal to n , on a non-empty open of S (and then they turn to be constant on the whole S). If this is the case, then the local isomorphisms ϕ_s originate indeed from a unique global k^i -isomorphism $\phi : X^i(n) \rightarrow Y^i$, which defines the given G_k -isomorphism $\Phi : \pi_1^\dagger(X) \rightarrow \pi_1^\dagger(Y)$.

Here is the way STIX shows that the exponents n_s are indeed constant: First, by replacing X by a properly chosen tame étale cover, we can suppose that the smooth completion X_0 of X is hyperbolic too. Next we fix some $m > 2$ relatively prime to $p = \text{char}(k)$, and replace k by its finite extension over which the m -torsion of Jac_{X_0} becomes rational. And choose an m -level structure on X_0 by

fixing an isomorphism

$$\varphi_{X,m} : {}_m\text{Jac}_{X_0} = \pi_1^{\text{ab}}(\overline{X}_0)/m \rightarrow (\mathbb{Z}/m)^{2g}$$

Then X_0 endowed with $\varphi_{X,m}$ is classified by a k -rational point $\psi_X : k \rightarrow \mathcal{M}_g[m]$. Moreover, using the “standard reduction technique”, the level structure $\varphi_{X,m}$ gives rise via the specialization homomorphisms $\text{sp}_s : \pi_1(\overline{X}_0) \rightarrow \pi_1(\overline{X}_{0,s})$ canonically to level structures

$$\varphi_{X_s,m} : {}_m\text{Jac}_{X_{0,s}} = \pi_1^{\text{ab}}(\overline{X}_{0,s})/m \rightarrow (\mathbb{Z}/m)^{2g}.$$

And this happens in such a way that $\psi_X : k \rightarrow \mathcal{M}_g[m]$ defined above becomes the generic fiber of a morphism $\psi_{X_S} : S \rightarrow \mathcal{M}_g[m]$ whose special fibers classify the curves $X_s \rightarrow \kappa(s)$ endowed with the level structures $\varphi_{X_s,m}$.

Now let us come back to the G_k -isomorphism $\Phi : \pi_1^{\dagger}(X) \rightarrow \pi_1^{\dagger}(Y)$. Clearly, Φ transports the m -level structure $\varphi_{X,m}$ of X_0 to an m -level structure $\varphi_{Y,m}$ for Y_0 . And the local $G_{\kappa(s)}$ -isomorphisms $\Phi_s : \pi_1^{\dagger}(X_s) \rightarrow \pi_1^{\dagger}(Y_s)$ transport the m -level structures $\varphi_{X_s,m}$ to m -level structures $\varphi_{Y_s,m}$ which are compatible with the specialization morphisms $\text{sp}_s : \pi_1(\overline{Y}_0) \rightarrow \pi_1(\overline{Y}_{0,s})$.

Now let us suppose that there exist some exponents n_s and $\kappa(s)$ -isomorphisms $\phi_s : X_s^i(n_s) \rightarrow Y_s^i$ which define the G_{κ} -isomorphisms $\Phi_s : \pi_1^{\dagger}(X_s) \rightarrow \pi_1^{\dagger}(Y_s)$. Then ϕ_s prolongs to an $\kappa(s)$ -isomorphism $\phi_{0,s} : X_{0,s}(n_s) \rightarrow Y_{0,s}$. Next remark that $X_{0,s}$ and its relative Frobenius twists endowed with the same m -level structure $\varphi_{X_s,m} = \varphi_{X_s(n_s),m}$ factor through the same closed point of $\mathcal{M}_g[m]$. Thus we have: The classifying morphisms $\psi_{X_S} : S \rightarrow \mathcal{M}_g[n]$ for $X_{0,S}$ and $\psi_{Y_S} : S \rightarrow \mathcal{M}_g[n]$ for $Y_{0,S}$ defined above coincide (topologically) on the closed points $s \in S$.

In order to conclude, STIX proves the following:

Proposition (STIX [St1]).

Let S and \mathcal{M} be irreducible \mathbb{Z} -varieties. Let $f, g : S \rightarrow \mathcal{M}$ be two morphisms which coincide topologically on the closed points of S . Suppose that $f \neq g$. Then S is defined over \mathbb{F}_p for some p , and f and g differ by a power of Frobenius, which is unique if f is not constant.

Thus applying the Proposition above we conclude that the classifying morphisms ϕ_X and ϕ_Y differ by a power Frob^n of Frobenius. In particular, fiber wise the same is the case. From this one finally deduces that $\Phi : \pi_1^{\dagger}(X) \rightarrow \pi_1^{\dagger}(Y)$ is defined by some k -isomorphisms $\phi : X^i(n) \rightarrow Y^i$ for some integer n .

This completes the proof of the case when X is an affine hyperbolic curve.

Case 2) $X \rightarrow k$ is a complete hyperbolic curve.

Let us try to mimic MOCHIZUKI’s strategy from the case of complete hyperbolic curves over finitely generated fields of characteristic zero. Then we run immediately into the following difficulty: If k has positive characteristic $p > 0$,

there is no obvious reason that some finite properly chosen étale (Galois) covers $X' \rightarrow X$ have bad reduction at points $s \in S$ where X has good reduction. Note that the “trick” used by MOCHIZUKI in [Mzk2] in the case $\text{char}(k) = 0$ does definitely not work in positive characteristic. (This follows from Grothendieck’s Specialization Theorem: Let $X_{R_s} \rightarrow R_s$ be a complete smooth curve, and let $X' \rightarrow X$ be an étale Galois cover whose geometric part has degree prime to p . Then X' has potentially good reduction.)

In order to avoid this difficulty, one can nevertheless use the Raynaud, Pop–Saidi, Tamagawa Theorem, see Part III) of these notes. A consequence of that result is the following: Let a closed point $s \in S$ be given. Then there exists a finite étale cover $X^{(s)} \rightarrow X$ whose geometric part is a cyclic étale cover of \overline{X} of degree prime to p having the property: Any finite étale cover $X' \rightarrow X^{(s)}$ whose geometric part factors through the maximal p -elementary étale cover of $\overline{X}^{(s)}$ *does not have potentially good reduction*. With this input, STIX uses the theory of log-étale fundamental groups in order to conclude the proof in the same style as MOCHIZUKI [Mzk1], but using the methods developed to treat the case of affine hyperbolic curves.

PART III: Beyond Grothendieck’s anabelian Geometry

It is/was a widespread believe that the reason for the existence of anabelian schemes is strong interaction between the arithmetic and a rich algebraic fundamental group, and that this interaction makes étale fundamental groups so rigid, that the only way isomorphisms, respectively open homomorphisms, can occur is the geometrical one. (To say so, morphisms between étale fundamental groups which do not have a reason to exist, do not exist indeed...)

On the other hand, some developments from the 1990’s showed evidence for very strong anabelian phenomena for curves and higher dimensional varieties over *algebraically closed fields*, thus in a total absence of a Galois action of the base field. We mention here the following:

a) *Bogomolov’s Program* (see [Bo])

Let ℓ be a fixed rational prime number. For algebraically closed base fields k of characteristic $\neq \ell$, we consider integral k -varieties $X \rightarrow k$, with function field $k(X)|k$. It turns out that there is a major difference between the cases $\dim(X) = 1$ and $\dim(X) > 1$. Indeed, if $\dim(X) = 1$, then the absolute Galois group $G_{k(X)}$ is profinite free on $|k|$ generators. This is the so called *Geometric case of a Conjecture of Shafarevich*, proved by HARBATER [Ha2], and POP [Po]. On the other hand, if $d = \dim(X) > 1$, then $G_{k(X)}$ is very complicated (in particular, having $\text{cd}_\ell G_{k(X)} = d$, etc.).

The guess of BOGOMOLOV is that in the latter case, i.e., if $\dim(X) > 1$, the Galois group $G_{k(X)}$ should encode the birational class of X up to pure inseparable

covers and Frobenius twists. More precisely, BOGOMOLOV proposes and gives strong evidence for the following: In the context above, let $G_K(\ell)$ be the maximal pro- ℓ quotient of G_K , i.e., $G_K(\ell)$ is the Galois group of the maximal Galois pro- ℓ sub-extension $K(\ell)$ of $K^s|K$. Further let $G_K^{(n)}$ denote the n^{th} factor in the central series of $G_K(\ell)$, and by $K^{(n)}$ the corresponding fixed fields inside $K(\ell)$. In particular, $K^{(1)} = K^{\ell, \text{ab}}$ is the maximal pro- ℓ Abelian extension of K , and $K^{(2)}$ is the maximal central extension of $K^{\ell, \text{ab}}$ inside $K(\ell)$. Now BOGOMOLOV claims and gives evidence for the fact that the isomorphism type of the function field $k(X)^i|k$ is encoded in the *second factor group* $\text{PGal}_K^c := G_K(\ell)/G_K^{(2)}$. The starting point in this development was BOGOMOLOV's observation that if a subgroup $\Gamma \cong \mathbb{Z}_\ell \times \mathbb{Z}_\ell$ of $G_K^{\ell, \text{ab}}$ which can be lifted to a commutative subgroup of PGal_K^c must contain inertia elements for some non-trivial valuation v on K .

b) *Tamagawa's Theorem concerning $\mathbb{P}_{k_0}^1 \setminus \{0, 1, \infty, x_1, \dots, x_n\}$*

In the mid 1990's TAMAGAWA gave evidence for the fact that some curves over the algebraic closure $k_0 = \mathbb{F}_p$ are weakly anabelian, i.e., their isomorphism type as a scheme can be recovered from π_1 or even π_1^\dagger . The first precursor of this fact is Tamagawa's result that given a smooth curve $X \rightarrow k$, the type (g, r) of the curve is encoded in the algebraic fundamental group $\pi_1(\overline{X})$; and moreover, the canonical projections $\pi_1(\overline{X}) \rightarrow \pi_1^\dagger(\overline{X}) \rightarrow \pi_1(X_0)$ are encoded in the algebraic fundamental group of X . This answered a question raised by HARBATER. And finally, TAMAGAWA [T2] showed the following:

Let $U = \mathbb{P}_{k_0}^1 \setminus \{0, 1, \infty, x_1, \dots, x_n\}$ be an affine open. Then the isomorphism type of U as a scheme can be recovered from $\pi_1^\dagger(U)$. Moreover, if X is any other curve over some algebraically closed field k , and $\pi_1(X) \cong \pi_1(U)$, then $k = k_0$, and $X \cong U$ as schemes.

The kind of results above show that one can expect anabelian phenomena over algebraically closed base fields, thus in a complete absence of arithmetical Galois action. This kind of anabelian phenomena go *beyond Grothendieck's anabelian Geometry*. Here is a short list of the kind of such anabelian results.

A) Small Galois groups and valuations

Let ℓ be a fixed prime number. We consider fields K of characteristic $\neq \ell$, such that $\mu_\ell \subset K$. We denote by $K(\ell)$ the maximal Galois pro- ℓ extension of K in some fixed algebraic closure K^a of K , and denote by $G_K(\ell)$ the Galois group of $K(\ell)|K$. In order to avoid complications arising from orderings in case $\ell = 2$, we will also suppose that $\mu_4 \subset K$ if $\ell = 2$.

In the above context, let v be a non-trivial valuation of $K(\ell)$ such that value group vK is not ℓ -divisible and the residue field Kv has characteristic $\neq \ell$. Let $V_v \subseteq T_v \subseteq Z_v$ be respectively the ramification, the inertia, and the decomposition groups of v in $G_K(\ell)$. Then by the Hilbert decomposition theory for valuations

one has, see e.g., [BOU]: $V_v = \{1\}$, as $\text{char}(Kv) \neq \ell$. Thus $T_v = T_v/V_v$ is an Abelian pro- ℓ group. Further, $K(\ell)v = (Kv)(\ell)$, and one has the canonical exact sequence:

$$1 \rightarrow T_v \rightarrow Z_v \rightarrow G_{Kv}(\ell) \rightarrow 1$$

Finally, $vK(\ell)$ is the ℓ -divisible hull of vK . And denoting by \widehat{vK} the ℓ -adic completion of vK , there is an isomorphism of G_{Kv} -modules $T_v \cong \text{Hom}(\widehat{vK}, \mathbb{T}_\ell)$ where $\mathbb{T}_\ell = \varprojlim^m \mu_{\ell^m}$ is the Tate module of Kv . This reduces the problem of describing Z_v to that of describing $Kv(\ell)$. But the essential observation here is that T_v is a non-trivial Abelian normal subgroup of Z_v .

The following result is based on work by WARE [W] if $\ell = 2$, and KOENIGSMANN [Ko1] if $\ell \neq 2$, see also EFRAT [Ef1], [Ef2]. It is the best possible converse to the above description of Z_v :

Theorem (ENGLER–KOENIGSMANN [E–K]).

In the above notations let $Z \subset G_K(\ell)$ be a closed non-procyclic subgroup having a non-trivial Abelian normal subgroup T . Then there exists a valuation w of $K(\ell)$ with the following properties:

- (i) $Z \subset Z_w$ and $T \subset T_w$.
- (ii) The residue field Kw has $\text{char}(Kw) \neq \ell$.

The proof of the Theorem above is based on a fine analysis of the multiplicative structure of fields with *very small pro- ℓ Galois group*. We will say namely that K has a very small pro- ℓ Galois group, if $K(\ell)$ is non pro-cyclic, but fits into an exact sequence of the form $0 \rightarrow \mathbb{Z}_\ell \rightarrow K(\ell) \rightarrow \mathbb{Z}_\ell \rightarrow 0$. In such a case one simply can write down the valuation ring of a valuation w on K satisfying the properties (i), (ii) above, see loc.cit.. The rest is just valuation theory techniques.

The above assertion concerning fields with very small pro- ℓ Galois group has a parallel assertion by BOGOMOLOV which was suggested in [Bo], and finally proved by BOGOMOLOV–TSCHINKEL. The assertion is as follows:

Theorem (BOGOMOLOV–TSCHINKEL [B–T1]).

Suppose that K contains an algebraic closure k of its prime field. Suppose that $\Gamma \subseteq G_K^{\ell, \text{ab}}$ is a non-procyclic closed subgroup which can be lifted to an Abelian subgroup of $\text{PGal}^c(K)$. Then there exists a valuation w of K and a non-trivial subgroup $T \subset \Gamma$ such that denoting by T_w the inertia group of w in $G_K^{\ell, \text{ab}}$ the following hold:

- (i) $T \subseteq T_w$
- (ii) The residue field Kw has $\text{char}(Kw) \neq \ell$.

The proof relies on a very ingenious idea of BOGOMOLOV to compare maps between affine geometries and projective geometries. The two kind of geometries

arise as follows: First, by Kummer Theory one has an canonical map

$$K^\times \xrightarrow{j} \mathrm{Hom}_{\mathrm{cont}}(G_K^{\ell, \mathrm{ab}}, \mathbb{Z}_\ell)$$

which is trivial on k^\times , as k is algebraically. This allows us to interpret j as a map from the projectivization K^\times/k^\times of the k -vector space $(K, +)$ to the “affine” space on the right, which is $\mathrm{Hom}_{\mathrm{cont}}(G_K^{\ell, \mathrm{ab}}, \mathbb{Z}_\ell)$, or even $\mathrm{Hom}_{\mathrm{cont}}(G_K^{\ell, \mathrm{ab}}, \mathbb{F}_p)$. And in particular, if $G_K^{\ell, \mathrm{ab}}$ is very small, then on the right we do really have an affine geometry. Finally, since such maps between projective and affine geometries are of very special shape, BOGOMOLOV–TSCHINKEL show that a liftable non pro-cyclic subgroup Γ of $G_K^{\ell, \mathrm{ab}}$ must contain an element σ which —by duality— defines a *flag function* on K^\times . Strictly speaking, this means that σ is an inertia element to a valuation w with the claimed properties.

It is interesting to remark that as a by-product of the theory of very small pro- ℓ Galois groups, one obtains a p -adic analog of the Artin–Schreier Theorem for the Galois characterization of the real closed fields. The result is:

Theorem (See POP [P0], KOENIGSMANN [Ko1], EFRAT [Ef1]).

Let K be a field having G_K isomorphic to some open subgroup of $G_{\mathbb{Q}_p}$. Then K is p -adically closed, i.e., K is Henselian with respect to a valuation v having divisible value group, and residue field Kv contained and relatively algebraically closed in some finite extension $k|\mathbb{Q}_p$ of \mathbb{Q}_p .

An interesting consequence of the Theorem above is a positive result of the birational Section Conjecture over the p -adics as follows:

Theorem (See POP [P0], KOENIGSMANN [Ko3]).

Let $k|\mathbb{Q}_p$ be a finite extension, and $K|k$ an arbitrary regular field extension. Then for every section $s : G_k \rightarrow G_K$ of the canonical projection $\mathrm{pr}_K : G_K \rightarrow G_k$ one has: The fixed field $K^{(s)}$ of $s(G_k)$ in K^{a} is p -adically closed. Moreover, if v_s is the valuation of $K^{(s)}$ defining it as a p -adically closed field, then $Kv_s = k$.

In particular, if $k = k(X)$ is the function field of a complete k -variety, then every section $s : G_k \rightarrow G_{k(X)}$ is defined by a k -rational point $x_s \in X(k)$. The point x_s is exactly the center of v_s on the complete k -variety X .

This is so far the best un-conditional result concerning the (birational) Section Conjecture we have. But it is not at all clear how to “globalize” such p -adic results in order to get the birational Section Conjecture over number fields.

B) The Raynaud / Pop–Saidi / Tamagawa Theorem

As we have seen at the beginning of Part III, one might/should expect strong anabelian phenomena for curves (maybe even more general varieties) over algebraically closed fields of positive characteristic. Maybe a good hint in that

direction is the fact that *we do not have a description of the algebraic fundamental group of any potentially hyperbolic curve*. Indeed, if from the fundamental group of a curve X we can recover X in a functorial way, then the fundamental group of X must encode the “moduli” for X , thus an information of a completely different nature than crude profinite group theory. See TAMAGAWA [T3], [T4] for more about this conjectural world.

Now let me explain the content of the announced Theorem. Recall that for a complete smooth connected curve X of genus $g \geq 2$ over a field of characteristic 0 one has $\pi_1(\overline{X}) \cong \widehat{\Pi}_g$, thus $\pi_1(\overline{X})$ depends only on g . As mentioned above, in positive characteristic $\pi_1(\overline{X})$ is unknown, and it *depends on the isomorphism type* of \overline{X} . By Grothendieck’s Specialization Theorem, $\pi_1(\overline{X})$ is a quotient of $\widehat{\Gamma}_g$, thus it is topologically finitely generated. In particular, $\pi_1(\overline{X})$ is completely determined by its set of their finite quotients. (Terminology: $\pi_1(\overline{X})$ is a *Pfaffian group*.)

Let $\mathcal{M}_g \rightarrow \mathbb{F}_p$ be the coarse moduli space of proper and smooth curves of genus g in characteristic p . One knows that \mathcal{M}_g is a quasi-projective and geometrically irreducible variety. And if k is an algebraically closed field of characteristic p , then $\mathcal{M}_g(k)$ is the set of isomorphism classes of curves of genus g over k . For $\overline{x} \in \mathcal{M}_g(k)$ let $C_{\overline{x}} \rightarrow k$ be a curve classified by \overline{x} , and let $x \in \mathcal{M}_g$ such that $\overline{x} : k \rightarrow \mathcal{M}_g$ factors through x . We set

$$\pi_1(x) := \pi_1(C_{\overline{x}}),$$

and remark that the structure of $\pi_1(x)$ as a profinite group depends on x only, and not on the concrete geometric point $\overline{x} \in \mathcal{M}_g(k)$ used to define it. In particular, the fundamental group functor gives rise to a map as follows:

$$\pi_1 : \mathcal{M}_g \rightarrow \mathcal{G}, \quad x \mapsto \pi_1(x).$$

To finish our preparation we remark that for points $x, y \in \mathcal{M}_g$ such that x is a specialization of y , by Grothendieck’s Specialization Theorem there exists a surjective continuous homomorphism $\text{sp} : \pi_1(y) \rightarrow \pi_1(x)$. In particular, if η is the generic point of \mathcal{M}_g , then C_η is the generic curve of genus g ; and every point x of \mathcal{M}_g is a specialization of η . For every $x \in \mathcal{M}_g$, there is a surjective homomorphism $\text{sp}_x : \pi_1(\eta) \rightarrow \pi_1(x)$ which is determined up to Galois-conjugacy by the choice of the local ring of x in the algebraic closure of $\kappa(\eta)$. For every $x \in \mathcal{M}_g$ we fix such a map once for all; in particular, if x is a specialization of y , there exists a specialization homomorphism $\text{sp}_{y,x} : \pi_1(y) \rightarrow \pi_1(x)$ such that $\text{sp}_{y,x} \circ \text{sp}_y = \text{sp}_x$.

Theorem (RAYNAUD [R2], POP–SAIDI [P–S], TAMAGAWA [T5]).

For all points $s \neq x$ in \mathcal{M}_g with s closed and specialization of x , the specialization homomorphism $\text{sp}_{x,s} : \pi_1(x) \rightarrow \pi_1(s)$ is not an isomorphism.

More precisely, there exist cyclic étale covers of X_x of order prime to p , which do not have good reduction under the specialization $x \mapsto s$.

As an application one the following answer to a question raised by HARBATER:

Corollary. *There is no non-empty open subset $U \subset \mathcal{M}_g$ such that the isomorphism type of the geometric fundamental group $\pi_1(x)$ is constant on U .*

Concerning the proof of the above Theorem: In the case $g = 2$ the above Theorem was proved by RAYNAUD, by introducing a new kind of *theta divisor* (called by now the Raynaud theta divisor). Using this tool, he showed that given a projective curve $X \rightarrow k_0$ of genus 2, there exist only finitely many curves $X' \rightarrow k_0$ with $\pi_1(X) \cong \pi_1(X')$, see RAYNAUD [R2]. Around the same time POP–SAIDI proposed a way of generalizing RAYNAUD’s result to all genera, by combining the theory of Raynaud’s theta divisor with the results by HRUSHOVSKI [Hr] on the geometric case of the Manin-Mumford Conjecture as follows: First suppose that $g = 2$. Then for points $x_0 \neq x_1$ in \mathcal{M}_g such that x_0 is a specialization of x_1 , it turns out that Raynaud’s Result follows from: *If x_0 is a closed point, then sp_{x_1, x_0} is not an isomorphism.* POP–SAIDI showed in [P–S] that this is the case for arbitrary genera $g > 1$, provide x_0 has some special properties, see loc.cit. Finally, TAMAGAWA [T5] elaborating on the method proposed in [P–S] showed that sp_{x_1, x_0} is not an isomorphism, provided $x \neq x_0$ and x_0 is a closed point.

C) Geometric pro- ℓ birational anabelian Geometry

Here I want to mention the new results concerning some progress on Bogomolov’s Program mentioned at the beginning of Part III. Recall that for a fixed prime number ℓ , we denote by $G_K(\ell)$ the Galois group of a maximal pro- ℓ Galois extension of K . The short spelling out of the story is the following:

Theorem (See POP [P5]).

Let ℓ be a fixed prime number. Consider all the function fields $K|k$ with k algebraic closures of finite fields, $\mathrm{char}(K) \neq \ell$ and $\mathrm{td}(K|k) > 1$. Then there exists a group theoretic recipe by which one can recover any $K|k$ from $G_K(\ell)$ in a functorial way.

Moreover, given a further function field $L|l$ with l an algebraically closed field, every isomorphism $\Phi : G_K(\ell) \rightarrow G_L(\ell)$ is defined by some field isomorphism $\phi : L^i \rightarrow K^i$, and ϕ is unique up to Frobenius twists.

I mention right away that BOGOMOLOV–TSCHINKEL [B–T2] announced a similar result in the case $K = k(X)$ is the function field of a surface X which has trivial (Abelian?) fundamental group. Also, their assertion is somewhat different from the one of the Theorem above, as they deal with the case PGal_K^c in stead of $G_K(\ell)$.

The Theorem above is a far reaching extension of Grothendieck’s birational conjecture in positive characteristic, as it implies the latter one if $\dim(K) > 1$. Second it implies an “arithmetic by pro- ℓ ” form of Grothendieck’s birational anabelian Conjecture.

A general strategy/approach in order to prove birational pro- ℓ anabelian type results was described in Part II of POP [P4], where the *arithmetic by pro- ℓ* case in positive characteristic was considered. Among other things, loc.cit. contains an “abstract Galois theory” (similar to the abstract class field theory), which under certain hypothesis is shown to *originate from geometry*. Such an approach (in the context of finitely generated fields) was suggested to me by DELIGNE (private communication). The new tools needed/developed in the [P5] is a *local theory* similar to the one in [P1], by using the results by KOENIGSMANN [Ko1], EFRAT [Ef2], WARE [Wa], etc., mentioned above. (Naturally, this could be done using BOGOMOLOV–TSCHINKEL [B–T1] too.) And a quite surprising new but very basic fact which is the following: *The set of all the inertia elements in $G_K(\ell)$ is topologically closed in $G_K(\ell)$. And further, the divisorial inertia elements are dense in the set of all the inertia elements.* This is in contrast to the behavior of the set of all the *Frobenius elements* of finitely generated fields, which is dense in the absolute Galois group of such fields.

Some major open Questions/Problems:

Q1: Let k be an algebraically closed field of positive characteristic. Can one recover $\text{td}(k)$ from $\pi_1(\mathbb{A}_k^1)$?

Q2: For k as above, give a non tautological description of $\pi_1(\mathbb{A}_k^1)$.

Q3: For k as above, give a non tautological description of $\pi_1(X)$ and/or $\pi_1^{\text{t}}(X)$ for some hyperbolic curve $X \rightarrow k$.

Q4: Give an algebraic proof of the fact that $\pi_1(\mathbb{P}_{\mathbb{C}}^1 \setminus \{0, 1, \infty\})$ is generated by inertia elements c_0, c_1, c_{∞} over $0, 1, \infty$ with a single relation $c_0 c_1 c_{\infty} = 1$.

Q5: Give a proof of the Hom-form of Grothendieck’s birational anabelian Conjecture in positive characteristic.

Q6: Prove the Isom-form and/or the Hom-form of the anabelian Conjecture for curves over finite fields.

Q7: Let $x_1 \neq x_0$ in $\mathcal{M}_g \rightarrow \mathbb{F}_p$ be such that x_0 is a specialization of x_1 . Show that $\text{sp}_{x_1 x_0} : \pi_1(x_1) \rightarrow \pi_1(x_0)$ is not an isomorphism.

Q8: Prove the Section Conjecture, say over number fields and/or p -adic fields...

REFERENCES

- [Bo] BOGOMOLOV, F. A., *On two conjectures in birational algebraic geometry*, in Algebraic Geometry and Analytic Geometry, ICM-90 Satellite Conference Proceedings, ed A. Fujiki et al, Springer Verlag Tokyo 1991.
- [B–T1] BOGOMOLOV, F. A., and TSCHINKEL, Y. *Commuting elements in Galois groups of function fields*, in: Motives, Polylogs and Hodge Theory, International Press 2002, 75–120.
- [B–T2] BOGOMOLOV, F. A., and TSCHINKEL, Y. *Reconstruction of function fields*, Manuscript, March 2003.
- [BOU] BOURBAKI, *Algèbre commutative*, Hermann Paris 1964.

- [D1] DELIGNE, P., *Le groupe fondamental de la droite projective moins trois points*, in: Galois groups over \mathbf{Q} , Math. Sci. Res. Inst. Publ. **16**, 79–297, Springer 1989.
- [Ef1] EFRAT, I., *Construction of valuations from K -theory*, Mathematical Research Letters **6** (1999), 335–344.
- [Ef2] ———, *Recovering higher global and local fields from Galois groups—an algebraic approach*, Invitation to higher local fields (Münster, 1999), 273–279 (electronic), Geom., Topol. Monogr. **3**, Geom. Topol. Publ., Coventry, 2000.
- [E–K] ENGLER, A. J. and KOENIGSMANN, J., *Abelian subgroups of pro- p Galois groups*, Trans. AMS **350** (1998), no. 6, 2473–2485.
- [F1] FALTINGS, G., *p -adic Hodge Theory*, Journal AMS **1** (1988), 255–299.
- [F2] ———, *Hodge–Tate structures and modular forms*, Math. Annalen **278** (1987), 133–149.
- [F3] ———, *Curves and their fundamental groups [following Grothendieck, Tamagawa, Mochizuki]*, Séminaire Bourbaki, Vol 1997–98, Exposé 840, Mars 1998. [F–J] FRIED, M. D. and JARDEN, M., *Field Arithmetic*, in: Ergebnisse der Mathematik und ihre Grenzgebiete, 3. Folge, Vol 11, Springer Verlag 2004.
- [G1] GROTHENDIECK, A., Letter to Faltings, June 1983. See [GGA].
- [G2] ———, *Esquisse d’un programme*, 1984. See [GGA].
- [GGA] Geometric Galois Actions I, LMS LNS Vol **242**, eds L. Schneps – P. Lochak, Cambridge Univ. Press 1998.
- [Ha1] HARBATER, D., *Abhyankar’s conjecture on Galois groups over curves*, Invent. Math. **117** (1994), 1–25.
- [Ha2] ———, *Fundamental groups of curves in characteristic p* , in: Proceedings of the ICM (Zürich, 1994), Birkhser, Basel, 1995, 656–666.
- [Hr] HRUSHOVSKI, E., *The Mordell–Lang conjecture for function fields*, Journal AMS **9** (1996), 667–690.
- [Ih] IHARA, Y., *On beta and gamma functions associated with the Grothendieck–Teichmüller group II*, J. reine angew. Math. **527** (2000), 1–11.
- [I–M] IHARA, Y. and MATSUMOTO, M., *On Galois actions on profinite completions of braid groups*, in: Recent developments in the inverse Galois problem (Seattle, WA, 1993), 173–200, Contemp. Math. **186** AMS Providence, RI, 1995.
- [I–N] IHARA, Y., and NAKAMURA, H., *Some illustrative examples for anabelian geometry in high dimensions*, in: Geometric Galois actions I, 127–138, LMS LNS **242**, Cambridge Univ. Press, Cambridge, 1997.
- [Ik] IKEDA, M., *Completeness of the absolute Galois group of the rational number field*, J. reine angew. Math. **291** (1977), 1–22.
- [J–W] JANNSEN, U. and WINGBERG, K., *Die Struktur der absoluten Galoisgruppe p -adischer Zahlkörper*, Invent. Math. **70** (1982/83), no. 1, 71–98.
- [J] DE JONG, A. J., *Families of curves and alterations*, Annales de l’institute Fourier, **47** (1997), pp. 599–621.
- [Ka] KATO, K., *Logarithmic Structures of Fontaine–Illusie*, Proceedings of the First JAMI Conference, Johns Hopkins Univ. Press (1990), 191–224.
- [K–L] KATZ, N., and LANG, S., *Finiteness theorems in geometric class field theory*, with an Appendix by K. Ribet, Enseign. Math. **27** (1981), 285–319.
- [Ko1] KOENIGSMANN J., *From p -rigid elements to valuations (with a Galois-characterization of p -adic fields). With an Appendix by Florian Pop*, J. Reine Angew. Math. **465** (1995), 165–182.
- [Ko2] ———, *Solvable absolute Galois groups are metabelian*, Invent. Math. **144** (2001), no. 1, 1–22.
- [Ko3] ———, *On the Section Conjecture*, manuscript, 2002.
- [Ko] KOCH, H., *Die Galoissche Theorie der p -Erweiterungen*, Math. Monographien **10**, Berlin 1970.

- [La] LANG, S., Algebra, Springer-Verlag 2001.
- [L-T] LANG, S. and TATE, J., Principal homogeneous spaces over Abelian varieties, Am. J. Math. **80** (1958), 659–684.
- [Mzk1] MOCHIZUKI, SH., *The profinite Grothendieck Conjecture for closed hyperbolic curves over number fields*, J. Math. Sci. Univ Tokyo **3** (1966), 571–627.
- [Mzk2] ———, *A version of the Grothendieck conjecture for p -adic local fields*, Internat. J. Math. no. 4, **8** (1997), 499–506.
- [Mzk3] ———, *The local pro- p Grothendieck conjecture for hyperbolic curves*, Invent. Math. **138** (1999), 319–423.
- [Mzk4] ———, *The absolute anabelian geometry of hyperbolic curves*, Galois theory and modular forms, 77–122, Dev. Math., 11, Kluwer Acad. Publ., Boston, MA, 2004.
- [N] NAGATA, M., *A theorem on valuation rings and its applications*, Nagoya Math. J. **29** (1967), 85–91.
- [Na1] NAKAMURA, H., *Galois rigidity of the étale fundamental groups of punctured projective lines*, J. reine angew. Math. **411** (1990) 205–216.
- [Na2] ———, *Galois rigidity of algebraic mappings into some hyperbolic varieties*, Int. J. Math. **4** (1993), 421–438.
- [N1] NEUKIRCH, J., *Über eine algebraische Kennzeichnung der Henselkörper*, J. reine angew. Math. **231** (1968), 75–81.
- [N2] ———, *Kennzeichnung der p -adischen und endlichen algebraischen Zahlkörper*, Inventiones math. **6** (1969), 269–314.
- [N3] ———, *Kennzeichnung der endlich-algebraischen Zahlkörper durch die Galoisgruppe der maximal auflösbaren Erweiterungen*, J. für Math. **238** (1969), 135–147.
- [O] ODA, T., *A note on ramification of the Galois representation of the fundamental group of an algebraic curve I*, J. Number Theory (1990) 225–228.
- [Pa] PARSHIN, A. N., *Finiteness Theorems and Hyperbolic Manifolds*, in: The Grothendieck Festschrift III, ed P. Cartier et al, PM Series Vol 88, Birkhäuser Boston Basel Berlin 1990.
- [P0] POP, F., *Galoissche Kennzeichnung p -adisch abgeschlossener Körper*, Dissertation, Heidelberg 1986.
- [P1] ———, *On Grothendieck’s conjecture of birational anabelian Geometry*, Ann. of Math. **139** (1994), 145–182.
- [P2] ———, *On Grothendieck’s conjecture of birational anabelian geometry II*, preprint, 1995
- [P3] ———, *Alterations and birational anabelian geometry*, in Resolution of Singularities (Obergrugl, Austria, 1997), PM Series 181, Birkhauser, Basel, 2000.
- [P4] ———, *The birational anabelian conjecture — r e v i s i t e d —*, Manuscript, Princeton/Bonn 2002. See <http://www.math.leidenuniv.nl/gtem/view.php>
- [P5] ———, *Pro- ℓ birational anabelian geometry over alg. closed fields I*, Manuscript, Bonn 2003. See http://arxiv.org/PS_cache/math/pdf/0307/0307076
- [P6] ———, *Glimpses of Grothendieck’s anabelian geometry*, in: Geometric Galois Actions I, LMS LNS Vol **242**, 133–126; eds L. Schneps – P. Lochak, Cambridge Univ. Press 1998.
- [Po] POP, F., *Étale Galois covers of affine, smooth curves*, Invent. Math. **120** (1995), 555–578.
- [P-S] POP, F., and SAIDI, M. *On the specialization homomorphism of fundamental groups of curves in positive characteristic*, in: Galois groups and Fundamental groups (ed. Schneps), MSRI Pub. Series **41**, 2003, pp.107–118.
- [R1] RAYNAUD, M., *Revêtements des courbes en caractéristique $p > 0$ et ordinarité*, Compositio Math. **123** (2000), no. 1, 73–88.

- [R2] ———, *Sur le groupe fondamental d'une courbe complète en caractéristique $p > 0$* , in: Arithmetic fundamental groups and noncommutative algebra (Berkeley, CA, 1999), 335–351, Proc. Sympos. Pure Math., 70, AMS, Providence, RI, 2002.
- [Ro] ROQUETTE, P., *Some tendencies in contemporary algebra*, in: Perspectives in Math. Anniversary of Oberwolfach 1984, Basel 1984, 393–422.
- [Sa] SAIDI, M., *Revêtements modérés et groupe fondamental de graphes de groupes*, Compositio Math. **107** (1997), 319–338.
- [S1] SERRE, J.-P., Cohomologie Galoisienne, 5th ed., LNM 5, Springer Verlag, Berlin, 1994
- [S2] ———, *Zeta and L functions*, in: Arithmetical Algebraic Geometry (Proc. Conf. Purdue Univ., 1963), pp.82–92; Harper & Row, New York 1965.
- [S3] ———, *Corps locaux*, Herman Paris 1962.
- [Sp] SPIESS, M., *An arithmetic proof of Pop's Theorem concerning Galois groups of function fields over number fields*, J. reine angew. Math. **478** (1996), 107–126.
- [St1] STIX, J., *Affine anabelian curves in positive characteristic*, Compositio Math. **134** (2002), no. 1, 75–85
- [St2] ———, *Projective anabelian curves in positive characteristic and descent theory for log-étale covers*, Dissertation, Bonner Mathematische Schriften **354**, Universit Bonn, Math. Inst. Bonn, 2002; see <http://www.math.uni-bonn.de/people/stix>
- [T1] TAMAGAWA, A., *The Grothendieck conjecture for affine curves*, Compositio Math. **109** (1997), 135–194.
- [T2] ———, *On the fundamental groups of curves over algebraically closed fields of characteristic > 0* , Internat. Math. Res. Notices 1999, no. 16, 853–873.
- [T3] ———, *On the tame fundamental groups of curves over algebraically closed fields of characteristic > 0* , in: Galois groups and fundamental groups, 47–105, MSRI Publ., **41**, Cambridge Univ. Press, Cambridge, 2003.
- [T4] ———, *Fundamental groups and geometry of curves in positive characteristic*, in: Arithmetic fundamental groups and noncommutative algebra (Berkeley, CA, 1999), 297–333, Proc. Sympos. Pure Math., 70, AMS Providence, RI, 2002.
- [T5] ———, *Finiteness of isomorphism classes of curves in positive characteristic with prescribed fundamental groups*, J. Algebraic Geom. **13** (2004), no. 4, 675–724.
- [U1] UCHIDA, K., *Isomorphisms of Galois groups of algebraic function fields*, Ann. of Math. **106** (1977), 589–598.
- [U2] ———, *Isomorphisms of Galois groups of solvably closed Galois extensions*, Tôhoku Math. J. **31** (1979), 359–362.
- [U3] ———, *Homomorphisms of Galois groups of solvably closed Galois extensions*, Journal Math. Soc. Japan **33**, No.4, 1981.
- [Ta1] TATE, J., *Relations between K_2 and Galois cohomology*, Invent. Math. **36** (1976), 257–274.
- [Ta2] ———, *p -divisible Groups*, Proceeding of a Conference on local fields, Driebergen, Springer Verlag 1969, 158–183.
- [W] WARE, R., *Valuation Rings and rigid Elements in Fields*, Can. J. Math. **33** (1981), 1338–1355.