

An Introduction to Trimmed Serendipity Finite Element Spaces

Andrew Gillette - University of Arizona

*joint work with
Tyler Kloefkorn, AAAS STP Fellow, hosted at NSF*




Table of Contents

- 1 Four well-known families of finite elements
- 2 Trimmed serendipity spaces: a new, smaller family of elements
- 3 A computational basis for trimmed serendipity spaces

- 1 Four well-known families of finite elements
- 2 Trimmed serendipity spaces: a new, smaller family of elements
- 3 A computational basis for trimmed serendipity spaces

Classification of conforming methods

Conforming finite element method types can be broadly classified by three integers:

- $n \rightarrow$ the spatial dimension of the domain
- $r \rightarrow$ the order of error decay
- $k \rightarrow$ the differential form order of the solution space


An element type is defined in part by its **degrees of freedom**. Typically:

the more degrees of freedom, the greater the computational cost of the method

Ex: $\mathcal{Q}_1^- \Lambda^2(\square_3)$ is an element for


- $n = 3 \rightarrow$ domains in $\mathbb{R}^3$
- $r = 1 \rightarrow$ linear order of error decay
- $k = 2 \rightarrow$ conformity in $\Lambda^2(\mathbb{R}^3) \rightsquigarrow H(\text{div})$

$\mathcal{Q}_1^- \Lambda^2(\square_3)$ is part of the $\mathcal{Q}^-$ 'column' of elements,
is defined on geometry $\square_3$ (i.e. a cube),
has a **6** dimensional space of test functions,
and has an associated set of **6** degrees of freedom
that are unisolvent for the test function space.



The ‘Periodic Table of the Finite Elements’

ARNOLD, LOGG, “Periodic table of the finite elements,” *SIAM News*, 2014.



Classification of many common conforming finite element types.

- $n \rightarrow$ Domains in $\mathbb{R}^2$ (top half) and in $\mathbb{R}^3$ (bottom half)
- $r \rightarrow$ Order 1, 2, 3 of error decay (going down columns)
- $k \rightarrow$ Conformity type $k = 0, \dots, n$ (going across a row)



Geometry types: Simplices (left half) and cubes (right half).

An abbreviated reading list (50 years of theory!)

- RAVIART, THOMAS, "A mixed finite element method for 2nd order elliptic problems" *Lecture Notes in Mathematics*, 1977 ← 3172 citations, including 150 from 2017!
- NÉDÉLEC, "Mixed finite elements in $\mathbb{R}^3$," *Numerische Mathematik*, 1980
- BREZZI, DOUGLAS JR., MARINI, "Two families of mixed finite elements for second order elliptic problems," *Numerische Mathematik*, 1985
- NÉDÉLEC, "A new family of mixed finite elements in $\mathbb{R}^3$," *Numerische Mathematik*, 1986
- ARNOLD, FALK, WINTHER "Finite element exterior calculus, homological techniques, and applications," *Acta Numerica*, 2006
- CHRISTIANSEN, "Stability of Hodge decompositions in finite element spaces of differential forms in arbitrary dimension," *Numerische Mathematik*, 2007
- ARNOLD, FALK, WINTHER "Finite element exterior calculus: from hodge theory to numerical stability," *Bulletin of the AMS*, 2010
- ARNOLD, AWANOU "The serendipity family of finite elements ", *Found. Comp Math*, 2011
- ARNOLD, AWANOU "Finite element differential forms on cubical meshes", *Math Comp.*, 2013
- ARNOLD, BOFFI, BONIZZONI "Finite element differential forms on curvilinear meshes and their approximation properties," *Numerische Mathematik*, 2014

Stable pairs of elements for mixed methods


Picking elements from the table for a mixed method for the Poisson problem:




Example and images on right from:

ARNOLD, FALK, WINTHER "Finite Element Exterior Calculus. . ." *Bulletin of the AMS*, 47:2, 2010.


Method selection and cochain complexes



$\subset H(\text{div})$



$\subset L^2$



Provably stable method

converges to
 $u = x(1-x)y(1-y)$

Stable pairs of elements for mixed Hodge-Laplacian problems are found by choosing consecutive spaces in compatible discretizations of the L^2 deRham Diagram.

$$H^1 \xrightarrow[\text{grad}]{\nabla} H(\text{curl}) \xrightarrow[\text{curl}]{\nabla \times} H(\text{div}) \xrightarrow[\text{div}]{\nabla \cdot} L^2$$

vector Poisson

σ

μ

Maxwell's eqn's

h

b

Darcy / Poisson

u

p


The Periodic Table of Finite Elements lets us 'read off' stable pairs visually.

Stable pairs for tetrahedral meshes



Problem: Darcy / Poisson
Dimension: $n = 3$
Mesh type: tetrahedral
Convergence: quadratic ($r = 2$)

Stable pairs for tetrahedral meshes




Problem: Darcy / Poisson
Dimension: $n = 3$
Mesh type: tetrahedral
Convergence: cubic ($r = 3$)

Stable pairs for tetrahedral meshes



Problem: Maxwell's
Dimension: $n = 3$
Mesh type: tetrahedral
Convergence: quadratic ($r = 2$)

Stable pairs for tetrahedral meshes




Problem: Maxwell's
Dimension: $n = 3$
Mesh type: tetrahedral
Convergence: cubic ($r = 3$)

Stable pairs for tetrahedral meshes




Problem: vector Poisson
Dimension: $n = 3$
Mesh type: tetrahedral
Convergence: quadratic ($r = 2$)

Stable pairs for tetrahedral meshes




Problem: vector Poisson
Dimension: $n = 3$
Mesh type: tetrahedral
Convergence: cubic ($r = 3$)

Stable pairs for cubical meshes





Problem: Darcy / Poisson
Dimension: $n = 3$
Mesh type: cubes
Convergence: quadratic ($r = 2$)

Stable pairs for cubical meshes




Problem: Darcy / Poisson
Dimension: $n = 3$
Mesh type: cubes
Convergence: cubic ($r = 3$)

Stable pairs for cubical meshes




Problem: Maxwell's
Dimension: $n = 3$
Mesh type: cubes
Convergence: quadratic ($r = 2$)

Stable pairs for cubical meshes




Problem:	Maxwell's
Dimension:	$n = 3$
Mesh type:	cubes
Convergence:	cubic ($r = 3$)

Stable pairs for cubical meshes



Problem: vector Poisson
Dimension: $n = 3$
Mesh type: cubes
Convergence: quadratic ($r = 2$)

Stable pairs for cubical meshes



Problem: vector Poisson
Dimension: $n = 3$
Mesh type: cubes
Convergence: cubic ($r = 3$)

Exact cochain complexes found in the table

On an n -simplex in $\mathbb{R}^n$:



$$\mathcal{P}_r^- \Lambda^0 \rightarrow \mathcal{P}_r^- \Lambda^1 \rightarrow \cdots \rightarrow \mathcal{P}_r^- \Lambda^{n-1} \rightarrow \mathcal{P}_r^- \Lambda^n \quad \text{‘trimmed’ polynomials}$$

$$\mathcal{P}_r \Lambda^0 \rightarrow \mathcal{P}_{r-1} \Lambda^1 \rightarrow \cdots \rightarrow \mathcal{P}_{r-n+1} \Lambda^{n-1} \rightarrow \mathcal{P}_{r-n} \Lambda^n \quad \text{polynomials}$$

On an n -dimensional cube in $\mathbb{R}^n$:

$$\mathcal{Q}_r^- \Lambda^0 \rightarrow \mathcal{Q}_r^- \Lambda^1 \rightarrow \cdots \rightarrow \mathcal{Q}_r^- \Lambda^{n-1} \rightarrow \mathcal{Q}_r^- \Lambda^n \quad \text{tensor product}$$

$$\mathcal{S}_r \Lambda^0 \rightarrow \mathcal{S}_{r-1} \Lambda^1 \rightarrow \cdots \rightarrow \mathcal{S}_{r-n+1} \Lambda^{n-1} \rightarrow \mathcal{S}_{r-n} \Lambda^n \quad \text{serendipity}$$









The ‘minus’ spaces proceed across rows of the PToFE (r is fixed) while the ‘regular’ spaces proceed along diagonals (r decreases)

Mysteriously, the degree of freedom count for mixed methods from the $\mathcal{P}_r^-$ spaces is smaller than those from the $\mathcal{P}_r$ spaces, while the opposite is true for the $\mathcal{Q}_r^-$ and $\mathcal{S}_r$ spaces.

Outline

- 1 Four well-known families of finite elements
- 2 **Trimmed serendipity spaces: a new, smaller family of elements**
- 3 A computational basis for trimmed serendipity spaces

The 5th column: Trimmed serendipity spaces



A new column for the PToFE:
the **trimmed serendipity** elements.

$\mathcal{S}_r^- \Lambda^k(\square_n)$ denotes
approximation order r ,
subset of k -form space $\Lambda^k(\Omega)$,
use on meshes of n -dim'l cubes.

Defined for any $n \geq 1$, $0 \leq k \leq n$, $r \geq 1$

Identical or analogous properties to all the
other columns in the table.

The advantage of the $\mathcal{S}_r^- \Lambda^k$ spaces is that
they have fewer degrees of freedom for mixed
methods than their tensor product and
serendipity counterparts.

Key properties of the trimmed serendipity spaces

$$\mathcal{Q}_r^- \Lambda^0 \rightarrow \mathcal{Q}_r^- \Lambda^1 \rightarrow \cdots \rightarrow \mathcal{Q}_r^- \Lambda^{n-1} \rightarrow \mathcal{Q}_r^- \Lambda^n \quad \text{tensor product}$$

$$\mathcal{S}_r \Lambda^0 \rightarrow \mathcal{S}_{r-1} \Lambda^1 \rightarrow \cdots \rightarrow \mathcal{S}_{r-n+1} \Lambda^{n-1} \rightarrow \mathcal{S}_{r-n} \Lambda^n \quad \text{serendipity}$$

$$\mathcal{S}_r^- \Lambda^0 \rightarrow \mathcal{S}_r^- \Lambda^1 \rightarrow \cdots \rightarrow \mathcal{S}_r^- \Lambda^{n-1} \rightarrow \mathcal{S}_r^- \Lambda^n \quad \text{trimmed serendipity}$$

Subcomplex: $d\mathcal{S}_r^- \Lambda^k \subset \mathcal{S}_r^- \Lambda^{k+1}$

Exactness: The above sequence is exact.
i.e. the image of incoming map = kernel of outgoing map

Inclusion: $\mathcal{S}_r \Lambda^k \subset \mathcal{S}_{r+1}^- \Lambda^k \subset \mathcal{S}_{r+1} \Lambda^k$

Trace: $\text{tr}_f \mathcal{S}_r^- \Lambda^k(\mathbb{R}^n) \subset \mathcal{S}_r^- \Lambda^k(f), \quad \text{for any } (n-1)\text{-hyperplane } f \text{ in } \mathbb{R}^n$

Special cases:

$$\begin{aligned} \mathcal{S}_r^- \Lambda^0 &= \mathcal{S}_r \Lambda^0 \\ \mathcal{S}_r^- \Lambda^n &= \mathcal{S}_{r-1} \Lambda^n \\ \mathcal{S}_r^- \Lambda^k + d\mathcal{S}_{r+1} \Lambda^{k-1} &= \mathcal{S}_r \Lambda^k. \end{aligned}$$

Replace 'S' by 'P' $\rightsquigarrow$ key properties about the first two columns for $\mathcal{P}_r^- \Lambda^k$ and $\mathcal{P}_r \Lambda^k$!

Dimension count and comparison

Formula for counting degrees of freedom of $\mathcal{S}_r^- \Lambda^k(\square_n)$:

$$\sum_{d=k}^{\min\{n, \lfloor r/2 \rfloor + k\}} 2^{n-d} \binom{n}{d} \left(\binom{r-d+2k-1}{r-d+k-1} \binom{r-d+k-1}{d-k} + \binom{r-d+2k}{k} \binom{r-d+k-1}{d-k-1} \right)$$

	k	r=1	2	3	4	5	6	7
n=2	0	4	8	12	17	23	30	38
	1	4	10	17	26	37	50	65
	2	1	3	6	10	15	21	28
n=3	0	8	20	32	50	74	105	144
	1	12	36	66	111	173	255	360
	2	6	21	45	82	135	207	301
	3	1	4	10	20	35	56	84
n=4	0	16	48	80	136	216	328	480
	1	32	112	216	392	656	1036	1563
	2	24	96	216	422	746	1227	1910
	3	8	36	94	200	375	644	1036
	4	1	5	15	35	70	126	210

Mixed Method dimension comparison 1

Mixed method for Darcy problem:

$$\begin{aligned}\mathbf{u} + K \nabla p &= 0 \\ \operatorname{div} \mathbf{u} - f &= 0\end{aligned}$$

We compare degree of freedom counts among the three families for use on meshes of affinely-mapped squares or cubes, when a conforming method with (at least) order r decay in the approximation of p , $\mathbf{u}$, and $\operatorname{div} \mathbf{u}$ is desired.

Total # of degrees of freedom on a square ($n = 2$):

r	$ \mathcal{Q}_r^- \Lambda^1 + \mathcal{Q}_r^- \Lambda^2 $	$ \mathcal{S}_r \Lambda^1 + \mathcal{S}_{r-1} \Lambda^2 $	$ \mathcal{S}_r^- \Lambda^1 + \mathcal{S}_r^- \Lambda^2 $
1	4+1 = 5	8+1 = 9	4+1 = 5
2	12+4 = 16	14+3 = 17	10+3 = 13
3	24+9 = 33	22+6 = 28	17+6 = 23

Total # of degrees of freedom on a cube ($n = 3$):

r	$ \mathcal{Q}_r^- \Lambda^2 + \mathcal{Q}_r^- \Lambda^3 $	$ \mathcal{S}_r \Lambda^2 + \mathcal{S}_{r-1} \Lambda^3 $	$ \mathcal{S}_r^- \Lambda^2 + \mathcal{S}_r^- \Lambda^3 $
1	6+1 = 7	18+1 = 19	6+1 = 7
2	36+8 = 44	39+4 = 43	21+4 = 25
3	108+27 = 135	72+10 = 82	45+10 = 55

Mixed Method dimension comparison 2

Mixed method for Darcy problem:

$$\begin{aligned} \mathbf{u} + K \nabla p &= 0 \\ \operatorname{div} \mathbf{u} - f &= 0 \end{aligned}$$

The number of interior degrees of freedom is reduced from tensor product, to serendipity, to trimmed serendipity:

of **interior** degrees of freedom on a square ($n = 2$):

r	$ \mathcal{Q}_r^- \Lambda_0^1 + \mathcal{Q}_r^- \Lambda_0^2 $	$ \mathcal{S}_r \Lambda_0^1 + \mathcal{S}_{r-1} \Lambda_0^2 $	$ \mathcal{S}_r^- \Lambda_0^1 + \mathcal{S}_r^- \Lambda_0^2 $
1	$0+1 = 1$	$0+1 = 1$	$0+1 = 1$
2	$4+4 = 8$	$2+3 = 5$	$2+3 = 5$
3	$12+9 = 21$	$6+6 = 12$	$5+6 = 11$

of **interior** degrees of freedom on a cube ($n = 3$):

r	$ \mathcal{Q}_r^- \Lambda_0^2 + \mathcal{Q}_r^- \Lambda_0^3 $	$ \mathcal{S}_r \Lambda_0^2 + \mathcal{S}_{r-1} \Lambda_0^3 $	$ \mathcal{S}_r^- \Lambda_0^2 + \mathcal{S}_r^- \Lambda_0^3 $
1	$0+1 = 1$	$0+1 = 1$	$0+1 = 1$
2	$12+8 = 20$	$3+4 = 7$	$3+4 = 7$
3	$54+27 = 81$	$12+10 = 22$	$9+10 = 19$

Mixed Method dimension comparison 3

Mixed method for Darcy problem:

$$\begin{aligned} \mathbf{u} + K \nabla p &= 0 \\ \operatorname{div} \mathbf{u} - f &= 0 \end{aligned}$$

Assuming interior degrees of freedom could be dealt with efficiently (e.g. by static condensation), trimmed serendipity elements *still* have the fewest DoFs:

of **interface** (edge) degrees of freedom on a square ($n = 2$):

r	$ \mathcal{Q}_r^- \Lambda^1(\partial \square_2) $	$ S_r \Lambda^1(\partial \square_2) $	$ S_r^- \Lambda^1(\partial \square_2) $
1	4	8	4
2	8	12	8
3	12	16	12

of **interface** (edge+face) degrees of freedom on a cube ($n = 3$):

r	$ \mathcal{Q}_r^- \Lambda^2(\partial \square_3) $	$ S_r \Lambda^2(\partial \square_3) $	$ S_r^- \Lambda^2(\partial \square_3) $
1	6	18	6
2	24	36	18
3	54	60	36

Outline

- 1 Four well-known families of finite elements
- 2 Trimmed serendipity spaces: a new, smaller family of elements
- 3 A computational basis for trimmed serendipity spaces

Decomposition by polynomial subspace

$\mathcal{S}_r^- \Lambda^k(\square_n)$ is a space of differential k -forms whose coefficients are polynomials in $\mathbb{R}^n$.

$$\mathcal{S}_r^- \Lambda^k = \mathcal{P}_r^- \Lambda^k \oplus \mathcal{J}_r \Lambda^k \oplus d\mathcal{J}_r \Lambda^{k-1}$$

Polynomial coefficients in each summand:

$\mathcal{P}_r^- \Lambda^k$: anything up to degree $r - 1$ and some degree r

$\mathcal{J}_r \Lambda^k$: certain polynomials whose degree is between $r+1$ and $r+n-k-1$

$d\mathcal{J}_r \Lambda^{k-1}$: certain polynomials whose degree is between r and $r+n-k-2$

The “regular” serendipity space has an analogous decomposition:

$$\mathcal{S}_r \Lambda^k = \mathcal{P}_r \Lambda^k \oplus \mathcal{J}_r \Lambda^k \oplus d\mathcal{J}_{r+1} \Lambda^{k-1}$$

This decomposition provides a direct sum into some precise but elaborate subspaces:

$$\mathcal{J}_r \Lambda^k(\mathbb{R}^n) := \sum_{l \geq 1} \kappa \mathcal{H}_{r+l-1, l} \Lambda^{k+1}(\mathbb{R}^n),$$

$$\text{where } \mathcal{H}_{r, l} \Lambda^k(\mathbb{R}^n) := \{ \omega \in \mathcal{H}_r \Lambda^k(\mathbb{R}^n) \mid \text{ldeg } \omega \geq l \},$$

$$\text{where } \text{ldeg}(x^\alpha dx_\sigma) := \#\{i \in \sigma^* : \alpha_i = 1\}.$$

Decomposition by Cubical Geometry

We can also decompose $\mathcal{S}_r^- \Lambda^k(\square_n)$ by the subspace of “zero trace”:

$$\mathcal{S}_r^- \Lambda^k = \mathcal{S}_r^- \Lambda_{\mathbf{0}}^k \oplus \left(\mathcal{S}_r^- \Lambda_{\mathbf{0}}^k \right)^\perp$$

We use this decomposition to prove that $\mathcal{S}_r^- \Lambda^\bullet(\square_n)$ is a **minimal compatible finite element system** containing $\mathcal{P}_{r-1} \Lambda^\bullet(\square_n)$.

A computational basis for $\mathcal{S}_r^- \Lambda_{\mathbf{0}}^k$ would aid in the construction of bases for $\mathcal{S}_r^- \Lambda^k$.

Building such a basis is non-trivial. Consider

$$\alpha := \begin{array}{l} (z-1)(y^2-1) \, dx \\ + y(x+1)(z-1) \, dy \\ + (x+1)(y^2-1) \, dz \end{array} \qquad \beta := \begin{array}{l} (z-1)(y^2-1) \, dx \\ - 2y(x+1)(z-1) \, dy \\ + (x+1)(y^2-1) \, dz \end{array}$$

Both α and β have a natural association to the approximation of y on the edge $\{x=1, z=-1\}$, and both are elements of $\mathcal{Q}_2 \Lambda^1(\square_3)$. But **only** β is in $\mathcal{S}_1 \Lambda^1(\square_3)$!

Decompositions shared insight

Why is $\beta \in \mathcal{S}_1 \Lambda^1(\square_3)$?

$$\begin{aligned} \beta &= \begin{aligned} &(z-1)(y^2-1) dx \\ &-2y(x+1)(z-1) dy \\ &+(x+1)(y^2-1) dz \end{aligned} \\ &= \underbrace{\begin{aligned} &y^2 z dx & -y^2 dx & 0 dx \\ &-2xyz dy & 2xy dy & -2yz dy \\ &xy^2 dz & 0 dz & y^2 dz \end{aligned}}_{\text{basis elements for } d\mathcal{J}_2\Lambda^0} + \underbrace{\begin{aligned} &(-z+1) dx \\ &2y dy \\ &(-x-1) dz \end{aligned}}_{\in \mathcal{P}_1\Lambda^1}. \end{aligned}$$

$$\beta \in d\mathcal{J}_2\Lambda^0 \oplus \mathcal{P}_1\Lambda^1 \subset \mathcal{S}_1\Lambda^1$$

From the polynomial subspace decompositions:

$$\beta \in d\mathcal{J}_2\Lambda^0 \oplus \mathcal{P}_1\Lambda^1 \subset \mathcal{S}_2^-\Lambda^1$$

Full report on this approach coming soon!

Acknowledgments

Related Publication

G., Kloefkorn “Trimmed Serendipity Finite Element Differential Forms”
Mathematics of Computation, to appear. See `arXiv:1607.00571`

Related talk

Victoria Sanders, University of Arizona
*Using Sage to Create Lists of Shape Functions for Trimmed Serendipity and
Serendipity Finite Elements*
Saturday 3:00 p.m. Room 19

Research Funding

Supported in part by the National Science Foundation grant DMS-1522289.

Slides and Pre-prints

`http://math.arizona.edu/~agillette/`

Thanks to the organizers for the invitation to speak!