

Methods of Applied Mathematics
Second Semester Lecture Notes

William G. Faris

January 2, 2009

Contents

1	Distributions	7
1.1	Properties of distributions	7
1.2	Mapping distributions	9
1.3	Radon measures	10
1.4	Approximate delta functions	10
1.5	Problems	11
1.6	Tempered distributions	12
1.7	Poisson equation	14
1.8	Diffusion equation	15
1.9	Wave equation	16
1.10	Homogeneous solutions of the wave equation	17
1.11	Problems	18
1.12	Answers to first two problems	19
2	Bounded Operators	21
2.1	Introduction	21
2.2	Bounded linear operators	21
2.3	Compact operators	24
2.4	Hilbert-Schmidt operators	27
2.5	Problems	28
2.6	Finite rank operators	29
2.7	Problems	30
3	Densely Defined Closed Operators	33
3.1	Introduction	33
3.2	Subspaces	33
3.3	Graphs	34
3.4	Operators	35
3.5	The spectrum	35
3.6	Spectra of inverse operators	36
3.7	Problems	37
3.8	Self-adjoint operators	38
3.9	First order differential operators with a bounded interval: point spectrum	38

3.10	Spectral projection and reduced resolvent	40
3.11	Generating second-order self-adjoint operators	41
3.12	First order differential operators with a semi-infinite interval: residual spectrum	42
3.13	First order differential operators with an infinite interval: contin- uous spectrum	42
3.14	Problems	43
3.15	A pathological example	44
4	Normal operators	45
4.1	Spectrum of a normal operator	45
4.2	Problems	46
4.3	Variation of parameters and Green's functions	47
4.4	Second order differential operators with a bounded interval: point spectrum	48
4.5	Second order differential operators with a semibounded interval: continuous spectrum	49
4.6	Second order differential operators with an infinite interval: con- tinuous spectrum	50
4.7	The spectral theorem for normal operators	50
4.8	Examples: compact normal operators	52
4.9	Examples: translation invariant operators and the Fourier trans- form	52
4.10	Examples: Schrödinger operators	53
4.11	Subnormal operators	54
4.12	Examples: forward translation invariant operators and the Laplace transform	56
4.13	Quantum mechanics	58
4.14	Problems	59
5	Calculus of Variations	61
5.1	The Euler-Lagrange equation	61
5.2	A conservation law	62
5.3	Second variation	63
5.4	Interlude: The Legendre transform	64
5.5	Lagrangian mechanics	66
5.6	Hamiltonian mechanics	67
5.7	Kinetic and potential energy	68
5.8	Problems	69
5.9	The path integral	70
5.10	Appendix: Lagrange multipliers	71
6	Perturbation theory	73
6.1	The implicit function theorem: scalar case	73
6.2	Problems	74
6.3	The implicit function theorem: systems	76

CONTENTS

5

6.4	Nonlinear differential equations	77
6.5	A singular perturbation example	78
6.6	Eigenvalues and eigenvectors	79
6.7	The self-adjoint case	81
6.8	The anharmonic oscillator	82

Chapter 1

Distributions

1.1 Properties of distributions

Consider the space $C_c^\infty(\mathbb{R})$ of complex test functions. These are complex functions defined on the real line, infinitely differentiable and with compact support. A distribution is a linear functional from this space to the complex numbers. It must satisfy a certain continuity condition, but we shall ignore that. The value of the distribution F on the test function ϕ is written $\langle F, \phi \rangle$.

If f is a locally integrable function, then we may define a distribution

$$\langle F, \phi \rangle = \int_{-\infty}^{\infty} f(x)\phi(x) dx. \quad (1.1)$$

Thus many functions define distributions. This is why distributions are also called generalized functions.

A sequence of distributions F_n is said to converge to a distribution F if for each test function ϕ the numbers $\langle F_n, \phi \rangle$ converge to the number $\langle F, \phi \rangle$.

Example. The distribution δ_a is defined by

$$\langle \delta_a, \phi \rangle = \phi(a). \quad (1.2)$$

This is not given by a locally integrable function. However a distribution may be written as a limit of functions. For instance, let $\epsilon > 0$ and consider the function

$$\delta_\epsilon(x - a) = \frac{1}{\pi} \frac{\epsilon}{(x - a)^2 + \epsilon^2}. \quad (1.3)$$

The limit of the distributions defined by these locally integrable functions as $\epsilon \downarrow 0$ is δ_a . For this reason the distribution is often written in the incorrect but suggestive notation

$$\langle \delta_a, \phi \rangle = \int_{-\infty}^{\infty} \delta(x - a)\phi(x) dx. \quad (1.4)$$

Operations on distributions are defined by looking at the example of a distribution defined by a function and applying the integration by parts formula

in that case. Thus, for instance, the derivative is defined by

$$\langle F', \phi \rangle = -\langle F, \phi' \rangle. \quad (1.5)$$

Example: Consider the locally integrable Heaviside function given by $H(x) = 1$ for $x > 0$, $H(x) = 0$ for $x < 0$. Then $H' = \delta$. Here δ is given by

$$\langle \delta, \phi \rangle = \phi(0). \quad (1.6)$$

Example: Consider the locally integrable log function $f(x) = \log|x|$. Then $f'(x) = \text{PV } 1/x$. Here the principal value $\text{PV } 1/x$ is given by

$$\langle \text{PV } \frac{1}{x}, \phi \rangle = \lim_{\epsilon \downarrow 0} \int_{-\infty}^{\infty} \frac{x}{x^2 + \epsilon^2} \phi(x) dx. \quad (1.7)$$

This can be seen by writing $\log|x| = \lim_{\epsilon} \log \sqrt{x^2 + \epsilon^2}$.

A distribution can be approximated by more than one sequence of functions. For instance, for each $a > 0$ let $\log_a(|x|) = \log(|x|)$ for $|x| \geq a$ and $\log_a(|x|) = \log(a)$ for $|x| < a$. Then $\log_a(|x|)$ also approaches $\log(|x|)$ as $a \rightarrow 0$. So its derivative is another approximating sequence for $\text{PV } 1/x$. This says that

$$\langle \text{PV } \frac{1}{x}, \phi \rangle = \lim_{a \downarrow 0} \int_{|x| > a} \frac{1}{x} \phi(x) dx. \quad (1.8)$$

We can use this sequence to compute the derivative of $\text{PV } 1/x$. We get

$$\int_{-\infty}^{\infty} \left(\frac{d}{dx} \text{PV } \frac{1}{x} \right) \phi(x) dx = - \lim_{a \rightarrow 0} \int_{|x| > a} \frac{1}{x} \phi'(x) dx = \lim_{a \rightarrow 0} \left[\int_{|x| > a} -\frac{1}{x^2} \phi(x) dx + \frac{\phi(a) + \phi(-a)}{a} \right]. \quad (1.9)$$

But $[\phi(a) + \phi(-a) - 2\phi(0)]/a$ converges to zero, so this is

$$\int_{-\infty}^{\infty} \left(\frac{d}{dx} \text{PV } \frac{1}{x} \right) \phi(x) dx = \lim_{a \rightarrow 0} \int_{|x| > a} -\frac{1}{x^2} [\phi(x) - \phi(0)] dx. \quad (1.10)$$

Another operator is multiplication of a distribution by a smooth function g in C^∞ . This is defined in the obvious way by

$$\langle g \cdot F, \phi \rangle = \langle F, g\phi \rangle. \quad (1.11)$$

Distributions are not functions. They may not have values at points, and in general nonlinear operations are not defined. For instance, the square of a distribution is not always a well defined distribution.

Also, some algebraic operations involving distributions are quite tricky. Consider, for instance, the associative law. Apply this to the three distributions $\delta(x)$, x , and $\text{PV } 1/x$. Clearly the product $\delta(x) \cdot x = 0$. On the other hand, the product $x \cdot \text{PV } 1/x = 1$ is one. So if the associate law were to hold, we would get

$$0 = 0 \cdot \text{PV } \frac{1}{x} = (\delta(x) \cdot x) \cdot \text{PV } \frac{1}{x} = \delta(x) \cdot (x \cdot \text{PV } \frac{1}{x}) = \delta(x) \cdot 1 = \delta(x). \quad (1.12)$$

1.2 Mapping distributions

A test function ϕ is naturally viewed as a covariant object, so the distribution F is contravariant. A proper function is a function such that the inverse image of each compact set is compact. It is natural to define the forward push of a distribution F by a smooth proper function g by $\langle g[F], \phi \rangle = \langle F, \phi \circ g \rangle$. Example: If F is the distribution $\delta(x-3)$ and if $u = g(x) = x^2 - 4$, then the forward push is $\delta(u-5)$. This is because $\int \delta(u-5)\phi(u) du = \int \delta(x-3)\phi(x^2-4) dx$.

On the other hand, it is actually more common to think of a distribution as being a covariant object, since a distribution is supposed to be a generalized function. The backward pull of the distribution by a smooth function g is defined in at least some circumstances by

$$\langle F \circ g, \phi \rangle = \langle F, g[\phi] \rangle. \quad (1.13)$$

Here

$$g[\phi](u) = \sum_{g(x)=u} \frac{\phi(x)}{|g'(x)|}. \quad (1.14)$$

Example. Let $u = g(x) = x^2 - 4$, with $a > 0$. Then the backward pull of $\delta(u)$ under g is $\delta(x^2 - 4) = (1/4)(\delta(x-2) + \delta(x+2))$. This is because

$$g[\phi](u) = \frac{\phi(\sqrt{u^2+4}) + \phi(-\sqrt{u^2+4})}{2\sqrt{u^2+4}}. \quad (1.15)$$

So if $F = \delta$, then

$$F \circ g = \frac{1}{4}(\delta_2 + \delta_{-2}) \quad (1.16)$$

Example: The backward pull is not always defined. To consider a distribution as a covariant object is a somewhat awkward act in general. Let $u = h(x) = x^2$. The backward pull of $\delta(u)$ by h is $\delta(x^2)$, which is not defined.

Example: The general formula for the pull back of the delta function is

$$\delta(g(x)) = \sum_{g(a)=0} \frac{1}{|g'(a)|} \delta(x-a). \quad (1.17)$$

The most important distributions are δ , PV $1/x$, $1/(x-i0)$, and $1/(x+i0)$. These are the limits of the functions $\delta_\epsilon(x)$, $x/(x^2 + \epsilon^2)$, $1/(x-i\epsilon)$, $1/(x+i\epsilon)$ as $\epsilon \downarrow 0$. The relations between these functions are given by

$$\delta_\epsilon(x) = \frac{1}{\pi} \frac{\epsilon}{x^2 + \epsilon^2} = \frac{1}{2\pi i} \left(\frac{1}{x-i\epsilon} - \frac{1}{x+i\epsilon} \right). \quad (1.18)$$

and

$$\frac{x}{x^2 + \epsilon^2} = \frac{1}{2} \left(\frac{1}{x-i\epsilon} + \frac{1}{x+i\epsilon} \right). \quad (1.19)$$

1.3 Radon measures

A Radon measure is a positive distribution. That is, it is a linear functional μ on $C_c^\infty(\mathbb{R})$ such that for each test function ϕ the condition $\phi \geq 0$ implies that the value $\langle \mu, \phi \rangle \geq 0$. Every Radon measure extends uniquely by continuity to a linear function on $C_c(\mathbb{R})$, the space of continuous functions with compact support. Each positive locally integrable function h defines a Radon measure by integrating $\phi(x)$ times $h(x) dx$. Also, the point mass distributions δ_a are Radon measures.

It is common to write the value of a Radon measure in the form

$$\langle \mu, \phi \rangle = \int \phi(x) d\mu(x). \quad (1.20)$$

What is remarkable is that the theory of Lebesgue integration works for Radon measures. That is, given a real function $f \geq 0$ that is only required to be Borel measurable, there is a natural definition of the integral such that

$$0 \leq \int f(x) d\mu(x) \leq +\infty. \quad (1.21)$$

Furthermore, if f is a complex Borel measurable function such that $|f|$ has finite integral, then the integral of f is defined and satisfies.

$$\left| \int f(x) d\mu(x) \right| \leq \int |f(x)| d\mu(x) < +\infty. \quad (1.22)$$

1.4 Approximate delta functions

It might seem that one could replace the notion of distribution by the notion of a sequence of approximating functions. This is true in some sense, but the fact is that many different sequences may approximate the same distribution. Here is a result of that nature.

Theorem. Let $\delta_1(u) \geq 0$ be a positive function with integral 1. For each $\epsilon > 0$ define $\delta_\epsilon(x) = \delta(x/\epsilon)/\epsilon$. Then the functions δ_ϵ converge to the δ distribution as ϵ tends to zero.

The convergence takes place in the sense of distributions (smooth test functions with compact support) or even in the sense of Radon measures (continuous test functions with compact support). Notice that there are no continuity or symmetry assumptions on the initial function.

The proof of this theorem is simple. Each δ_ϵ has integral one. Consider a bounded continuous function ϕ . Then

$$\int_{-\infty}^{\infty} \delta_\epsilon(x) \phi(x) dx = \int_{-\infty}^{\infty} \delta_1(u) \phi(\epsilon u) du. \quad (1.23)$$

The dominated convergence theorem shows that this approaches the integral

$$\int_{-\infty}^{\infty} \delta_1(u) \phi(0) du = \phi(0). \quad (1.24)$$

Here is an even stronger result.

Theorem. For each $\epsilon > 0$ let $\delta_\epsilon \geq 0$ be a positive function with integral 1. Suppose that for each $a > 0$ that

$$\int_{|x|>a} \delta_\epsilon(x) dx \rightarrow 0 \quad (1.25)$$

as $\epsilon \rightarrow 0$. Then the functions δ_ϵ converge to the δ distribution as ϵ tends to zero.

Here is a proof of this more general theorem. Let $H_\epsilon(a) = \int_0^a \delta_\epsilon(x) dx$. Then for each $a < 0$ we have $H_\epsilon(a) \rightarrow 0$, and for each $a > 0$ we have $1 - H_\epsilon(a) \rightarrow 0$. In other words, for each $a \neq 0$ we have $H_\epsilon(a) \rightarrow H(a)$ as $\epsilon \rightarrow 0$. Since the functions H_ϵ are uniformly bounded, it follows from the dominated convergence theorem that $H_\epsilon \rightarrow H$ in the sense of distributions. It follows by differentiation that $\delta_\epsilon \rightarrow \delta$ in the sense of distributions.

1.5 Problems

If F and G are distributions, and if at least one of them has compact support, then their convolution $F * G$ is defined by

$$\langle F * G, \phi \rangle = \langle F_x G_y, \phi(x + y) \rangle.$$

This product is commutative. It is also associative if at least two of the three factors have compact support.

1. If F and G are given by locally integrable functions f and g , and at least one has compact support, then $F * G$ is given by a locally integrable function

$$(f * g)(z) = \int_{-\infty}^{\infty} f(x)g(z - x) dx = \int_{-\infty}^{\infty} f(z - y)g(y) dy.$$

2. If G is given by a test function g , then $F * g$ is given by a smooth function

$$(F * g)(z) = \langle F_x, g(z - x) \rangle.$$

3. Calculate the convolution $1 * \delta'$.
4. Calculate the convolution $\delta' * H$, where H is the Heaviside function.
5. Calculate the convolution $(1 * \delta') * H$ and also calculate the convolution $1 * (\delta' * H)$. What does this say about the associative law for convolution?
6. Let L be a constant coefficient linear differential operator. Let u be a distribution that is a fundamental solution, that is, let $Lu = \delta$. Let G be a distribution with compact support. Show that the convolution $F = u * G$ satisfies the equation $LF = G$. Hint: Write $\langle LF, \phi \rangle = \langle F, L^\dagger \phi \rangle$, where L^\dagger is adjoint to L .

7. Take $L = -d^2/dx^2$. Is there a fundamental solution that has support in a bounded interval? Is there a fundamental solution that has support in a semi-infinite interval?

1.6 Tempered distributions

Let d/dx be the operator of differentiation, and let x be the operator of multiplication by the coordinate x . The space \mathcal{S} of rapidly decreasing smooth test functions consists of the functions ϕ in L^2 such that every finite product of the operators d/dx and x applied to ϕ is also in L^2 . The advantage of this definition is that the space \mathcal{S} is clearly invariant under Fourier transformation.

A tempered distribution is a linear functional on \mathcal{S} satisfying the appropriate continuity property. Every tempered distribution restricts to define a distribution. So tempered distributions are more special.

The advantage of tempered distributions is that one can define Fourier transforms \hat{F} of tempered distributions F . The definition is

$$\langle \hat{F}, \phi \rangle = \langle F, \hat{\phi} \rangle. \quad (1.26)$$

Here $\hat{\phi}$ is the Fourier transform of ϕ .

Here are some Fourier transforms for functions. The first two are easy. They are

$$\int_{-\infty}^{\infty} e^{-ikx} \frac{1}{x - i\epsilon} dx = 2\pi i e^{\epsilon k} H(-k) \quad (1.27)$$

and

$$\int_{-\infty}^{\infty} e^{-ikx} \frac{1}{x + i\epsilon} dx = -2\pi i e^{-\epsilon k} H(k). \quad (1.28)$$

Subtract these and divide by $2\pi i$. This gives

$$\int_{-\infty}^{\infty} e^{-ikx} \delta_{\epsilon}(x) dx = e^{-\epsilon|k|}. \quad (1.29)$$

Instead, add these and divide by 2. This gives

$$\int_{-\infty}^{\infty} e^{-ikx} \frac{x}{x^2 + \epsilon^2} dx = -\pi i e^{-\epsilon|k|} \text{sign}(k). \quad (1.30)$$

The corresponding Fourier transforms for distributions are

$$F[1/(x - i0)] = 2\pi i H(-k) \quad (1.31)$$

and

$$F[1/(x + i0)] = -2\pi i H(k). \quad (1.32)$$

Also,

$$F[\delta(x)] = 1 \quad (1.33)$$

and

$$F[\text{PV } \frac{1}{x}] = -\pi i \text{sign}(k). \quad (1.34)$$

Example: Here is a more complicated calculation. The derivative of PV $1/x$ is the distribution

$$\frac{d}{dx} \text{PV } \frac{1}{x} = -\frac{1}{x^2} + c\delta(x), \quad (1.35)$$

where c is the infinite constant

$$c = \int_{-\infty}^{\infty} \frac{1}{x^2} dx. \quad (1.36)$$

This makes rigorous sense if interprets it as

$$\int_{-\infty}^{\infty} \left(\frac{d}{dx} \text{PV } \frac{1}{x} \right) \phi(x) dx = \lim_{a \rightarrow 0} \int_{|x| > a} -\frac{1}{x^2} [\phi(x) - \phi(0)] dx. \quad (1.37)$$

One can get an intuitive picture of this result by graphing the approximating functions. The key formula is

$$\frac{d}{dx} \frac{x}{x^2 + \epsilon^2} = -\frac{x^2}{(x^2 + \epsilon^2)^2} + c_\epsilon \frac{2\epsilon^3}{\pi(x^2 + \epsilon^2)^2}, \quad (1.38)$$

where $c_\epsilon = \pi/(2\epsilon)$. This is an approximation to $-1/x^2$ plus a big constant times an approximation to the delta function.

The Fourier transform of the derivative is obtained by multiplying the Fourier transform of PV $1/x$ by ik . Thus the Fourier transform of $-1/x^2 + c\delta(x)$ is ik times $-\pi i \text{sign}(k)$ which is $\pi|k|$.

This example is interesting, because it looks at first glance that the derivative of PV $1/x$ should be $-1/x^2$, which is negative definite. But the correct answer for the derivative is $-1/x^2 + c\delta(x)$, which is actually positive definite. And in fact its Fourier transform is positive.

For each of these formula there is a corresponding inverse Fourier transform. For, instance, the inverse Fourier transform of 1 is

$$\delta(x) = \int_{-\infty}^{\infty} e^{ikx} \frac{dk}{2\pi} = \int_0^{\infty} \cos(kx) \frac{dk}{2\pi}. \quad (1.39)$$

Of course such an equation is interpreted by integrating both sides with a test function.

Another formula of the same type is gotten by taking the inverse Fourier transform of $-\pi i \text{sign}(k)$. This is

$$\text{PV } \frac{1}{x} = -\pi i \int_{-\infty}^{\infty} e^{ikx} \text{sign}(k) \frac{dk}{2\pi} = \int_0^{\infty} \sin(kx) dk. \quad (1.40)$$

1.7 Poisson equation

We begin the study of fundamental solutions of partial differential equations. These are solutions of the equation $Lu = \delta$, where L is the differential operator, and δ is a point source.

Let us start with the equation in one space dimension:

$$\left(-\frac{d^2}{dx^2} + m^2\right)u = \delta(x). \quad (1.41)$$

This is an equilibrium equation that balances a source with losses due to diffusion and to dissipation (when $m > 0$). Fourier transform. This gives

$$(k^2 + m^2)\hat{u}(k) = 1. \quad (1.42)$$

The solution is

$$\hat{u}(k) = \frac{1}{k^2 + m^2}. \quad (1.43)$$

There is no problem of division by zero. The inverse Fourier transform is

$$u(x) = \frac{1}{2m}e^{-m|x|}. \quad (1.44)$$

This is the only solution that is a tempered distribution. (The solutions of the homogeneous equation all grow exponentially.)

What happens when $m = 0$? This is more subtle. The equation is

$$-\frac{d^2}{dx^2}u = \delta(x). \quad (1.45)$$

Fourier transform. This gives

$$k^2\hat{u}(k) = 1. \quad (1.46)$$

Now there is a real question about division by zero. Furthermore, the homogeneous equation has solutions that are tempered distributions, namely linear combinations of $\delta(k)$ and $\delta'(k)$. The final result is that the inhomogeneous equation does have a tempered distribution solution, but it needs careful definition. The solution is

$$\hat{u}(k) = \frac{1}{k^2} + \infty\delta(k). \quad (1.47)$$

This may be thought of as the derivative of $-\text{PV } 1/k$. The inverse Fourier transform of $\text{PV } 1/k$ is $(1/2)i\text{sign}(x)$. So the inverse Fourier transform of $-d/dk\text{PV } 1/k$ is $-(-ix)(1/2)i\text{sign}(x) = -(1/2)|x|$. Thus

$$u(x) = -\frac{1}{2}|x| \quad (1.48)$$

is a solution of the inhomogeneous equation. The solutions of the homogeneous equation are linear combinations of 1 and x . None of these solutions are a

good description of diffusive equilibrium. In fact, in one dimension there is no diffusive equilibrium.

The next case that is simple to compute and of practical importance is the equation in dimension 3. This is

$$(-\nabla^2 + m^2)u = \delta(x). \quad (1.49)$$

This is an equilibrium equation that balances a source with losses due to diffusion and to dissipation (when $m > 0$). Fourier transform. This gives

$$(k^2 + m^2)\hat{u}(k) = 1. \quad (1.50)$$

The solution is

$$\hat{u}(k) = \frac{1}{k^2 + m^2}. \quad (1.51)$$

The inverse Fourier transform in the three dimension case may be computed by going to polar coordinates. It is

$$u(x) = \frac{1}{4\pi|x|}e^{-m|x|}. \quad (1.52)$$

What happens when $m = 0$? The situation is very different in three dimensions. The equation is

$$\nabla^2 u = \delta(x). \quad (1.53)$$

Fourier transform. This gives

$$k^2\hat{u}(k) = 1. \quad (1.54)$$

The inhomogeneous equation has a solution

$$\hat{u}(k) = \frac{1}{k^2}. \quad (1.55)$$

But now this is a locally integrable function. It defines a tempered distribution without any regularization. Thus

$$u(x) = \frac{1}{4\pi|x|} \quad (1.56)$$

is a solution of the inhomogeneous equation. In three dimensions there is diffusive equilibrium. There is so much room that the effect of the source can be completely compensated by diffusion alone.

1.8 Diffusion equation

The diffusion equation or heat equation is

$$\left(\frac{\partial}{\partial t} - \frac{1}{2}\sigma^2\frac{\partial^2}{\partial x^2}\right)u = \delta(x)\delta(t). \quad (1.57)$$

It says that the time rate of change is entirely due to diffusion. Fourier transform. We get

$$(i\omega + \frac{1}{2}\sigma^2 k^2)\hat{u}(k, \omega) = 1. \quad (1.58)$$

This has solution

$$\hat{u}(k, \omega) = \frac{1}{i\omega + \frac{1}{2}\sigma^2 k^2} = \frac{1}{i} \frac{1}{\omega - i\frac{1}{2}\sigma^2 k^2}. \quad (1.59)$$

Here the only division by zero is when both ω and k are zero. But this is not so serious, because it is clear how to regularize. We can use the fact that the inverse Fourier transform of $1/(\omega - i\epsilon)$ with $\epsilon > 0$ is $iH(t)e^{-\epsilon t}$. So we have

$$\hat{u}(k, t) = H(t)e^{-\frac{\sigma^2 t k^2}{2}}. \quad (1.60)$$

This is a Gaussian, so the fundamental solution is

$$u(x, t) = H(t) \frac{1}{\sqrt{2\pi\sigma^2 t}} e^{-\frac{x^2}{2\sigma^2 t}}. \quad (1.61)$$

1.9 Wave equation

We will look for the solution of the wave equation with a point source at time zero that lies in the forward light cone. The wave equation in 1 + 1 dimensions is

$$\left(\frac{\partial^2}{\partial t^2} - c^2 \frac{\partial^2}{\partial x^2} \right) u = \delta(x)\delta(t). \quad (1.62)$$

Fourier transform. We get

$$(-\omega^2 + c^2 k^2)\hat{u}(k, \omega) = 1. \quad (1.63)$$

This has solution

$$\hat{u}(k, \omega) = -\frac{1}{(\omega - i0)^2 - c^2 k^2}. \quad (1.64)$$

The division by zero is serious, but it is possible to regularize. The choice of regularization is not the only possible one, but we shall see that it is the convention that gives propagation into the future. We can write this also as

$$\hat{u}(k, \omega) = -\frac{1}{2c|k|} \left(\frac{1}{\omega - i0 - c|k|} - \frac{1}{\omega - i0 + c|k|} \right). \quad (1.65)$$

We can use the fact that the inverse Fourier transform of $1/(\omega - i0)$ is $iH(t)$. So we have

$$\hat{u}(k, t) = -\frac{1}{2c|k|} iH(t) \left[e^{ic|k|t} - e^{-ic|k|t} \right] = \frac{1}{c|k|} \sin(c|k|t)H(t). \quad (1.66)$$

It is easy to check that this is the Fourier transform of

$$u(x, t) = \frac{1}{2c} [H(x + ct) - H(x - ct)] H(t). \quad (1.67)$$

The wave equation in 3 + 1 dimensions is

$$\left(\frac{\partial^2}{\partial t^2} - c^2 \nabla^2 \right) u = \delta(x) \delta(t). \quad (1.68)$$

Fourier transform. We get

$$(-\omega^2 + c^2 k^2) \hat{u}(k, \omega) = 1. \quad (1.69)$$

This has solution

$$\hat{u}(k, \omega) = -\frac{1}{(\omega - i0)^2 - c^2 k^2}. \quad (1.70)$$

Again we can write this as

$$\hat{u}(k, \omega) = -\frac{1}{2c|k|} \left(\frac{1}{\omega - i0 - c|k|} - \frac{1}{\omega - i0 + c|k|} \right). \quad (1.71)$$

Thus we have again

$$\hat{u}(k, t) = -\frac{1}{2c|k|} iH(t) [e^{ic|k|t} - e^{-ic|k|t}] = \frac{1}{c|k|} \sin(c|k|t) H(t). \quad (1.72)$$

However the difference is that the k variable is three dimensional. It is easy to check that this is the Fourier transform of

$$u(x, t) = \frac{1}{4\pi c|x|} \delta(|x| - ct) H(t). \quad (1.73)$$

This is a beautiful formula. It represents an expanding sphere of influence, going into the future. Inside the sphere it is dark. The solution has an even more beautiful expression that exhibits the symmetry:

$$u(x, t) = \frac{1}{2\pi c} \delta(x^2 - c^2 t^2) H(t). \quad (1.74)$$

1.10 Homogeneous solutions of the wave equation

The polynomial $\omega^2 - k^2$ vanishes on an entire cone, so it is not surprising that the wave equation has a number of interesting homogeneous solutions. The most important ones are

$$\hat{u}(k, t) = \frac{1}{c|k|} \sin(c|k|t) \quad (1.75)$$

and its derivative

$$\hat{v}(k, t) = \cos(c|k|t). \quad (1.76)$$

These are the solutions that are used in constructing solutions of the initial value problem.

It is interesting to see what these solutions look like in the frequency representation. The result is

$$\hat{u}(k, \omega) = -\pi i \frac{1}{c|k|} [\delta(\omega - c|k|) - \delta(\omega + c|k|)] = -2\pi i \delta(\omega^2 - c^2 k^2) \text{sign}(\omega) \quad (1.77)$$

and

$$\hat{v}(k, \omega) = i\omega \hat{u}(k, \omega) = \pi [\delta(\omega - c|k|) + \delta(\omega + c|k|)] = 2\pi |\omega| \delta(\omega^2 - c^2 k^2). \quad (1.78)$$

1.11 Problems

1. Show that $\frac{1}{x^{\frac{1}{3}}}$ is a locally integrable function and thus defines a distribution. Show that its distribution derivative is

$$\frac{d}{dx} \frac{1}{x^{\frac{1}{3}}} = -\frac{1}{3} \frac{1}{x^{\frac{4}{3}}} + c\delta(x), \quad (1.79)$$

where

$$c = \frac{1}{3} \int_{-\infty}^{\infty} \frac{1}{x^{\frac{4}{3}}} dx \quad (1.80)$$

Hint: To make this rigorous, consider $x/(x^2 + \epsilon^2)^{\frac{2}{3}}$.

2. Show that $\frac{1}{|x|^{\frac{1}{3}}}$ is a locally integrable function and thus defines a distribution. Show that its distribution derivative is

$$\frac{d}{dx} \frac{1}{|x|^{\frac{1}{3}}} = -\frac{1}{3} \frac{1}{|x|^{\frac{4}{3}}} \text{sign}(x). \quad (1.81)$$

The right hand side is not locally integrable. Explain the definition of the right hand side as a distribution. Hint: To make this rigorous, consider $1/(x^2 + \epsilon^2)^{\frac{1}{6}}$.

3. Discuss the contrast between the results in the last two problems. It may help to draw some graphs of the functions that approximate these distributions.
4. Let $m > 0$. Use Fourier transforms to find a tempered distribution u that is the fundamental solution of the equation

$$-\frac{d^2}{dx^2} u + m^2 u = \delta(x). \quad (1.82)$$

Is this a unique tempered distribution solution? Explain.

5. Let $m > 0$. Consider Euclidian space of dimension $n = 3$. Use Fourier transforms to find a tempered distribution u that is the fundamental solution of the equation

$$-\nabla^2 u + m^2 u = \delta(x). \quad (1.83)$$

1.12 Answers to first two problems

1. The function $\frac{1}{x^{\frac{1}{3}}}$ is locally integrable, while its pointwise derivative $-\frac{1}{3} \frac{1}{x^{\frac{4}{3}}}$ is not. But we do have

$$\frac{d}{dx} \frac{x}{(x^2 + \epsilon^2)^{\frac{2}{3}}} = -\frac{1}{3} \frac{x^2}{(x^2 + \epsilon^2)^{\frac{5}{3}}} + \frac{\epsilon^2}{(x^2 + \epsilon^2)^{\frac{5}{3}}}. \quad (1.84)$$

Let

$$c_\epsilon = \int_{-\infty}^{\infty} \frac{1}{3} \frac{x^2}{(x^2 + \epsilon^2)^{\frac{5}{3}}} dx \quad (1.85)$$

which is easily seen to be proportional to $1/\epsilon^{\frac{1}{3}}$. From the fundamental theory of calculus

$$\int_{-\infty}^{\infty} \frac{\epsilon^2}{(x^2 + \epsilon^2)^{\frac{5}{3}}} dx = c_\epsilon. \quad (1.86)$$

Write

$$\delta_\epsilon(x) = \frac{1}{c_\epsilon} \frac{\epsilon^2}{(x^2 + \epsilon^2)^{\frac{5}{3}}}. \quad (1.87)$$

Then $\delta_\epsilon(x) \rightarrow \delta(x)$ in the sense of distributions. Furthermore,

$$\int_{-\infty}^{\infty} \frac{d}{dx} \frac{1}{x^{\frac{1}{3}}} \phi(x) dx = \lim_{\epsilon \rightarrow 0} \int_{-\infty}^{\infty} \left[-\frac{1}{3} \frac{x^2}{(x^2 + \epsilon^2)^{\frac{5}{3}}} + c_\epsilon \delta_\epsilon(x) \right] \phi(x) dx. \quad (1.88)$$

Since for ϕ smooth enough

$$\int c_\epsilon \delta_\epsilon(x) [\phi(x) - \phi(0)] dx \rightarrow 0, \quad (1.89)$$

this may be written in the simple form

$$\int_{-\infty}^{\infty} \frac{d}{dx} \frac{1}{x^{\frac{1}{3}}} \phi(x) dx = \int_{-\infty}^{\infty} -\frac{1}{3} \frac{1}{x^{\frac{4}{3}}} [\phi(x) - \phi(0)] dx. \quad (1.90)$$

Notice that the integrand is integrable for each test function ϕ .

2. The function $\frac{1}{|x|^{\frac{1}{3}}}$ is locally integrable, while its pointwise derivative $-\frac{1}{3} \frac{1}{x^{\frac{4}{3}}} \text{sign}(x)$ is not. We have

$$\frac{d}{dx} \frac{1}{(x^2 + \epsilon^2)^{\frac{1}{6}}} = -\frac{1}{3} \frac{x}{(x^2 + \epsilon^2)^{\frac{7}{6}}}. \quad (1.91)$$

Since this derivative is an odd function, we have

$$\int_{-\infty}^{\infty} \frac{d}{dx} \frac{1}{|x|^{\frac{1}{3}}} \phi(x) dx = \lim_{\epsilon \rightarrow 0} \int_{-\infty}^{\infty} -\frac{1}{3} \frac{x}{(x^2 + \epsilon^2)^{\frac{7}{6}}} [\phi(x) - \phi(0)] dx. \quad (1.92)$$

We can write this as

$$\int_{-\infty}^{\infty} \frac{d}{dx} \frac{1}{|x|^{\frac{1}{3}}} \phi(x) dx = \int_{-\infty}^{\infty} -\frac{1}{3} \frac{1}{x^{\frac{4}{3}}} \text{sign}(x) [\phi(x) - \phi(0)] dx. \quad (1.93)$$

Again the integrand is integrable.

Chapter 2

Bounded Operators

2.1 Introduction

This chapter deals with bounded linear operators. These are bounded linear transformations of a Hilbert space into itself. In fact, the chapter treats four classes of operators: finite rank, Hilbert-Schmidt, compact, and bounded. Every finite rank operator is Hilbert-Schmidt. Every Hilbert-Schmidt operator is compact. Every compact operator is bounded.

We shall see in the next chapter that it is also valuable to look at an even broader class of operators, those that are closed and densely defined. Every bounded operator (everywhere defined) is closed and densely defined. However the present chapter treats only bounded operators.

2.2 Bounded linear operators

Let H be a Hilbert space (a vector space with an inner product that is a complete metric space). A linear transformation $K : H \rightarrow H$ is said to be bounded if it maps bounded sets into bounded sets. This is equivalent to there being a constant M with

$$\|Ku\| \leq M\|u\|. \tag{2.1}$$

Let M_2 be the least such M . This is called uniform norm of K and is written $\|K\|_\infty$ or simply $\|K\|$ when the context makes this clear. [The subscript in M_2 is supposed to remind us that we are dealing with Hilbert spaces like L^2 . The subscript in $\|K\|_\infty$, on the other hand, tells that we are looking at a least upper bound.]

If K and L are bounded operators, their sum $K + L$ and product KL are bounded. Furthermore, we have $\|K + L\|_\infty \leq \|K\|_\infty + \|L\|_\infty$ and $\|KL\|_\infty \leq \|K\|_\infty\|L\|_\infty$.

A bounded operator K always has an adjoint K^* that is a bounded operator.

It is the unique operator with the property that

$$(u, K^*v) = (Ku, v) \quad (2.2)$$

for all u, v in H . Furthermore $K^{**} = K$, and $\|K^*\|_\infty = \|K\|_\infty$. For products we have $(KL)^* = L^*K^*$, in the opposite order.

It is also possible to define the adjoint of an operator from one Hilbert space to another. Here is a special case. Let g be a vector in the Hilbert space H . Then g defines a linear transformation from \mathbf{C} to H by sending z to the vector zg . The adjoint is a linear transformation from H to \mathbf{C} , denoted by g^* . It is the transformation that sends v to (g, v) , so $g^*v = (g, v)$. The adjointness relation is $\bar{z}g^*v = (zg, v)$ which is just $\bar{z}(g, v) = (zg, v)$.

Let f be another vector. Define the operator K from H to H by $Ku = f(g, u)$, that is, $K = fg^*$. Then the adjoint of K is $K^* = gf^*$.

Example: Hilbert-Schmidt integral operators. Let

$$(Kf)(x) = \int k(x, y)f(y) dy. \quad (2.3)$$

with k in L^2 , that is,

$$\|k\|_2^2 = \iint |k(x, y)|^2 dx dy < \infty. \quad (2.4)$$

Then K is bounded with norm $\|K\|_\infty \leq \|k\|_2$.

Proof: Fix x . Apply the Schwarz inequality to the integral over y in the definition of the operator. This gives that $|(Kf)(x)|^2 \leq \int |k(x, y)|^2 dy \int |f(y)|^2 dy$. Now integrate over x .

Example: Interpolation integral operators. Let K be an integral operator such that

$$M_1 = \sup_y \int |k(x, y)| dx < \infty \quad (2.5)$$

and

$$M_\infty = \sup_x \int |k(x, y)| dy < \infty. \quad (2.6)$$

(Here we choose to think of $k(x, y)$ as a function of x and y , not as a Schwartz distribution like $\delta(x - y)$.) Let $1/p + 1/q = 1$. Then for each p with $1 \leq p \leq \infty$ the norm M_p of K as an operator on L^p is bounded by $M_p \leq M_1^{\frac{1}{p}} M_\infty^{\frac{1}{q}}$. In particular as an operator on L^2 the norm $M_2 = \|K\|_\infty$ of K is bounded by

$$M_2 \leq \sqrt{M_1 M_\infty}. \quad (2.7)$$

The reason for the name interpolation operator (which is not standard) is that the bound interpolates for all p from the extreme cases $p = 1$ (where the bound is M_1) and $p = \infty$ (where the bound is M_∞).

Proof: Write

$$|(Kf)(x)| \leq \int |k(x, y)||f(y)| dy = \int |k(x, y)|^{\frac{1}{q}} \cdot |k(x, y)|^{\frac{1}{p}} |f(y)| dy. \quad (2.8)$$

Apply the Hölder inequality to the integral. This gives

$$|(Kf)(x)| \leq \left(\int |k(x, y)| dy \right)^{\frac{1}{q}} \left(\int |k(x, y)||f(y)|^p dy \right)^{\frac{1}{p}} \leq M_{\infty}^{\frac{1}{q}} \left(\int |k(x, y)||f(y)|^p dy \right)^{\frac{1}{p}}. \quad (2.9)$$

It follows that

$$\int |(Kf)(x)|^p dx \leq M_{\infty}^{\frac{p}{q}} \int \int |k(x, y)||f(y)|^p dy dx \leq M_{\infty}^{\frac{p}{q}} M_1 \int |f(y)|^p dy. \quad (2.10)$$

Take the p th root to get the bound

$$\left(\int |(Kf)(x)|^p dx \right)^{\frac{1}{p}} \leq M_{\infty}^{\frac{1}{q}} M_1^{\frac{1}{p}} \left(\int |f(y)|^p dy \right)^{\frac{1}{p}}. \quad (2.11)$$

For a bounded operator K the set of points μ such that $(\mu I - K)^{-1}$ is a bounded operator is called the resolvent set of K . The complement of the resolvent set is called the spectrum of K . When one is only interested in values of $\mu \neq 0$ it is common to define $\lambda = 1/\mu$ and write

$$\mu(\mu I - K)^{-1} = (I - \lambda K)^{-1} \quad (2.12)$$

While one must be alert to which convention is being employed, this should be recognized as a trivial relabeling.

Theorem. If complex number μ satisfies $\|K\|_{\infty} < |\mu|$, then μ is in the resolvent set of K .

Proof: This is the Neumann series expansion

$$(\mu I - K)^{-1} = \frac{1}{\mu} \sum_{j=0}^{\infty} \frac{1}{\mu^j} K^j. \quad (2.13)$$

The j th term has norm $\|K^j\|_{\infty}/|\mu|^j \leq (\|K\|_{\infty}/|\mu|)^j$. This is the j th term of a convergent geometric series.

Theorem. If for some $n \geq 1$ the complex number μ satisfies $\|K^n\|_{\infty} < |\mu|^n$, then μ is in the resolvent set of K .

This is the Neumann series again. But now we only require the estimate for the n th power of the operator. Write $j = an + b$, where $0 \leq b < n$. Then $\|K^j\|_{\infty}/|\mu|^j$ is bounded by $(\|K\|_{\infty}/|\mu|)^b (\|K^n\|_{\infty}/|\mu|^n)^a$. So this is the sum of n convergent geometric series, one for each value of b between 0 and $n - 1$.

The spectral radius of an operator is the largest value of $|\mu|$, where μ is in the spectrum. The estimate of this theorem gives an upper bound on the spectral radius.

This is a remarkable result, since it has no analog for scalars. However for a matrix the norm of a power can be considerably smaller than the corresponding power of the norm, so this result can be quite useful. The most spectacular application is to Volterra integral operators.

Example: Volterra integral operators. Let $H = L^2(0, 1)$ and

$$(Kf)(x) = \int_0^x k(x, y)f(y) dy. \quad (2.14)$$

Suppose that $|k(x, y)| \leq C$ for $0 \leq y \leq x \leq 1$ and $k(x, y) = 0$ for $0 \leq x < y \leq 1$. Then

$$\|K^n\|_\infty \leq \frac{C^n}{(n-1)!}. \quad (2.15)$$

As a consequence every complex number $\mu \neq 0$ is in the resolvent set of K .

Proof: Each power K^n with $n \geq 1$ is also a Volterra integral operator. We claim that it has kernel $k^n(x, y)$ with

$$|k^n(x, y)| \leq \frac{C^n}{(n-1)!} (x-y)^{n-1}. \quad (2.16)$$

for $0 \leq y \leq x \leq 1$ and zero otherwise. This follows by induction.

Since $|k^n(x, y)| \leq C^n/(n-1)!$, the Hilbert-Schmidt norm of K is also bounded by $C^n/(n-1)!$. However this goes to zero faster than every power. So the Neumann series converges.

The norm of a bounded operator can always be found by calculating the norm of a self-adjoint bounded operator. In fact, $\|K\|_\infty^2 = \|K^*K\|_\infty$. Furthermore, the norm of a self-adjoint operator is its spectral radius. This shows that to calculate the norm of a bounded operator K exactly, one needs only to calculate the spectral radius of K^*K and take the square root. Unfortunately, this can be a difficult problem. This is why it is good to have other ways of estimating the norm.

2.3 Compact operators

Let H be a Hilbert space. A subset S is totally bounded if for every $\epsilon > 0$ there is a cover of S by a finite number N of ϵ balls. A totally bounded set is bounded.

Thus for instance, if H is finite dimensional with dimension n and S is a cube of side L , then $N \approx (L/\epsilon)^n$. This number N is finite, though it increases with ϵ . One expects a cube or a ball to be totally bounded only in finite dimensional situations.

However the situation is different for a rectangular shaped region in infinite dimensions. Say that the sides are $L_1, L_2, L_3, \dots, L_k, \dots$ and that these decrease to zero. Consider $\epsilon > 0$. Pick k so large that L_{k+1}, L_{k+1}, \dots are all less than ϵ . Then $N \approx (L_1/\epsilon)(L_2/\epsilon) \cdots (L_k/\epsilon)$. This can increase very rapidly with ϵ . But such a region that is fat only in finitely many dimensions and increasingly thin in all the others is totally bounded.

A linear transformation $K : H \rightarrow H$ is said to be compact if it maps bounded sets into totally bounded sets. (One can take the bounded set to be the unit ball.) A compact operator is bounded.

If K and L are compact operators, then so is their sum $K + L$. If K is compact and L is bounded, then KL and LK are compact.

If the self-adjoint operator K^*K is compact, then K is compact.

Proof: Let $\epsilon > 0$. Then there are u_1, \dots, u_k in the unit ball such that for each u in the unit ball there is a j with $\|K^*K(u - u_j)\| < \epsilon/2$. But this says that

$$\|K(u - u_j)\| = (K(u - u_j), K(u - u_j)) = ((u - u_j), K^*K(u - u_j)) \leq \|u - u_j\| \|K^*K(u - u_j)\| < \epsilon. \quad (2.17)$$

The adjoint K^* of a compact operator K is a compact operator.

Proof: Say that K is compact. Then since K^* is bounded, it follows that KK^* is compact. It follows from the last result that K^* is compact.

Approximation theorem. If K_n is a sequence of compact operators, and K is a bounded operator, and if $\|K_n - K\|_\infty \rightarrow 0$ and $n \rightarrow \infty$, then K is also compact.

The proof is a classical $\epsilon/3$ argument. Let $\epsilon > 0$. Choose n so large that $\|K - K_n\|_\infty < \frac{\epsilon}{3}$. Since K_n is compact, there are finitely many vectors u_1, \dots, u_k in the unit ball such that every vector $K_n u$ with u in the unit ball is within $\epsilon/3$ of some $K_n u_j$. Consider an arbitrary u in the unit ball and pick the corresponding u_j . Since

$$Ku - Ku_j = (K - K_n)u + (K_n u - K_n u_j) + (K_n - K)u_j, \quad (2.18)$$

it follows that

$$\|Ku - Ku_j\| \leq \|(K - K_n)u\| + \|K_n u - K_n u_j\| + \|(K_n - K)u_j\| \leq \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon. \quad (2.19)$$

This shows that the image of the unit ball under K is totally bounded.

Spectral properties of compact operators. Let K be a compact operator. The only non-zero points in the spectrum of K are eigenvalues of finite multiplicity. The only possible accumulation point of the spectrum is 0.

Notice that there is no general claim that the eigenvectors of K form a basis for the Hilbert space. The example of a Volterra integral operator provides a counterexample: The only point in the spectrum is zero.

Let K be a compact operator. We want to consider equations of the form

$$\mu u = f + Ku, \quad (2.20)$$

where μ is a parameter. If $\mu = 0$, then this is an equation of the first kind. If $\mu \neq 0$, then this is an equation of the second kind. Very often an equation of the second kind is written $u = f_1 + \lambda Ku$, where $\lambda = 1/\mu$, and where $f_1 = \lambda f$.

The condition for a unique solution of an equation of the first kind is the existence of the inverse K^{-1} , and the solution is $u = -K^{-1}f$. Thus the issue for an equation of the first kind is whether $\mu = 0$ is not an eigenvalue of K . If $\mu = 0$ is not an eigenvalue, the operator K^{-1} will be typically be an unbounded operator that is only defined on a linear subspace of the Hilbert space.

The condition for a unique solution of an equation of the second kind is the existence of the inverse $(\mu I - K)^{-1}$ for a particular value of $\mu \neq 0$, and the solution is $u = (\mu I - K)^{-1}f$. The issue for an equation of the second kind is whether $\mu \neq 0$ is not an eigenvalue of K . In this case, if $\mu \neq 0$ is not an eigenvalue, then $(\mu I - K)^{-1}$ will be a bounded operator. Thus equations of the second kind are much nicer. This is because compact operators have much better spectral properties away from zero.

Spectral theorem for compact self-adjoint operators. Let K be a compact self-adjoint operator. Then there is an orthonormal basis u_j of eigenvectors of K . The eigenvalues μ_j of K are real. Each non-zero eigenvalue is of finite multiplicity. The only possible accumulation point of the eigenvalues is zero. The operator K has the representation

$$Kf = \sum_j \mu_j u_j (u_j, f). \quad (2.21)$$

In abbreviated form this is

$$K = \sum_j \mu_j u_j u_j^*. \quad (2.22)$$

There is yet another way of writing the spectral theorem for compact operators. Define the unitary operator U from H to ℓ^2 by $(Uf)_j = (u_j, f)$. Its inverse is the unitary operator from ℓ^2 to H given by $(U^*c) = \sum c_j u_j$. Let M be the diagonal operator from ℓ^2 to ℓ^2 defined by multiplication by μ_j . Then

$$K = U^* M U. \quad (2.23)$$

The norm of a compact operator can always be found by calculating the norm of a self-adjoint compact operator. In fact, $\|K\|_\infty^2 = \|K^*K\|_\infty$. Furthermore, the norm of a compact self-adjoint operator is its spectral radius, which in this case is the largest value of $|\mu|$, where μ is an eigenvalue. Unfortunately, this can be a difficult computation.

Singular value decomposition for compact operators. Let K be a compact operator. Then there is an orthonormal family u_j and an orthonormal family w_j and a sequence of numbers $\chi_j \geq 0$ (singular values of K) approaching zero such that the operator K has the representation

$$Kf = \sum_j \chi_j w_j (u_j, f). \quad (2.24)$$

In abbreviated form this is

$$K = \sum_j \chi_j w_j u_j^*. \quad (2.25)$$

Sketch of proof: The operator K^*K is self-adjoint with positive eigenvalues χ_j^2 . We can write

$$K^*Kf = \sum_j \chi_j^2 u_j (u_j, f). \quad (2.26)$$

Then

$$\sqrt{K^*K}f = \sum_j \chi_j u_j(u_j, f). \quad (2.27)$$

Since $\|Kf\| = \|\sqrt{K^*K}f\|$ for each f , we can write $K = V\sqrt{K^*K}$, where $\|Vg\| = \|g\|$ for all $g = \sqrt{K^*K}f$ in the range of $\sqrt{K^*K}$. This is the well-known polar decomposition. Then

$$Kf = \sum_j \chi_j w_j(u_j, f), \quad (2.28)$$

where $w_j = Vu_j$.

There is another way of writing the singular value decomposition of K . Let $\sqrt{K^*K} = U^*DU$ be the spectral representation of $\sqrt{K^*K}$, where D is diagonal with entries $\chi_j \geq 0$. Then $K = VU^*DU = WDU$.

It follows from the approximation theorem and from the singular value decomposition that an operator is compact if and only if it is a norm limit of a sequence of finite rank operators.

Notice that this theorem gives a fairly clear picture of what a compact operator acting on L^2 looks like. It is an integral operator with kernel

$$k(x, y) = \sum_j \chi_j w_j(x) \overline{u_j(y)}, \quad (2.29)$$

where the $\chi_j \rightarrow 0$. Of course this representation may be difficult to find in practice. What happens in the case of a Hilbert-Schmidt integral operator is special: the χ_j go to zero sufficiently rapidly that $\sum_j |\chi_j|^2 < \infty$.

2.4 Hilbert-Schmidt operators

Let H be a Hilbert space. For a positive bounded self-adjoint operator B the trace is defined by $\text{tr}B = \sum_j (e_j, Be_j)$, where the e_j form an orthonormal basis. A bounded linear operator $K : H \rightarrow H$ is said to be Hilbert-Schmidt if $\text{tr}(K^*K) < \infty$. The Hilbert-Schmidt norm is $\|K\|_2 = \sqrt{\text{tr}(K^*K)}$.

Theorem. If K is a Hilbert-Schmidt integral operator with $(Kf)(x) = \int k(x, y)f(y) dy$ then

$$\|K\|_2^2 = \iint |k(x, y)|^2 dx dy. \quad (2.30)$$

Proof: Let $\{e_1, e_2, \dots\}$ be an orthonormal basis for H . We can always expand a vector in this basis via

$$u = \sum_j e_j(e_j, u). \quad (2.31)$$

(Remember the convention adopted here that inner products are linear in the second variable, conjugate linear in the first variable.) Write

$$Kf = \sum_i e_i \sum_j (e_i, Ke_j)(e_j, f). \quad (2.32)$$

The matrix elements (e_i, Ke_j) satisfy

$$\sum_i \sum_j |(e_i, Ke_j)|^2 = \sum_j \sum_i (Ke_j, e_i)(e_i, Ke_j) = \sum_j \|Ke_j\|^2 = \sum_j (e_j, K^*Ke_j) = \text{tr}(K^*K). \quad (2.33)$$

So the kernel of the integral operator K is

$$k(x, y) = \sum_i \sum_j (e_i, Ke_j) e_i(x) \overline{e_j(y)}. \quad (2.34)$$

This sum is convergent in $L^2(dx dy)$.

A Hilbert-Schmidt operator is a bounded operator. It is always true that $\|K\|_\infty \leq \|K\|_2$.

Theorem. A Hilbert-Schmidt operator is compact.

Proof: Let K be a Hilbert-Schmidt operator. Then K is given by a square-summable matrix. So K may be approximated by a sequence of finite-rank operators K_n such that $\|K_n - K\|_2 \rightarrow 0$. In particular, $\|K_n - K\|_\infty \rightarrow 0$. Since each K_n is compact, it follows from the approximation theorem that K is compact.

If K and L are Hilbert-Schmidt operators, their sum $K + L$ is a Hilbert-Schmidt operator and $\|K + L\|_2 \leq \|K\|_2 + \|L\|_2$. If K is a Hilbert-Schmidt operator and L is a bounded operator, then the products KL and LK are Hilbert-Schmidt. Furthermore, $\|KL\|_2 \leq \|K\|_2 \|L\|_\infty$ and $\|LK\|_2 \leq \|L\|_\infty \|K\|_2$.

The adjoint of a Hilbert-Schmidt operator is a Hilbert-Schmidt operator. Furthermore, $\|K^*\|_2 = \|K\|_2$.

Notice that the Hilbert-Schmidt norm of a bounded operator is defined in terms of a self-adjoint operator. In fact, $\|K\|_2$ is the square root of the trace of the self-adjoint operator K^*K , and the trace is the sum of the eigenvalues. However we do not need to calculate the eigenvalues, since, as we have seen, there are much easier ways to calculate the trace.

2.5 Problems

1. Let $H = L^2$ be the Hilbert space of square integrable functions on the line. Create an example of a Hilbert-Schmidt operator that is not an interpolation operator.
2. Let $H = L^2$ be the Hilbert space of square integrable functions on the line. Create an example of an interpolation operator that is not a Hilbert-Schmidt operator. Make the example so that the operator is not compact.
3. Find an example of a compact bounded operator on $H = L^2$ that is neither a Hilbert-Schmidt or an interpolation operator.
4. Find an example of a bounded operator on $H = L^2$ that is neither a Hilbert-Schmidt nor an interpolation operator, and that is also not a compact operator.

5. Let H be a Hilbert space. Give an example of a Hilbert-Schmidt operator for which the spectral radius is equal to the uniform norm.
6. Let H be a Hilbert space. Give an example of a Hilbert-Schmidt operator for which the spectral radius is very different from the uniform norm.
7. Let H be a Hilbert space. Is it possible for a Hilbert-Schmidt operator to have its Hilbert-Schmidt norm equal to its uniform norm? Describe all possible such situations.

2.6 Finite rank operators

Let H be a Hilbert space. A bounded linear transformation $K : H \rightarrow H$ is said to be finite rank if its range is finite dimensional. The dimension of the range of K is called the rank of K . A finite rank operator may be represented in the form

$$Kf = \sum_j z_j(u_j, f), \quad (2.35)$$

where the sum is finite. In a more abbreviated notation we could write

$$K = \sum_j z_j u_j^*. \quad (2.36)$$

Thus in L^2 this is an integral operator with kernel

$$k(x, y) = \sum_j z_j(x) \overline{u_j(y)}. \quad (2.37)$$

If K and L are finite rank operators, then so is their sum $K + L$. If K is finite rank and L is bounded, then KL and LK are finite rank.

The adjoint K^* of a finite rank operator K is finite rank. The two operators have the same rank.

Every finite rank operator is Hilbert-Schmidt and hence compact. If K is a compact operator, then there exists a sequence of finite rank operators K_n such that $\|K_n - K\|_\infty \rightarrow 0$ as $n \rightarrow \infty$.

If K is a finite rank operator and $\mu \neq 0$, then the calculation of $(\mu I - K)^{-1}$ or of $(I - \lambda K)^{-1}$ may be reduced to a finite-dimensional matrix problem in the finite dimensional space $R(K)$. This is because K leaves $R(K)$ invariant. Therefore if $\mu = 1/\lambda$ is not an eigenvalue of K acting in $R(K)$, then there is an inverse $(I - \lambda K)^{-1}$ acting in $R(K)$. However this gives a corresponding inverse in the original Hilbert space, by the formula

$$(I - \lambda K)^{-1} = I + (I - \lambda K)^{-1} \lambda K. \quad (2.38)$$

Explicitly, to solve

$$u = \lambda K u + f, \quad (2.39)$$

write

$$u = (I - \lambda K)^{-1}f = f + (I - \lambda K)^{-1}\lambda Kf. \quad (2.40)$$

To solve this, let w be the second term on the right hand side, so that $u = f + w$. Then $(I - \lambda K)w = \lambda Kf$. Write $w = \sum_j a_j z_j$. Then

$$\sum_j a_j z_j - \lambda \sum_j z_j(u_j, \sum_r a_r z_r) = \lambda \sum_j z_j(u_j, f). \quad (2.41)$$

Thus

$$a_j - \lambda \sum_r (u_j, z_r) a_r = (u_j, f). \quad (2.42)$$

This is a matrix equation that may be solved whenever $1/\lambda$ is not an eigenvalue of the matrix with entries (u_j, z_r) .

2.7 Problems

It may help to recall that the problem of inverting $I - \lambda K$ is the same as the problem of showing that $\mu = 1/\lambda$ is not in the spectrum of K .

1. Consider functions in $L^2(-\infty, \infty)$. Consider the integral equation

$$f(x) - \lambda \int_{-\infty}^{\infty} \cos(\sqrt{x^2 + y^4}) e^{-|x|-|y|} f(y) dy = g(x).$$

It is claimed that there exists $r > 0$ such that for every complex number λ with $|\lambda| < r$ the equation has a unique solution. Prove or disprove. Interpret this as a statement about the spectrum of a certain linear operator.

2. Consider functions in $L^2(-\infty, \infty)$. Consider the integral equation

$$f(x) - \lambda \int_{-\infty}^{\infty} \cos(\sqrt{x^2 + y^4}) e^{-|x|-|y|} f(y) dy = g(x).$$

It is claimed that there exists $R < \infty$ such that for every complex number λ with $|\lambda| > R$ the equation does not have a unique solution. Prove or disprove. Interpret this as a statement about the spectrum of a certain linear operator.

3. Consider functions in $L^2(-\infty, \infty)$. Consider the integral equation

$$f(x) - \lambda \int_{-\infty}^{\infty} e^{-|x|-|y|} f(y) dy = g(x).$$

Find all complex numbers λ for which this equation has a unique solution. Find the solution. Interpret this as a statement about the spectrum of a certain linear operator.

4. Consider functions in $L^2(0, 1)$. Consider the integral equation

$$f(x) - \lambda \int_0^x f(y) dy = g(x).$$

Find all complex numbers λ for which this equation has a unique solution. Find the solution. Interpret this as a statement about the spectrum of a certain linear operator. Hint: Differentiate. Solve a first order equation with a boundary condition.

5. Consider functions in $L^2(0, 1)$. Consider the integral equation

$$f(x) - \lambda \left[\int_0^x y(1-x)f(y) dy + \int_x^1 x(1-y)f(y) dy \right] = g(x).$$

Find all complex numbers λ for which this equation has a unique solution. Interpret this as a statement about the spectrum of a certain linear operator. Hint: The integral operator K has eigenfunctions $\sin(n\pi x)$. Verify this directly. This should also determine the eigenvalues.

Chapter 3

Densely Defined Closed Operators

3.1 Introduction

This chapter deals primarily with densely defined closed operators. Each everywhere defined bounded operator is in particular a densely defined closed operator.

If L is a densely defined closed operator, then so is its adjoint L^* . Furthermore, $L^{**} = L$. If both L and L^* have trivial null spaces, then both L^{-1} and L^{*-1} are densely defined closed operators.

A complex number λ is in the resolvent set of a densely defined closed operator L if $(L - \lambda I)^{-1}$ is an everywhere defined bounded operator. A complex number λ is in the spectrum of L if it is not in the resolvent set. A complex number λ is in the spectrum of L if and only if $\bar{\lambda}$ is in the spectrum of the adjoint L^* .

It is common to divide the spectrum into three disjoint subsets: point spectrum, continuous spectrum, and residual spectrum. (This terminology is misleading, in that it treats limits of point spectrum as continuous spectrum.) In this treatment we divide the spectrum into four disjoint subsets: standard point spectrum, pseudo-continuous spectrum, anomalous point spectrum, and residual spectrum. The adjoint operation maps the first two kind of spectra into themselves, but it reverses the latter two.

3.2 Subspaces

Let H be a Hilbert space. Let M be a vector subspace of H . The closure \bar{M} is also a vector subspace of H . The subspace M is said to be closed if $M = \bar{M}$. The orthogonal complement M^\perp is a closed subspace of H . Furthermore, $(\bar{M})^\perp = M^\perp$. Finally $M^{\perp\perp} = \bar{M}$. The nicest subspaces are closed subspaces.

For a closed subspace M we always have $M^{\perp\perp} = M$.

A subspace M is dense in H if $\bar{M} = H$. This is equivalent to the condition $M^\perp = \{0\}$.

3.3 Graphs

Let H be a Hilbert space. The direct sum $H \oplus H$ with itself consists of all ordered pairs $[u, v]$, where u, v are each vectors in H . The inner product of $[u, v]$ with $[u', v']$ is

$$([u, v], [u', v']) = (u, u') + (v, v'). \quad (3.1)$$

A graph is a linear subspace of $H \oplus H$. If we have two graphs L_1 and L_2 and if $L_1 \subset L_2$, then we say that L_1 is a *restriction* of L_2 or L_2 is an *extension* of L_1 .

If L is a graph, then its domain $D(L)$ is the set of all u in H such that there exists a v with $[u, v]$ in L . Its range $R(L)$ is the set of all v in H such that there exists u with $[u, v]$ in L .

If L is a graph, then its null space $N(L)$ is the set of all u in H such that $[u, 0]$ is in L .

If L is a graph, then the inverse graph L^{-1} consists of all $[v, u]$ such that $[u, v]$ is in L .

We say that a graph L is an operator if $[0, v]$ in L implies $v = 0$. It is easy to see that this is equivalent to saying that $N(L^{-1}) = \{0\}$ is the zero subspace. When L is an operator and $[u, v]$ is in L , then we write $Lu = v$. We shall explore the properties of operators in the next section.

Write \bar{L} for the closure of L . We say L is closed if $L = \bar{L}$.

If L is a graph, then the adjoint graph L^* consists of the pairs $[w, z]$ such that for all $[u, v]$ in L we have $(z, u) = (w, v)$.

If L is a graph, then the adjoint graph L^* is always closed. Furthermore, the adjoint of its closure \bar{L} is the same as the adjoint of L .

Remark: One way to think of the adjoint graph is to define the negative inverse $-L^{-1}$ of a graph L to consist of all the ordered pairs $[v, -u]$ with $[u, v]$ in L . Then the adjoint L^* is the orthogonal complement in $H \oplus H$ of $-L^{-1}$. That is, the pair $[z, w]$ in the graph L^* is orthogonal to each $[v, -u]$ with $[u, v]$ in the graph of L . This says that $([z, w], [v, -u]) = (z, u) - (w, v) = 0$ for all such $[u, v]$. [This says to take the graph with negative reciprocal slope, and then take the perpendicular graph to that.]

Another way to think of this is to define the anti-symmetric form $\omega([z, w], [u, v]) = (z, u) - (w, v)$. Then the adjoint A^* consists of the orthogonal complement of A with respect to ω .

Theorem. If L is a graph, then $L^{**} = \bar{L}$.

Perhaps the nicest general class of graphs consists of the closed graphs L . For such a graph the adjoint L^* is a graph, and $L^{**} = L$.

It is not hard to check that $L^{*-1} = L^{-1*}$.

Theorem. $N(L^*) = R(L)^\perp$.

Corollary. $\overline{R(L)} = N(L^*)^\perp$.

This corollary is very important in the theory of linear equations. Let L be a linear operator. Suppose that $R(L)$ is a closed subspace of H . Then in this special case the corollary says that $R(L) = N(L^*)^\perp$. Thus a linear equation $Lu = g$ has a solution u if and only if g is orthogonal to all solutions v of $L^*v = 0$. This is called the Fredholm alternative.

3.4 Operators

If L is a graph, then L is an operator if $[0, v]$ in L implies $v = 0$. It is easy to see that L is an operator precisely when $N(L^{-1}) = \{0\}$ is the zero subspace. When L is an operator and $[u, v]$ is in L , then we write $Lu = v$.

Corollary. L is densely defined if and only if L^* is an operator.

Proof: Apply the last theorem of the previous section to L^{-1} . This gives $N(L^{*-1}) = D(L)^\perp$.

Corollary. \bar{L} is an operator if and only if L^* is densely defined.

Proof: Apply the previous corollary to L^* and use $L^{**} = \bar{L}$.

An operator L is said to be closable if \bar{L} is also an operator. For a densely defined closable operator the adjoint L^* is a densely defined closed operator, and $L^{**} = \bar{L}$. Furthermore, the definition of the adjoint is that w is in the domain of L^* and $L^*w = z$ if and only if for all u in the domain of L we have $(z, u) = (w, Lu)$, that is,

$$(L^*w, u) = (w, Lu). \quad (3.2)$$

Perhaps the nicest general class of operators consists of the densely defined closed operators L . For such an operator the adjoint L^* is a densely defined closed operator, and $L^{**} = L$.

3.5 The spectrum

Theorem. (Closed graph theorem) Let H be a Hilbert space. Let L be a closed operator with domain $D(L) = H$. Then L is a bounded operator. (The converse is obvious.)

0. Let L be a closed operator. We say that λ is in the resolvent set of L if $N(L - \lambda I) = \{0\}$ and $R(L - \lambda I) = H$. In that case $(L - \lambda I)^{-1}$ is a closed operator with domain $D((L - \lambda I)^{-1}) = H$. By the closed graph theorem, $(L - \lambda I)^{-1}$ is a bounded operator.

We shall usually refer to $(L - \lambda I)^{-1}$ as the resolvent of L . However in some contexts it is convenient to use instead $(\lambda I - L)^{-1}$, which is of course just the negative. There is no great distinction between these two possible definitions of resolvent. However it is important to be alert to which one is being used.

1. Let L be a closed operator. We say that λ is in the standard point spectrum of L if $N(L - \lambda I) \neq \{0\}$ and $R(L - \lambda I)^\perp \neq \{0\}$.

2. Let L be a closed operator. We say that λ is in the anomalous point spectrum of L if $N(L - \lambda I) \neq \{0\}$ and $R(L - \lambda I)^\perp = \{0\}$ (that is, $R(L - \lambda I)$ is dense in H).

3. Let L be a closed operator. We say that λ is in the pseudo-continuous spectrum of L if $N(L - \lambda I) = \{0\}$ and $R(L - \lambda I)^\perp = \{0\}$ (so $R(L - \lambda I)$ is dense in H) but $R(L - \lambda I) \neq H$. In that case $(L - \lambda I)^{-1}$ is a closed operator with dense domain $D((L - \lambda I)^{-1})$ not equal to H .

4. Let L be a closed operator. We say that λ is in the residual spectrum of L if $N(L - \lambda I) = \{0\}$ and $R(L - \lambda I)^\perp \neq \{0\}$. In that case $(L - \lambda I)^{-1}$ is a closed operator with a domain that is not dense.

Theorem. Let L be a densely defined closed operator and let L^* be its adjoint operator. Then:

0. The number λ is in the resolvent set of L if and only if $\bar{\lambda}$ is in the resolvent set of L^* .

1. The number λ is in the standard point spectrum of L if and only if $\bar{\lambda}$ is in the standard point spectrum of L^* .

2. The number λ is in the anomalous point spectrum of L if and only if $\bar{\lambda}$ is in the residual spectrum of L^* .

3. λ is in the pseudo-continuous spectrum of L if and only if $\bar{\lambda}$ is in the pseudo-continuous spectrum of L^* .

4. λ is in the residual spectrum of L if and only if $\bar{\lambda}$ is in the anomalous point spectrum of L^* .

For finite dimensional vector spaces only cases 0 and 1 can occur.

Summary: Let L be a closed, densely defined operator. The complex number λ is in the point spectrum of L is equivalent to λ being an eigenvalue of L . Similarly, λ in the pseudo-continuous spectrum of L is equivalent to $(L - \lambda I)^{-1}$ being a densely defined, closed, but unbounded operator. Finally, λ in the residual spectrum of L is equivalent to $(L - \lambda I)^{-1}$ being a closed operator that is not densely defined.

3.6 Spectra of inverse operators

Consider a closed, densely defined operator with an inverse L^{-1} that is also a closed, densely defined operator. Let $\lambda \neq 0$. Then λ is in the resolvent set of L if and only if $1/\lambda$ is in the resolvent set of L^{-1} . In fact, we have the identity

$$(I - \lambda^{-1}L)^{-1} + (I - \lambda L^{-1})^{-1} = I. \quad (3.3)$$

One very important situation is when $K = L^{-1}$ is a compact operator. Then we know that all non-zero elements μ of the spectrum of $K = L^{-1}$ are eigenvalues of finite multiplicity, with zero as their only possible accumulation point. It follows that all elements $\lambda = 1/\mu$ of the spectrum of L are eigenvalues of finite multiplicity, with infinity as their only possible accumulation point.

3.7 Problems

If K is a bounded everywhere defined operator, then in particular K is a closed densely defined operator.

If K is a closed densely defined operator, and if both K and K^* have trivial nullspaces, then $L = K^{-1}$ is also a closed densely defined operator.

1. Let $K = L^{-1}$ be as above. Let $\lambda \neq 0$ and let $\mu = 1/\lambda$. Find a formula relating the resolvent $(L - \lambda)^{-1}$ to the resolvent $(K - \mu)^{-1}$.
2. Consider functions in $L^2(0, 1)$. Consider the integral operator K given by

$$(Kf)(x) = \int_0^x f(y) dy.$$

Show that $L = K^{-1}$ exists and is closed and densely defined. Describe the domain of L . Be explicit about boundary conditions. Describe how L acts on the elements of this domain. Show that $L^* = K^{*-1}$ is closed and densely defined. Describe the domain of L^* . Describe how L^* acts on the elements of this domain. Hint: Differentiate.

3. In the preceding problem, find the spectrum of L . Also, find the resolvent $(L - \lambda)^{-1}$ of L . Hint: Solve a first order linear ordinary differential equation.
4. Consider functions in $L^2(0, 1)$. Consider the integral operator K given by

$$(Kf)(x) = \left[\int_0^x y(1-x)f(y) dy + \int_x^1 x(1-y)f(y) dy \right].$$

Show that $L = K^{-1}$ exists and is closed and densely defined. Describe the domain of L . Be explicit about boundary conditions. Describe how L acts on the elements of this domain. Hint: Differentiate twice.

5. In the preceding problem, find the spectrum of L . Find the resolvent $(L - \lambda)^{-1}$ of L . Hint: Use $\sin(\sqrt{\lambda}x)$ and $\sin(\sqrt{\lambda}(1-x))$ as a basis for the solutions of a homogeneous second order linear ordinary differential equation. Solve the inhomogeneous equation by variation of parameters.
6. Let K be a compact operator. Suppose that K and K^* have trivial nullspaces, so that $L = K^{-1}$ is a closed densely defined operator. Prove that the spectrum of $L = K^{-1}$ consists of isolated eigenvalues of finite multiplicity. To what extent does this result apply to the examples in the previous problems?
7. Let K be a compact self-adjoint operator. Suppose that K has trivial null-space, so that $L = K^{-1}$ is a self-adjoint operator. Prove that there exists an orthogonal basis consisting of eigenvectors of L . To what extent does this result apply to the examples in the previous problems?

3.8 Self-adjoint operators

It is difficult to do algebraic operations with closed, densely defined operators, because their domains may differ. It is always true that $(zL)^* = \bar{z}L^*$. If K is bounded everywhere defined, then $(L+K)^* = L^*+K^*$ and $(K+L)^* = K^*+L^*$. Furthermore, if K is bounded everywhere defined, then $(KL)^* = L^*K^*$.

An operator A is self-adjoint if $A = A^*$. A self-adjoint operator is automatically closed and densely defined. (Every adjoint A^* is automatically closed. If A^* is an operator, then A is densely defined.)

If a self-adjoint operator has trivial null space, then its inverse is also a self-adjoint operator.

If L is a closed, densely defined operator, then L^*L is defined on the domain consisting of all u in $D(L)$ such that Lu is in $D(L^*)$. It is not obvious that this is closed and densely defined, much less that it is self-adjoint. However this is all a consequence of the following theorem.

Theorem. If L is a closed and densely defined operator, then L^*L is a self-adjoint operator.

Proof: If L is a closed and densely defined operator, then L^* is also a closed and densely defined operator. Furthermore, LL^* is an operator with $LL^* \subset (LL^*)^*$.

The Hilbert space $H \oplus H$ may be written as the direct sum of the two closed graphs $-L^{-1}$ and L^* . Therefore an arbitrary $[0, h]$ for h in H may be written as the sum $[0, h] = [-Lf, f] + [g, L^*g]$. This says that $0 = -Lf + g$ and $h = f + L^*g$. As a consequence $h = f + L^*Lf$. Furthermore, by properties of projections we have $\|Lf\|^2 + \|f\|^2 \leq \|h\|^2$. We have shown that for each h we can solve $(I + L^*L)f = h$ and that $\|f\|^2 \leq \|h\|^2$. Thus $(I + L^*L)^{-1}$ is everywhere defined and is a bounded operator with norm bounded by one.

Since $L^*L \subset (L^*L)^*$, we have $(I + L^*L)^{-1} \subset (I + L^*L)^{-1*}$. It follows that $(I + L^*L)^{-1} = (I + L^*L)^{-1*}$ is a self-adjoint operator. The conclusion follows.

3.9 First order differential operators with a bounded interval: point spectrum

In this section we shall see examples of operators with no spectrum at all. However we shall also see a very pretty and useful example of an operator with standard point spectrum. This operator is the one behind the theory of Fourier series.

Example 1A: This example is one where the correct number of boundary conditions are imposed. In the case of a first order differential operator this number is one. Let H be the Hilbert space $L^2(0, 1)$. Let L_0 be the operator d/dx acting on functions of the form $f(x) = \int_0^x g(y) dy$ where g is in H . The value of L_0 on such a function is $g(x)$. Notice that functions in the domain of L_0 automatically satisfy the boundary condition $f(0) = 0$. This is an example of a closed operator. The reason is that $(L_0^{-1}g)(x) = \int_0^x g(y) dy$. This is a bounded operator defined on the entire Hilbert space. So L_0^{-1} and L_0 are both closed.

The adjoint of the inverse is given by $(L_0^{-1*})h(x) = \int_x^1 h(y) dy$. It follows that L_0^* is the operator $-L_1$, where L_1 is given by d/dx acting on functions of the form $f(x) = -\int_x^1 g(y) dy$ where g is in H . The value of L_1 on such a function is $g(x)$. Notice that functions in the domain of L_1 automatically satisfy the boundary condition $f(1) = 0$.

The operators L_0 and L_1 each have one boundary condition. They are negative adjoints of each other. They each have a spectral theory, but it is extremely pathological. For instance, the resolvent of L_0 is given by

$$((L_0 - \lambda I)^{-1}g)(x) = \int_0^x e^{\lambda(x-y)}g(y) dy.$$

So there are no points at all in the spectrum of L_0 . It is in some sense located all at infinity.

To see this, consider the operator L_0^{-1} . This operator has spectrum consisting of the point zero. All the spectral information is hidden at this one point. This is, by the way, an example of pseudo-continuous spectrum.

This is one important though somewhat technical point. The domain of L_0 consists precisely of the functions in the range of $K_0 = L_0^{-1}$. In the example where K_0 is the integration operator, this is all functions of the form

$$u(x) = \int_0^x f(y)dy, \quad (3.4)$$

where f is in $L^2(0, 1)$. These functions u need not be C^1 . They belong to a larger class of functions that are indefinite integrals of L^2 functions. Such functions are continuous, but they may have slope discontinuities. The functions u of course satisfy the boundary condition $u(0) = 0$. The action of L_0 on a function u is given by $L_0u = u'$, where the derivative exists except possibly on a set of measure zero. However L^2 functions such as f are defined only up to sets of measure zero, so this is not a problem.

Now for a really picky question: If u is also regarded as an L^2 function, then it is also defined only up to sets of measure zero. So what does $u(0)$ mean? After all, the set consisting of 0 alone is of measure zero. The answer is that the general indefinite integral is a function of the form

$$u(x) = \int_0^x f(y)dy + C. \quad (3.5)$$

Among all L^2 functions given by such an integral expression, there is a subclass of those for which $C = 0$. These are the ones satisfying the boundary condition $u(0) = 0$. There is another subclass for which $C = -\int_0^1 f(y)dy$. These are the ones satisfying $u(1) = 0$.

A densely defined operator L is said to be self-adjoint if $L = L^*$. Similarly, L is said to be skew-adjoint if $L = -L^*$.

Example 1B: Here is another example with the correct number of boundary conditions. Let H be the Hilbert space $L^2(0, 1)$. Let $L_=-$ be the operator d/dx

acting on functions of the form $f(x) = \int_0^x g(y) dy + C$ where g is in H and $\int_0^1 g(y) dy = 0$. The value of $L_=-$ on such a function is $g(x)$. Notice that functions in the domain of $L_=-$ automatically satisfy the boundary condition $f(0) = f(1)$. The operator $L_=-$ is skew-adjoint.

The operator $L_=-$ has periodic boundary conditions. Since it is skew-adjoint, it has an extraordinarily nice spectral theory. The resolvent is

$$((L_=- - \lambda I)^{-1}g)(x) = \frac{1}{1 - e^\lambda} \int_0^x e^{\lambda(x-y)} g(y) dy - \frac{1}{1 - e^{-\lambda}} \int_x^1 e^{-\lambda(y-x)} g(y) dy.$$

The spectrum consists of the numbers $2\pi in$. These are all point spectrum. The corresponding eigenvectors form a basis that gives the Fourier series expansion of an arbitrary periodic function with period one.

A densely defined operator L is said to be *Hermitian* if $L \subset L^*$. This is simply the algebraic property that

$$(Lu, w) = (u, Lw)$$

for all u, w in $D(L)$. Similarly, L is said to be skew-Hermitian if $L \subset -L^*$.

Example 2: The following example illustrates what goes wrong when one imposes the wrong number of boundary conditions. Let H be the Hilbert space $L^2(0, 1)$. Let L_{01} be the operator d/dx acting on functions of the form $f(x) = \int_0^x g(y) dy$ where g is in H and $\int_0^1 g(y) dy = 0$. The value of L_{01} on such a function is $g(x)$. Notice that functions in the domain of L_{01} automatically satisfy the boundary conditions $f(0) = 0$ and $f(1) = 0$. The adjoint of L_{01} is the operator $-L$, where L is given by d/dx acting on functions of the form $\int_0^x g(y) dy + C$, where g is in H . The value of L on such a function is $g(x)$. Notice that functions in the domain of L need not satisfy any boundary conditions. From this we see that L_{01} is skew-Hermitian.

The operator L is d/dx . It has too few boundary conditions. The operator L_{01} has a boundary condition at 0 and at 1. This is too many boundary conditions. Each of these operators is the negative of the adjoint of the other. The spectrum of L consists of the entire complex plane, and it is all point spectrum. The spectrum of L_{01} also consists of the entire complex plane, and it is all residual spectrum.

Remark: The operators L_0 and L_1 have $L_{01} \subset L_0 \subset L$ and with $L_{01} \subset L_1 \subset L$. Furthermore, the operator $L_=-$ has $L_{01} \subset L_=- \subset L$. Thus there are various correct choices of boundary conditions, but they may have different spectral properties.

3.10 Spectral projection and reduced resolvent

Consider a closed densely defined operator L . In this section we shall assume that the eigenvectors of L span the entire Hilbert space. Consider also an isolated eigenvalue λ_n . The spectral projection corresponding to λ_n is a (not

necessarily orthogonal) projection onto the corresponding eigenspace. It is given in terms of the resolvent by

$$P_n = \lim_{\lambda \rightarrow \lambda_n} (\lambda_n - \lambda)(L - \lambda I)^{-1}. \quad (3.6)$$

This is the negative of the residue of the resolvent at λ_n . The reason this works is that $L = \sum_m \lambda_m P_m$ and consequently

$$(L - \lambda I)^{-1} = \sum_m \frac{1}{\lambda_m - \lambda} P_m, \quad (3.7)$$

at least in the case under consideration, when the eigenvectors span the entire Hilbert space.

The reduced resolvent corresponding to λ_n is defined as the operator that inverts $(L - \lambda_n I)$ in the range of $(I - P_n)$ and is zero in the range of P_n . It is a solution of the equations

$$S_n P_n = P_n S_n = 0 \quad (3.8)$$

and

$$(L - \lambda_n I) S_n = 1 - P_n \quad (3.9)$$

It may be expressed in terms of the resolvent by

$$S_n = \lim_{\lambda \rightarrow \lambda_n} (L - \lambda I)^{-1} (1 - P_n). \quad (3.10)$$

When the eigenvectors span this is

$$S_n = \sum_{m \neq n} \frac{1}{\lambda_m - \lambda_n} P_m. \quad (3.11)$$

Example: Take the skew-adjoint operator $L = d/dx$ acting in $L^2(0, 1)$ with periodic boundary conditions. The spectral projection corresponding to eigenvalue $2\pi i n$ is the self-adjoint operator

$$(P_n g)(x) = \int_0^1 \exp(2\pi i n(x - y)) g(y) dy. \quad (3.12)$$

The reduced resolvent corresponding to eigenvalue 0 is the skew-adjoint operator

$$(S_0 g)(x) = \int_0^x \left(\frac{1}{2} - x + y\right) g(y) dy + \int_x^1 \left(-\frac{1}{2} - x + y\right) g(y) dy. \quad (3.13)$$

3.11 Generating second-order self-adjoint operators

This section exploits the theorem that says that if L is an arbitrary closed densely defined operator, then L^*L is a self-adjoint operator. Remember that $L^{**} = L$, so LL^* is also a self-adjoint operator.

It would be easy to conclude that first order differential operators such as L_{01} and its adjoint L are of no interest for spectral theory. This is not the case. From the general theorem LL_{01} and $L_{01}L$ are self-adjoint second-order differential operators. These, as we shall see, have a nice spectral theory. The operator LL_{01} is the operator $-d^2/dx^2$ with Dirichlet boundary conditions $u(0) = 0$ and $u(1) = 0$. The operator $L_{01}L$ is the operator $-d^2/dx^2$ with Neumann boundary conditions $u'(0) = 0$ and $u'(1) = 0$. It is amusing to work out other self-adjoint second order differential operators that may be generated from the first order differential operators of the preceding sections.

3.12 First order differential operators with a semi-infinite interval: residual spectrum

In this example we shall see examples of operators with anomalous point spectrum and residual spectrum. These operators underly the theory of the Laplace transform.

Example: Let H be the Hilbert space $L^2(0, \infty)$. Let L_0 be the operator d/dx acting on functions f in H of the form $f(x) = \int_0^x g(y) dy$ where g is in H . The value of L_0 on such a function is $g(x)$. Notice that functions in the domain of L_0 automatically satisfy the boundary condition $f(0) = 0$.

If $\Re\lambda < 0$, then we can always find a solution of the equation $(L_0 - \lambda I)f = g$. This solution is

$$f(x) = \int_0^x e^{\lambda(x-y)} g(y) dy. \quad (3.14)$$

This equation defines the bounded operator $(L_0 - \lambda I)^{-1}$ that sends g into f , at least when $\Re\lambda < 0$. Notice that if $\Re\lambda > 0$, then the formula gives a result in L^2 only if g is orthogonal to $e^{-\bar{\lambda}x}$. Thus $\Re\lambda > 0$ corresponds to residual spectrum.

Again let H be the Hilbert space $L^2(0, \infty)$. Let L be the operator d/dx with no boundary condition. If $\Re\lambda > 0$, then we can always find a solution of the equation $(L - \lambda I)f = g$. This solution is

$$f(x) = - \int_x^\infty e^{\lambda(x-y)} g(y) dy. \quad (3.15)$$

This equation defines the bounded operator $(L - \lambda I)^{-1}$ that sends g into f , at least when $\Re\lambda > 0$. On the other hand, if $\Re\lambda < 0$, then we have point spectrum.

The relation between these two operators is $L_0^* = -L$. This corresponds to the fact that $(L_0 - \lambda I)^{-1*} = (-L - \bar{\lambda})^{-1}$, which is easy to check directly.

3.13 First order differential operators with an infinite interval: continuous spectrum

In this section we shall see an example of an operator with continuous spectrum. This is the example that underlies the theory of the Fourier transform.

Example: Let H be the Hilbert space $L^2(-\infty, \infty)$. Let L be the operator d/dx acting on functions f in H with derivatives f' in H .

If $\Re\lambda < 0$, then we can find always find a solution of the equation $(L - \lambda I)f = g$. This solution is

$$f(x) = \int_{-\infty}^x e^{\lambda(x-y)} g(y) dy. \quad (3.16)$$

This equation defines the bounded operator $(L - \lambda I)^{-1}$ that sends g into f , at least when $\Re\lambda < 0$. If $\Re\lambda > 0$, then we can find always find a solution of the equation $(L - \lambda I)f = g$. This solution is

$$f(x) = - \int_x^{\infty} e^{\lambda(x-y)} g(y) dy. \quad (3.17)$$

This equation defines the bounded operator $(L - \lambda I)^{-1}$ that sends g into f , at least when $\Re\lambda > 0$.

The operator L is skew-adjoint, that is, $L^* = -L$. This corresponds to the fact that $(L - \lambda I)^{-1*} = (-L - \bar{\lambda})^{-1}$, which is easy to check directly.

3.14 Problems

- Let $H = L^2(0, 1)$. Let $L = -id/dx$ with periodic boundary conditions. Find an explicit formula for $(\lambda - L)^{-1}g$. Hint: Solve the first order ordinary differential equation $(\lambda - L)f = g$ with the boundary condition $f(0) = f(1)$.
- Find the eigenvalues and eigenvectors of L . For each eigenvalue λ_n , find the residue P_n of $(\lambda - L)^{-1}$ at λ_n .
- Find the explicit form of the formula $g = \sum_n P_n g$.
- Let $H = L^2(-\infty, \infty)$. Let $L = -id/dx$. Let k be real and $\epsilon > 0$. Find an explicit formula for $(L - k - i\epsilon)^{-1}g$. Also, find an explicit formula for $(L - k + i\epsilon)^{-1}g$. Find the explicit form of the expression

$$\delta_\epsilon(L - k)g = \frac{1}{2\pi i} [(L - k - i\epsilon)^{-1} - (L - k + i\epsilon)^{-1}]g.$$

- Find the explicit form of the formula

$$g = \int_{-\infty}^{\infty} \delta_\epsilon(L - k)g dk.$$

- Let $\epsilon \rightarrow 0$. Find the explicit form of the formula

$$g = \int_{-\infty}^{\infty} \delta(L - k)g dk.$$

3.15 A pathological example

Consider $H = L^2(\mathbb{R})$ and fix g in H . Define K as the integral operator with kernel $k(x, y) = g(x)\delta(y)$. Consider the domain $D(K)$ to be the set of all continuous functions in H . Then

$$(Ku)(x) = g(x)f(0). \quad (3.18)$$

This K is densely defined but not closed. It has a closure \bar{K} , but this is not an operator. To see this, let $u_n \rightarrow u$ and $Ku_n = u_n(0)g \rightarrow v$. Then the pair $[u, v]$ is in the graph \bar{K} . But we can take $u_n \rightarrow 0$ in the Hilbert space sense, yet with each $u_n(0) = C$. So this gives the pair $[0, Cg]$ in the graph \bar{K} . This is certainly not an operator!

Consider the adjoint K^* . This is the integral operator with kernel $\delta(x)\overline{g(y)}$. That is,

$$(K^*w)(x) = \delta(x) \int_{-\infty}^{\infty} \overline{g(y)}w(y) dy. \quad (3.19)$$

Since

$$\int_{-\infty}^{\infty} \delta(x)^2 dx = +\infty, \quad (3.20)$$

the $\delta(x)$ is not in L^2 . Hence the domain of K^* consists of all w with $(g, w) = 0$, and $K^* = 0$ on this domain. This is an operator that is closed but not densely defined. According to the general theory, its adjoint is $K^{**} = \bar{K}$, which is not an operator.

Chapter 4

Normal operators

4.1 Spectrum of a normal operator

Theorem (von Neumann) Let L be a densely defined closed operator. Then L^*L and LL^* are each self-adjoint operators.

A densely defined closed operator is said to be normal if $L^*L = LL^*$.

There are three particularly important classes of normal operators.

1. A self-adjoint operator is an operator L with $L^* = L$.
2. A skew-adjoint operator is an operator L with $L^* = -L$.
3. A unitary operator is an operator L with $L^* = L^{-1}$. A unitary operator is bounded.

For a self-adjoint operator the spectrum is on the real axis. For a skew-adjoint operator the spectrum is on the imaginary axis. For a unitary operator the spectrum is on the unit circle.

For normal operators there is a different classification of spectrum. Let L be a normal operator acting in a Hilbert space H . The point spectrum consists of the eigenvalues of L . The corresponding eigenvectors span a closed subspace M_p of the Hilbert space. The spectrum of L in this space consists of either what we have previously called standard point spectrum or of what we have previously called pseudo-continuous spectrum. This kind of pseudo-continuous spectrum is not really continuous at all, since it consists of limits of point spectrum.

Let M_c be the orthogonal complement in H of M_p . Then the spectrum of L restricted to M_c is called the continuous spectrum of L . In our previous classification the spectrum of L in this space would be pseudo-continuous spectrum.

With this classification for normal operators the point spectrum and continuous spectrum can overlap. But they really have nothing to do with each other, since they take place in orthogonal subspaces.

Spectral theorem for compact normal operators. Let K be a compact normal operator. (This includes the cases of self-adjoint operators and skew-adjoint operators.) Then K has an orthogonal basis of eigenvectors. The non-zero

eigenvalues have finite multiplicity. The only possible accumulation point of eigenvalues is zero.

Spectral theorem for normal operators with compact resolvent. Let L be a normal operator with compact resolvent. (This includes the cases of self-adjoint operators and skew-adjoint operators.) Then L has an orthogonal basis of eigenvectors. The eigenvalues have finite multiplicity. The only possible accumulation point of eigenvalues is infinity.

4.2 Problems

1. Perhaps the most beautiful self-adjoint operator is the spherical Laplacian

$$\Delta_S = \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \sin \theta \frac{\partial}{\partial \theta} + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2}.$$

Show by explicit computation that this is a Hermitian operator acting on L^2 of the sphere with surface measure $\sin \theta d\theta d\phi$. Pay explicit attention to what happens at the north pole and south pole when one integrates by parts.

2. Let r be the radius satisfying $r^2 = x^2 + y^2 + z^2$. Let

$$L = r \frac{\partial}{\partial r}$$

be the Euler operator. Show that the Laplace operator

$$\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$$

is related to L and Δ_S by

$$\Delta = \frac{1}{r^2} [L(L+1) + \Delta_S].$$

3. Let p be a polynomial in x, y, z that is harmonic and homogeneous of degree ℓ . Thus $\Delta p = 0$ and $Lp = \ell p$. Such a p is called a solid spherical harmonic. Show that each solid spherical harmonic is an eigenfunction of Δ_S and find the corresponding eigenvalue as a function of ℓ .
4. The restriction of a solid spherical harmonic to the sphere $r^2 = 1$ is called a surface spherical harmonic. The surface spherical harmonics are the eigenfunctions of Δ_S . Show that surface spherical harmonics for different values of ℓ are orthogonal in the Hilbert space of L^2 functions on the sphere.
5. The dimension of the eigenspace indexed by ℓ is $2\ell + 1$. For $\ell = 0$ the eigenspace is spanned by 1. For $\ell = 1$ it is spanned by z , $x + iy$, and $x - iy$. For $\ell = 2$ it is spanned by $3z^2 - r^2$, $z(x + iy)$, $z(x - iy)$, $(x + iy)^2$,

and $(x - iy)^2$. For $\ell = 3$ it is spanned by $5z^3 - 3zr^2$, $(5z^2 - r^2)(x + iy)$, $(5z^2 - r^2)(x - iy)$, $z(x + iy)^2$, $z(x - iy)^2$, $(x + iy)^3$, $(x - iy)^3$. Express the corresponding surface spherical harmonics in spherical coordinates.

6. In the case $\ell = 1$ we can write the general spherical harmonic as $ax + by + cz$. In the case $\ell = 2$ we can write it as $ax^2 + by^2 + cz^2 + dxy + eyz + fzx$ with an additional condition on the coefficients. What is this condition? In the case $\ell = 3$ we can write it as $a_1x^3 + b_1y^2x + c_1z^2x + a_2y^3 + b_2z^2y + c_2x^2y + a_3z^3 + b_3x^2z + c_3y^2z + dxyz$ with additional conditions. What are they?

4.3 Variation of parameters and Green's functions

First look at first order linear ordinary differential operators. Let $Lu = p(x)u' + r(x)u$. Let u_1 be a non-zero solution of the homogeneous equation $Lu = 0$. The general solution of the homogeneous equation $Lu = 0$ is $u(x) = c_1u_1(x)$, where c_1 is a parameter. The method of variation of parameters gives a solution of the inhomogeneous equation $Lu = f$ in the form $u(x) = c_1(x)u_1(x)$.

The condition on the parameter is given by plugging u into $Lu = f$. This gives $p(x)c_1'(x)u_1(x) = f(x)$. The solution is $c_1'(x) = f(x)/(p(x)u_1(x))$. The only difficult part is to integrate this to get the general solution

$$u(x) = \int_a^x \frac{u_1(x)}{p(y)u_1(y)} f(y) dy + Cu_1(x). \quad (4.1)$$

Now look at second order linear ordinary differential operators. Let $Lu = p(x)u'' + r(x)u' + q(x)u$. Let u_1 and u_2 be independent solutions of the homogeneous equation $Lu = 0$. The general solution of the homogeneous equation $Lu = 0$ is $u(x) = c_1u_1(x) + c_2u_2(x)$, where c_1 and c_2 are parameters. The method of variation of parameters gives a solution of the inhomogeneous equation $Lu = f$ in the form $u(x) = c_1(x)u_1(x) + c_2(x)u_2(x)$. Not only that, it has the property that the derivative has the same form, that is, $u'(x) = c_1(x)u_1'(x) + c_2(x)u_2'(x)$.

If this is to be so, then $c_1'(x)u_1(x) + c_2'(x)u_2(x) = 0$. This is the first equation. The second equation is given by plugging u into $Lu = f$. This gives $p(x)(c_1'u_1'(x) + c_2'u_2'(x)) = f(x)$. This system of two linear equations is easily solved. Let $w(x) = u_1(x)u_2'(x) - u_2(x)u_1'(x)$. The solution is $c_1'(x) = -u_2(x)f(x)/(p(x)w(x))$ and $c_2'(x) = u_1(x)f(x)/(p(x)w(x))$.

A solution of $Lu = f$ is thus

$$u(x) = \int_x^b \frac{u_1(x)u_2(y)}{p(y)w(y)} f(y) dy + \int_a^x \frac{u_2(x)u_1(y)}{p(y)w(y)} f(y) dy. \quad (4.2)$$

Furthermore,

$$u'(x) = \int_x^b \frac{u_1'(x)u_2(y)}{p(y)w(y)} f(y) dy + \int_a^x \frac{u_2'(x)u_1(y)}{p(y)w(y)} f(y) dy. \quad (4.3)$$

Notice that $u(a) = Au_1(a)$ and $u'(a) = Au'_1(a)$, while $u(b) = Bu_2(b)$ and $u'(b) = Bu'_2(b)$. So this form of the solution is useful for specifying boundary conditions at a and b .

The general solution is obtained by adding an arbitrary linear combination $C_1u_1(x) + C_2u_2(x)$. However often we want a particular solution with boundary conditions at a and b . Then we use the form above. This can also be written

$$u(x) = (Kf)(x) = \int_a^b k(x, y)f(y) dy, \quad (4.4)$$

where

$$k(x, y) = \begin{cases} \frac{u_1(x)u_2(y)}{p(y)w(y)} & \text{if } x < y \\ \frac{u_2(x)u_1(y)}{p(y)w(y)} & \text{if } x > y \end{cases} \quad (4.5)$$

Sometime one thinks of y as a fixed source and write the equation

$$L_x k(x, y) = \delta(x - y). \quad (4.6)$$

Of course this is just another way of saying that $LK = I$.

4.4 Second order differential operators with a bounded interval: point spectrum

Example. Consider the self-adjoint differential operator $L_D = -d^2/dx^2$ on $L^2(0, 1)$ with Dirichlet boundary conditions $f(0) = 0$ and $f(1) = 0$ at 0 and 1. Take the solutions $u_1(x) = \sin(\sqrt{\lambda}x)/\sqrt{\lambda}$ and $u_2(x) = \sin(\sqrt{\lambda}(1-x))/\sqrt{\lambda}$. These are defined in a way that does not depend on which square root of λ is taken. (Furthermore, they have obvious values in the limit $\lambda \rightarrow 0$.) Then $p(x) = -1$ and $w(x) = -\sin(\sqrt{\lambda})/\sqrt{\lambda}$. This also does not depend on the cut. The resolvent is thus $((L_D - \lambda)^{-1}g)(x) = f(x)$ where

$$f(x) = \frac{1}{\sqrt{\lambda} \sin(\sqrt{\lambda})} \left[\int_0^x \sin(\sqrt{\lambda}(1-x)) \sin(\sqrt{\lambda}y)g(y) dy + \int_x^1 \sin(\sqrt{\lambda}x) \sin(\sqrt{\lambda}(1-y))g(y) dy \right]. \quad (4.7)$$

The spectrum consists of the points $\lambda = n^2\pi^2$ for $n = 1, 2, 3, \dots$. This is standard point spectrum. It is amusing to work out the spectral projection at the eigenvalue $n^2\pi^2$. This is the negative of the residue and is explicitly

$$(P_n g)(x) = 2 \int_0^1 \sin(n\pi x) \sin(n\pi y)g(y) dy. \quad (4.8)$$

Example. Consider the self-adjoint differential operator $L_N = -d^2/dx^2$ on $L^2(0, 1)$ with Neumann boundary conditions $f'(0) = 0$ and $f'(1) = 0$ at 0 and 1. Take the solutions $u_1(x) = \cos(\sqrt{\lambda}x)$ and $u_2(x) = \cos(\sqrt{\lambda}(1-x))$. Then $p(x) = -1$ and $w(x) = \sqrt{\lambda} \sin(\sqrt{\lambda})$. This also does not depend on the cut. The

resolvent is thus

$$((L_N - \lambda)^{-1}g)(x) = -\frac{1}{\sqrt{\lambda} \sin(\sqrt{\lambda})} \left[\int_0^x \cos(\sqrt{\lambda}(1-x)) \cos(\sqrt{\lambda}y) g(y) dy + \int_x^1 \cos(\sqrt{\lambda}x) \cos(\sqrt{\lambda}(1-y)) g(y) dy \right]. \quad (4.9)$$

The spectrum consists of the points $\lambda = n^2\pi^2$ for $n = 0, 1, 2, 3, \dots$. This is standard point spectrum. The spectral projection at the eigenvalue $n^2\pi^2$ for $n = 1, 2, 3, \dots$ is

$$(P_n g)(x) = 2 \int_0^1 \cos(n\pi x) \cos(n\pi y) g(y) dy. \quad (4.10)$$

For $n = 0$ it is

$$(P_0 g)(x) = \int_0^1 g(y) dy. \quad (4.11)$$

It is interesting to compute the reduced resolvent of L_N at the eigenvalue 0. Thus we must compute $(L_N - \lambda)^{-1}\bar{g}$, where $\bar{g} = (1 - P)g$ has zero average, and then let λ approach zero. This is easy. Expand the cosine functions to second order. The constant terms may be neglected, since they are orthogonal to \bar{g} . This gives

$$(S_0 \bar{g})(x) = \int_0^x \left(\frac{1}{2}(1-x)^2 + \frac{1}{2}y^2 \right) \bar{g}(y) dy + \int_x^1 \left(\frac{1}{2}x^2 + \frac{1}{2}(1-y)^2 \right) \bar{g}(y) dy. \quad (4.12)$$

From this it is easy to work out that

$$(S_0 g)(x) = \int_0^x \left(\frac{1}{2}(1-x)^2 + \frac{1}{2}y^2 - \frac{1}{6} \right) g(y) dy + \int_x^1 \left(\frac{1}{2}x^2 + \frac{1}{2}(1-y)^2 - \frac{1}{6} \right) g(y) dy. \quad (4.13)$$

4.5 Second order differential operators with a semibounded interval: continuous spectrum

Example. Consider the self-adjoint differential operator $L_D = -d^2/dx^2$ on $L^2(0, \infty)$ with Dirichlet boundary condition $f(0) = 0$ at 0. Take the solutions $u_1(x) = \sinh(\sqrt{-\lambda}x)/\sqrt{-\lambda}$ and $u_2(x) = e^{-\sqrt{-\lambda}x}$. Since $\sinh(iz) = i \sin(z)$, this is the same $u_1(x)$ as before. In $u_2(x)$ the square root is taken to be cut on the negative axis. Then $p(x) = -1$ and $w(x) = -1$. The resolvent is

$$((L_D - \lambda)^{-1}g)(x) = \frac{1}{\sqrt{-\lambda}} \left[\int_0^x e^{-\sqrt{-\lambda}x} \sinh(\sqrt{-\lambda}y) g(y) dy + \int_x^\infty \sinh(\sqrt{-\lambda}x) e^{-\sqrt{-\lambda}y} g(y) dy \right]. \quad (4.14)$$

The spectrum consists of the positive real axis and is continuous.

It is instructive to compute the resolvent of the self-adjoint differential operator $L_N = -d^2/dx^2$ on $L^2(0, \infty)$ with Neumann boundary condition $f'(0) = 0$ at 0. Again the spectrum consists of the positive real axis and is continuous.

4.6 Second order differential operators with an infinite interval: continuous spectrum

Example. Consider the self-adjoint differential operator $L = -d^2/dx^2$ on $L^2(-\infty, \infty)$. There is now no choice of boundary conditions. The resolvent is

$$((L-\lambda)^{-1}g)(x) = \frac{1}{2\sqrt{-\lambda}} \left[\int_{-\infty}^x e^{-\sqrt{-\lambda}x} e^{\sqrt{-\lambda}y} g(y) dy + \int_x^{\infty} e^{\sqrt{-\lambda}x} e^{-\sqrt{-\lambda}y} g(y) dy \right]. \quad (4.15)$$

This can also be written in the form

$$((L-\lambda)^{-1}g)(x) = \frac{1}{2\sqrt{-\lambda}} \int_{-\infty}^{\infty} e^{-\sqrt{-\lambda}|x-y|} g(y) dy. \quad (4.16)$$

The spectrum consists of the positive real axis and is continuous.

4.7 The spectral theorem for normal operators

Throughout the discussion we make the convention that the inner product is conjugate linear in the first variable and linear in the second variable.

The great theorem of spectral theory is the following.

Let H be a Hilbert space. Let L be a normal operator. Then there exists a set K (which may be taken to be a disjoint union of copies of the line) and a measure μ on K and a unitary operator $U : H \rightarrow L^2(K, \mu)$ and a complex function λ on K such that

$$(ULf)(k) = \lambda(k)(Uf)(k). \quad (4.17)$$

Thus if we write Λ for the operator of multiplication by the function λ , we get the representation

$$L = U^* \Lambda U. \quad (4.18)$$

The theorem is a generalization of the theorem on diagonalization of normal matrices. If the measure μ is discrete, then the norm in the space $L^2(\mu)$ is given by $\|g\|^2 = \sum_k |g(k)|^2 \mu(\{k\})$. The λ_k are the eigenvalues of L . The equation then says

$$(ULf)_k = \lambda_k (Uf)_k. \quad (4.19)$$

The unitary operator U is given by

$$(Uf)_k = (\psi_k, f), \quad (4.20)$$

where the ψ_k are eigenvectors of L normalized so that $\mu(\{k\}) = 1/(\psi_k, \psi_k)$. The inverse of U is given by

$$U^* g = \sum_k g_k \psi_k \mu(\{k\}). \quad (4.21)$$

The equation

$$Lf = U^* \Lambda U f \quad (4.22)$$

says explicitly that

$$Lf = \sum_k \lambda_k(\psi_k, f) \psi_k \mu(\{k\}). \quad (4.23)$$

If the measure μ is continuous, then the norm in the space $L^2(K, \mu)$ is given by $\|g\|^2 = \int |g(k)|^2 d\mu(k)$. Then $\lambda(k)$ is a function of the continuous parameter k . The equation then says

$$(ULf)(k) = \lambda(k)(Uf)(k). \quad (4.24)$$

In quite general contexts the unitary operator U is given by

$$(Uf)(k) = (\psi_k, f), \quad (4.25)$$

but now this equation only makes sense for a dense set of f in the Hilbert space, and the ψ_k resemble eigenvectors of L , but do not belong to the Hilbert space, but instead to some larger space, such as a space of slowly growing functions or of mildly singular distributions. The inverse of U is given formally by

$$U^*g = \int g(k) \psi_k d\mu(k), \quad (4.26)$$

but this equation must be interpreted in some weak sense. The equation

$$Lf = U^* \Lambda U f \quad (4.27)$$

says formally that

$$Lf = \int \lambda(k) (\psi_k, f) \psi_k d\mu(k). \quad (4.28)$$

Since the eigenvectors ψ_k are not in the Hilbert space, it is convenient in many contexts to forget about them and instead refer to the measure μ and the function λ and to the operators U and U^* . The theorem says simply that every normal operator L is isomorphic to multiplication by a function λ . The simplicity and power of the equation $L = U^* \Lambda U$ cannot be overestimated.

The spectral theorem for normal operators says that every normal operator is isomorphic (by a unitary operator mapping the Hilbert space to an L^2 space) to a multiplication operator (multiplication by some complex valued function λ). The spectrum is the essential range of the function. This is the set of points w such that for each $\epsilon > 0$ the set of all points k such that $\lambda(k)$ is within ϵ of w has measure > 0 . This is obvious; a function $\lambda(k)$ has $1/(\lambda(k) - w)$ bounded if and only if w is not in the essential range of $\lambda(k)$.

4.8 Examples: compact normal operators

The theorem on compact normal operators is a corollary of the general spectral theorem. Consider a compact normal operator L . Say that it is isomorphic to multiplication by λ . Fix $\epsilon > 0$ and look at the part of the space where $|L| \geq \epsilon$. This is just the part of the space that is isomorphic to the part of L^2 where $|\lambda| \geq \epsilon$. More explicitly, this is the subspace consisting of all functions g in L^2 such that $g(k) \neq 0$ only where $|\lambda(k)| \geq \epsilon$. On this part of the space the operator L maps the unit ball onto a set that contains the ball of radius ϵ . Since L is compact, it follows that this part of the space is finite dimensional. This shows that the spectrum in the subspace where $|L| \geq \epsilon$ is finite dimensional. Therefore there are only finitely many eigenvectors of finite multiplicity in this space. Since $\epsilon > 0$ is arbitrary, it follows that there are only countably many isolated eigenvectors of finite multiplicity in the part of the space where $|L| > 0$. In the part of the space where $L = 0$ we can have an eigenvalue 0 of arbitrary multiplicity (zero, finite, or infinite).

4.9 Examples: translation invariant operators and the Fourier transform

The nicest examples for the continuous case are given by translation invariant operators acting in $H = L^2(\mathbb{R}, dx)$. In this case the Fourier transform maps H into $L^2(\mathbb{R}, dk/(2\pi))$. The Fourier transform is given formally by

$$(Ff)(k) = (\psi_k, f) = \int_{-\infty}^{\infty} e^{-ikx} f(x) dx. \quad (4.29)$$

Here $\psi_k(x) = e^{ikx}$, and we are using the convention that the inner product is linear in the second variable. The inverse Fourier transform is

$$(F^{-1}g)(x) = \int_{-\infty}^{\infty} e^{ikx} g(k) \frac{dk}{2\pi}. \quad (4.30)$$

Here are some examples:

Example 1: Translation. Let U_a be defined by

$$(T_a f)(x) = f(x - a). \quad (4.31)$$

Then T_a is unitary. The spectral representation is given by the Fourier transform. In fact

$$(FT_a f)(k) = \exp(-ika)(Ff)(k). \quad (4.32)$$

Example 2: Convolution. Let C be defined by

$$(Cf)(x) = \int_{-\infty}^{\infty} c(x - y)f(y) dy = \int_{-\infty}^{\infty} c(a)f(x - a) da, \quad (4.33)$$

where c is an integrable function. Then C is bounded normal. Then by integrating the first example we get

$$(FCf)(k) = \hat{c}(k)(Ff)(k), \quad (4.34)$$

where \hat{c} is the Fourier transform of c .

Example 3: Differentiation. Let D be defined by

$$(Df)(x) = \frac{df(x)}{dx}. \quad (4.35)$$

Then D is skew-adjoint. Furthermore, we get

$$(FDf)(k) = ik(Ff)(k). \quad (4.36)$$

Notice that the unitary operator in Example 1 may be written

$$U_a = \exp(-aD). \quad (4.37)$$

Example 4. Second differentiation. Let D^2 be defined by

$$(D^2f)(x) = \frac{d^2f(x)}{dx^2}. \quad (4.38)$$

Then D is self-adjoint. Furthermore, we get

$$(FDf)(k) = -k^2(Ff)(k). \quad (4.39)$$

We can take interesting functions of these operator. For instance $(-D^2 + m^2)^{-1}$ is convolution by $1/(2m)e^{-m|x|}$. And $\exp(tD^2)$ is convolution by $1/\sqrt{2\pi t}e^{-\frac{x^2}{2t}}$.

4.10 Examples: Schrödinger operators

If $V(x)$ is a real locally integrable function that is bounded below, then

$$H = -D^2 + V(x) \quad (4.40)$$

is a well-defined self-adjoint operator. Such an operator is called a Schrödinger operator.

If $V(x) \rightarrow \infty$ as $|x| \rightarrow \infty$, then the spectrum of H is point spectrum. Finding the eigenvalues is a challenge. One case where it is possible to obtain explicit formulas is when $V(x)$ is a quadratic function.

Also there is an interesting limiting case. If $V(x) = 0$ for $0 < x < 1$ and $V(x) = +\infty$ elsewhere, then we may think of this as the operator $H = -D^2$ with Dirichlet boundary conditions at the end points of the unit interval. We know how to find the spectrum in this case.

If on the other hand, $V(x)$ is integrable on the line, then the spectrum of H consists of positive continuous spectrum and possibly some strictly negative eigenvalues. A nice example of this is the square well, where there is a constant

$a > 0$ with $V(x) = -a$ for $0 < x < 1$ and $V(x) = 0$ otherwise. This is another case where computations are possible.

The calculation of the spectral properties of Schrödinger operators is the main task of quantum physics. However we shall see that Schrödinger operators play a role in other contexts as well. (One example will be in calculus of variations.)

4.11 Subnormal operators

The spectral theorem for normal operators is a landmark. However not every operator is normal. One important class of operators with fascinating spectral properties consists of the subnormal operators. A subnormal operator is an operator that is the restriction of a normal operator to an invariant subspace. First consider the case of bounded operators. An operator $S : H \rightarrow H$ is subnormal if there exists a larger Hilbert space H' with $H \subset H'$ as a closed subspace, a normal operator $N : H' \rightarrow H'$ that leaves H invariant, and such that N restricted to H is S .

Example: Let $H = L^2([0, \infty), dt)$. For each $a \geq 0$ define

$$(S_a f)(t) = f(t - a) \quad (4.41)$$

for $a \leq t$ and

$$(S_a f)(t) = 0 \quad (4.42)$$

for $0 \leq t < a$. Then S_a is subnormal.

To see this, consider the bigger space $H' = L^2(\mathbb{R}, dt)$ and consider H as the subspace of functions that vanish except on the positive reals. Let U_a be translation by a on H' . If $a \geq 0$, then the subspace H is left invariant by U_a . Then S_a is U_a acting in this subspace.

It follows from the spectral theorem that a subnormal operator S is isomorphic by a unitary operator $U : H \rightarrow M$ to a multiplication operator that sends $g(k)$ to $\lambda(k)g(k)$. Here M is a closed subspace of an L^2 space. For each f in H we have Uf in M and $USf(k) = \lambda(k)(Uf)(k)$ in M .

Now it is more difficult to characterize the spectrum. A number w is in the resolvent set if $1/(\lambda(k) - w)g(k)$ is in M for every function $g(k)$ in M . However it is not sufficient that this is a bounded function. If, for instance, every function $g(k)$ in M has an extension to an analytic function $g(z)$ defined on some larger region, then one would want $1/(\lambda(z) - w)g(z)$ to also be an analytic function in this region. So we need to require also that w is not in the range of the extension $\lambda(z)$.

Example: Let F be the Fourier transform applied to the Hilbert space H of L^2 functions that vanish except on the positive axis. Then the image of this transform consists of the subspace M of L^2 functions $g(k)$ that are boundary values of analytic functions $g(z)$ in the lower half plane. The operator S_a for $a > 0$ is isomorphic to multiplication by $\lambda(k) = \exp(-iak)$ acting in this subspace. This function extends to a function $\lambda(z) = \exp(-iaz)$ defined in the lower half

plane. So the spectrum is the range of this function. But the image of $\Im z \leq 0$ under $\exp(-iaz)$ is the unit circle $|w| \leq 1$. So this is the spectrum.

The adjoint of a subnormal operator $S : H \rightarrow H$ is another operator $S^* : H \rightarrow H$. The adjoint S^* need not be subnormal.

Lemma. If $N : H' \rightarrow H'$ is normal, $H \subset H'$, and $S : H \rightarrow H$ is the restriction of N to H , then $S^* : H \rightarrow H$ is given by $S^* = PN^*$, where P is the orthogonal projection of H' onto H .

Proof: If u is in H , for each v in H we have

$$(S^*u, v) = (u, Sv) = (u, Nv) = (N^*u, v). \quad (4.43)$$

This says that $N^*u - S^*u$ is orthogonal to every v in H . Since S^*u is in H , this implies that S^*u is the orthogonal projection of N^*u onto H .

Theorem. If S is a subnormal operator, then $SS^* \leq S^*S$ as quadratic forms, that is, $(u, SS^*u) \leq (u, S^*Su)$ for all u in H .

Proof: First note that for u in H' we have

$$(N^*u, N^*u) = (u, NN^*u) = (u, N^*Nu) = (Nu, Nu). \quad (4.44)$$

Then for u in H we have

$$(S^*u, S^*u) = (PN^*u, PN^*u) \leq (N^*u, N^*u) = (Nu, Nu) = (Su, Su). \quad (4.45)$$

Corollary. If S is a subnormal operator and $Su = 0$, then $S^*u = 0$. Thus the null space of S is contained in the null space of S^* .

Corollary. If S is a subnormal operator and $Su = \lambda u$, then $S^*u = \bar{\lambda}u$. Thus every eigenvector of S is an eigenvector of S^* .

It is not true that every eigenvalue of S^* is an eigenvalue of S . It is more typical that S^* has eigenvalues while S does not. In fact we shall see examples in which S^* has anomalous point spectrum, while S has residual spectrum.

Theorem. If the Hilbert space is finite dimensional, then every subnormal operator is normal.

Proof: Let S be a subnormal operator acting in H . Since the space H is finite dimensional, S has an eigenvector u in H with $Su = \lambda u$. Since S is subnormal, it follows that $S^*u = \bar{\lambda}u$. Let v be a vector in H that is orthogonal to u . Then $(Sv, u) = (v, S^*u) = \bar{\lambda}(v, u) = 0$. Thus the orthogonal complement of u in H is also an invariant space, so the operator S restricted to this smaller space is also subnormal. Continue in this way until one finds an orthogonal basis of eigenvectors for S .

There are examples in which one might want to consider unbounded subnormal operators. One possible definition might be the following. Consider an unbounded normal operator N . Thus there is a dense domain $D(N) \subset H'$ such that $N : D(N) \rightarrow H'$ is normal. Let H be a Hilbert space that is a closed subspace of H' . Suppose that $D(N) \cap H$ is dense in H and that N sends vectors in this dense subspace into H . Then if $D(S) = D(N) \cap H$ and S is the restriction of N to $D(S)$, the operator S is subnormal. Then $D(S)$ is dense in H , and $S : D(S) \rightarrow H$ is a closed, densely defined operator.

Example: Let $H = L^2([0, \infty), dt)$. Define

$$(Sf)(t) = f'(t) \quad (4.46)$$

on the domain consisting of all f in H such that f' is also in H and $f(0) = 0$. Then S is subnormal.

To see this, consider the bigger space $H' = L^2(\mathbb{R}, dt)$ and consider H as the subspace of functions that vanish except on the positive reals. Let N be differentiation on H' . Notice that if f is in $D(N)$ and is also in H , then $f(0) = 0$ automatically.

If N is a normal operator acting in H' , so $N : D(N) \rightarrow H'$, then its adjoint N^* is also a normal operator, and in fact $D(N^*) = D(N)$. So if $S : D(S) \rightarrow H$ is subnormal, then $D(S) = D(N) \cap H = D(N^*) \cap H$. Furthermore, if u is in $D(N^*) \cap H$, then u is in $D(S^*)$ and $S^*u = PN^*u$. This can be seen from the computation $(u, Sv) = (u, Nv) = (N^*u, v)$ for all v in $D(S)$.

We conclude that for a subnormal operator $D(S) \subset D(S^*)$ and $(S^*, S^*u) \leq (Su, Su)$ for all u in $D(S)$. This is also an easy computation: $(S^*u, S^*u) = (PN^*u, PN^*u) \leq (N^*u, N^*u) = (Nu, Nu) = (Su, Su)$.

Example: For the operator S in the last example we have

$$(S^*f)(t) = -f'(t) \quad (4.47)$$

on the domain consisting of all f in H such that f' is also in H . There is no boundary condition at zero. This is a case where $D(S) \subset D(S^*)$ and the two domains are not equal.

4.12 Examples: forward translation invariant operators and the Laplace transform

The example in the last section is the operator theory context for the theory of the Laplace transform.

The Laplace transform of a function f in $L^2([0, \infty), dt)$ is

$$(Lf)(z) = \int_0^\infty e^{-zt} f(t) dt. \quad (4.48)$$

If we think of $z = i\omega$ on the imaginary axis, then when regarded as a function of ω this is the Fourier transform. However it extends as an analytic function to ω in the lower half plane, that is, to z in the right half plane.

Let $a \geq 0$ and let S_a be the operator of right translation filling in with zero as defined in the previous section. Then

$$(LS_a f)(z) = \int_a^\infty e^{-zt} f(t-a) dt = e^{-az} (Lf)(z). \quad (4.49)$$

So S_a is isomorphic to multiplication by e^{-az} acting on the Hilbert space of functions analytic in the right half plane. Its spectrum consists of the closed unit disk.

4.12. EXAMPLES: FORWARD TRANSLATION INVARIANT OPERATORS AND THE LAPLACE TRANSFORM

Let c be an integrable function defined on the positive axis, and let the causal convolution C be defined by

$$(Cf)(t) = \int_0^t c(t-u)f(u) du = \int_0^t c(a)f(t-a) da. \quad (4.50)$$

Then C is subnormal, and

$$(LCf)(z) = \hat{c}(z)(Lf)(z), \quad (4.51)$$

where \hat{c} is the Laplace transform of c .

Let $D_0 = d/dt$ be differentiation with zero boundary conditions at the origin. Then

$$(LD_0f)(z) = z(Lf)(z). \quad (4.52)$$

Notice that the boundary condition is essential for integration by parts. The spectrum of D_0 consists of the closed right half plane. We can write $\exp(-aD_0) = S_a$ for $a \geq 0$. This operator satisfies the differential equation $dS_a f/da = -D_0 S_a f$ for $a \geq 0$, provided that f is in the domain of D_0 (and in particular satisfies the boundary condition).

The adjoint of a subnormal operator need not be subnormal. Thus, for instance, the adjoint of S_a is

$$(S_a^*f)(t) = f(t+a). \quad (4.53)$$

Its spectrum is also the closed unit disc, but the interior of the disk consists of point spectrum. Notice that the Laplace transform does not send this into a multiplication operator. In fact,

$$(LS_a^*f)(z) = \int_0^\infty e^{-zt} f(t+a) dt = e^{az} \left[(Lf)(z) - \int_0^a e^{-zt} f(t) dt \right]. \quad (4.54)$$

Similarly, the adjoint of D_0 is $-D$, where $D = d/dt$ with no boundary condition. Again the Laplace transform does not make this into a multiplication operator. In fact, we have

$$(LDf)(z) = z(Lf)(z) - f(0). \quad (4.55)$$

The spectrum of D consists of the closed left half plane. The interior consists of point spectrum. We can write $\exp(aD) = \exp(-aD_0^*) = S_a^*$ for $a \geq 0$. Even though D does not have a spectral representation as a multiplication operator, this operator satisfies the differential equation $dS_a^* f/da = DS_a^* f$ for $a \geq 0$, provided f is in the domain of D .

Example: Solve the differential equation $(D+k)f = g$, with $k > 0$, with boundary condition $f(0) = c$. The operator $D+k$ is not invertible, since $-k$ is an eigenvalue of D . However we can write $f = u - ce^{-kt}$ and solve

$$(D_0 + k)u = g. \quad (4.56)$$

The solution is $u = (D_0 + k)^{-1}g$. In terms of Laplace transforms this is $\hat{u}(z) = 1/(z + k)\hat{g}(z)$. It follows that u is given by the causal convolution

$$u(t) = \int_0^t e^{-k(t-u)}g(u) du. \quad (4.57)$$

Thus

$$f(t) = f(0)e^{-kt} + \int_0^t e^{-k(t-u)}g(u) du. \quad (4.58)$$

4.13 Quantum mechanics

Here is a dictionary of the basic concepts. Fix a Hilbert space. A *quantum observable* is a self-adjoint operator L . A *quantum state* is a unit vector u . The *expectation* of the observable L in the state u in $D(L)$ is

$$\mu = (u, Lu). \quad (4.59)$$

The *variance* of the observable L in the state u in $D(L)$ is

$$\sigma^2 = \|(L - \mu I)u\|^2. \quad (4.60)$$

Here are some observables. The Hilbert space is $L^2(R, dx)$. The *momentum observable* is $p = -i\hbar d/dx$. Here $\hbar > 0$ is Planck's constant. The *position observable* is q which is multiplication by the coordinate x . The *Heisenberg uncertainty principle* says that for every state the product $\sigma_p\sigma_q \geq \hbar/2$.

If L is an observable and f is a real function, then $f(L)$ is an observable. Take the case where the function is 1_A , the indicator function of a set A of position coordinate values. Then $1_A(q)$ has expectation

$$(u, 1_A(q)u) = \int_A |u(x)|^2 dx. \quad (4.61)$$

This is the probability that the position is in A . Similarly, take the case where the function is 1_B , the indicator function of a set B of momentum coordinate values. Since in the Fourier transform representation p is represented by multiplication by $\hbar k$, it follows that $1_B(p)$ has expectation

$$(u, 1_B(p)u) = \int_{\{\hbar k \in B\}} |\hat{u}(k)|^2 \frac{dk}{2\pi}. \quad (4.62)$$

This is the probability that the momentum is in B .

Energy observables are particularly important. The *kinetic energy observable* is $H_0 = p^2/(2m) = -\hbar^2/(2m)d^2/dx^2$. Here $m > 0$ is the mass. The spectrum of H_0 is the positive real axis. The *potential energy observable* is $V = v(q)$ which is multiplication by $v(x)$. Here v is a given real function that represents potential energy as a function of the space coordinate. The spectrum of V is the range of the function v . The *total energy observable* or *quantum Hamiltonian* is

$$H = H_0 + V = \frac{p^2}{2m} + v(q) = -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + v(x). \quad (4.63)$$

If we assume that the function v is bounded, then H is a self-adjoint operator. (In many cases when v is only bounded below it remains a self-adjoint operator.) The problem of investigating its spectrum is of the utmost importance for quantum mechanics. Since H_0 and V do not commute, this is not an easy problem.

Suppose that the total energy observable H is a self-adjoint operator. Then the *time evolution operator* is the unitary operator $\exp(-itH/\hbar)$. A central problem of quantum mechanics is to compute this operator. This is not easy, because while $\exp(-itH_0/\hbar)$ and $\exp(-itV/\hbar)$ are easy to compute, the operators H_0 and V do not commute. So there is no direct algebraic way to express $\exp(-itH/\hbar)$ in terms of the simpler operators. Nevertheless, we shall encounter a beautiful formula for this time evolution operator in the next chapter.

4.14 Problems

1. A particularly fascinating self-adjoint operator is the quantum harmonic oscillator. (This operator also occurs in disguised form in other contexts.) It is

$$N = \frac{1}{2} \left(-\frac{d^2}{dx^2} + x^2 - 1 \right)$$

acting in $L^2(-\infty, \infty)$. Show that it factors as

$$N = A^*A,$$

where

$$A = \frac{1}{\sqrt{2}} \left(x + \frac{d}{dx} \right)$$

and

$$A^* = \frac{1}{\sqrt{2}} \left(x - \frac{d}{dx} \right).$$

2. Show that $AA^* = A^*A + I$.
3. Solve the equation $Au_0 = 0$. Show that $Nu_0 = 0$.
4. Show that if $Nu_n = nu_n$ and $u_{n+1} = A^*u_n$, then $Nu_{n+1} = (n+1)u_{n+1}$. Thus the eigenvalues of N are the natural numbers. These are the standard type of point spectrum.
5. Show that each eigenfunction u_n is a polynomial in x times $u_0(x)$. Find the polynomials for the cases of $n = 0, 1, 2, 3$ explicitly (up to constant factors). Verify that each u_n belongs to the Hilbert space.
6. It may be shown that A^* is a subnormal operator and so A is the adjoint of a subnormal operator. Find all eigenvalues (point spectrum) of A . Find each corresponding eigenvector. Verify that it belongs to the Hilbert space.

7. Find all eigenvalues (point spectrum) of A^* . Find the spectrum of A^* . What kind of spectrum is it?
8. If A^* is indeed a subnormal operator, then we should have $A^*A \leq AA^*$ as quadratic forms. Is this the case?

Chapter 5

Calculus of Variations

5.1 The Euler-Lagrange equation

The problem is to find the critical points of

$$F(y) = \int_{x_1}^{x_2} f(y, y_x, x) dx. \quad (5.1)$$

The differential of F is

$$dF(y)h = \int_{x_1}^{x_2} (f_y h + f_{y_x} h_x) dx = \int_{x_1}^{x_2} (f_y - \frac{d}{dx} f_{y_x}) h dx + f_{y_x} h \Big|_{x_1}^{x_2}. \quad (5.2)$$

Thus for the differential to be zero we must have the Euler-Lagrange equation

$$f_y - \frac{d}{dx} f_{y_x} = 0. \quad (5.3)$$

This is an equation for the critical function y . It is second order, and it has the explicit form

$$f_y - f_{yy_x} \frac{dy}{dx} - f_{y_x y_x} \frac{d^2 y}{dx^2} - f_{xy_x} = 0. \quad (5.4)$$

This equation is linear in $d^2 y/dx^2$. However the coefficients are in general nonlinear expressions in y , dy/dx , and x .

If the y are required to have fixed values $y = y_1$ at $x = x_1$ and $y = y_2$ at $x = x_2$ at the end points, then the h are required to be zero at the end points. The corresponding boundary term is automatically zero.

If the y and h are free to vary at an end point, then at a critical point one must have $f_{y_x} = 0$ at at the end points.

Sometimes one wants to think of

$$\frac{\delta F}{\delta y(x)} = \frac{\partial f}{\partial y} - \frac{d}{dx} \frac{\partial f}{\partial y_x} \quad (5.5)$$

as the gradient of F , where the inner product is given by the integral. This expression is then known as the variational derivative. The Euler-Lagrange equation then says that the variational derivative is zero.

Example. Consider the problem of minimizing the length

$$F(y) = \int_{x_1}^{x_2} \sqrt{1 + y_x^2} dx \quad (5.6)$$

between the points (x_1, y_1) and (x_2, y_2) . The boundary conditions are $y(x_1) = y_1$ and $y(x_2) = y_2$. The solution of the Euler-Lagrange equation is $y_x = C$, a curve of constant slope. So the solution is $y - y_1 = m(x - x_1)$, where $m = (y_2 - y_1)/(x_2 - x_1)$.

Example. Consider the problem of minimizing the length

$$F(y) = \int_{x_1}^{x_2} \sqrt{1 + y_x^2} dx \quad (5.7)$$

between the lines $x = x_1$ and $x = x_2$. The boundary conditions for the Euler-Lagrange equation are $y_x(x_1) = 0$ and $y_x(x_2) = 0$. The solution of the Euler-Lagrange equation is $y_x = C$, a curve of constant slope. But to satisfy the boundary conditions $C = 0$. So the solution is $y = A$, where A is an arbitrary constant.

5.2 A conservation law

Say that one has a solution of the Euler-Lagrange equation. Then

$$\frac{d}{dx} f = f_y y_x + f_{y_x} y_{xx} + f_x = \frac{d}{dx} (y_x f_{y_x}) + f_x. \quad (5.8)$$

Thus

$$\frac{d}{dx} H + f_x = 0, \quad (5.9)$$

where

$$H = y_x f_{y_x} - f. \quad (5.10)$$

If $f_x = 0$, then this says that H is constant. This is the conservation law. It is a nonlinear equation for y and dy/dx . It takes the explicit form

$$y_x f_{y_x} - f = C. \quad (5.11)$$

Thus it is first order, but in general fully nonlinear in y_x . How in practice does one solve a problem in calculus of variations? There is a way when the function $f(y, y_x)$ does not depend on x . Use the conservation law $H(y, dy/dx) = C$ to solve for $dy/dx = \alpha(y)$. Perhaps this equation can be solved directly. Or write $dx = dy/\alpha(y)$. Integrate both sides to get x in terms of y . Or make a substitution expressing y in terms of a new variable u , and get x and y each in terms of u .

Example 1: This example comes from minimizing the area of a surface of revolution. The unknown is a function y of x between $-a$ and a . The value of y at $\pm a$ is r . The element of area is proportional to $2\pi y ds = 2\pi y \sqrt{1 + y_x^2} dx$. Let $f(y, y_x) = 2\pi y \sqrt{1 + y_x^2}$. Then $H = -2\pi y / \sqrt{1 + y_x^2} = -C$. The differential equation to be solved is $(dy/dx)^2 = (k^2 y^2 - 1)$, where $k = 2\pi/C$. This has solution $y = (1/k) \cosh(k(x - x_0))$. By symmetry $x_0 = 0$. So we need to solve $rk = \cosh(ak)$. This equation has to be solved for k . Fix r and vary a . When a is small enough, then $\cosh(ak)$ cuts the line rk in two points. Thus there are two solutions of the Euler-Lagrange equations, corresponding to $y = (1/k) \cosh(kx)$ with the two values of k . The derivative is $dy/dx = \sinh(kx)$. The smaller value of k gives the smaller derivative, so this is the minimum area surface satisfying the boundary conditions.

Example 2: This example comes up finding the maximum area for given arc length. For simplicity consider y as a function of x between $-a$ and a , with value 0 at the two end points. For simplicity let the arc length κ satisfy $2a < \kappa < \pi a$. The area is the integral from $-a$ to a of $y dx$, while the length is the integral from $-a$ to a of $ds = \sqrt{1 + y_x^2} dx$. This is a Lagrange multiplier problem. The function is $f(y, y_x) = y - \lambda \sqrt{1 + y_x^2}$. The conserved quantity is $H = -\lambda / \sqrt{1 + y_x^2} - y = -c_1$. The differential equation is $dx/dy = (y - c_1) / \sqrt{(\lambda^2 - (y - c_1)^2)}$. Thus $x - c_2 = \sqrt{\lambda^2 - (y - c_1)^2}$, which is the equation of a circle $(x - c_2)^2 + (y - c_1)^2 = \lambda^2$. The Lagrange multiplier turns out to be the radius of the circle. By symmetry $c_2 = 0$. Furthermore $c_1 = -\sqrt{\lambda^2 - a^2}$. If we let $\sin(\theta) = a/\lambda$, then the equation $\sin(\theta) < \theta < (\pi/2) \sin(\theta)$ translates into $2a < \kappa = 2\lambda\theta < \pi a$.

5.3 Second variation

Say that y is a solution of the Euler-Lagrange equation that has fixed values at the end points. Then the second differential of F is obtained by expanding $F(y + h) = F(y) + dF(y)h + \frac{1}{2}d^2F(y)(h, h) + \dots$. The result is

$$d^2F(y)(h, h) = \int_{x_1}^{x_2} (f_{yy}h^2 + 2f_{yy_x}hh_x + f_{y_x y_x}h_x^2) dx. \quad (5.12)$$

The functions h have value 0 at the end points, so we may freely integrate by parts. The result may thus also be written

$$d^2F(y)(h, h) = \int_{x_1}^{x_2} ((f_{yy} - \frac{d}{dx}f_{yy_x})h^2 + f_{y_x y_x}h_x^2) dx. \quad (5.13)$$

Yet another form is

$$d^2F(y)(h, h) = \int_{x_1}^{x_2} [-\frac{d}{dx}(f_{y_x y_x}h_x) + (f_{yy} - \frac{d}{dx}f_{yy_x})h]h dx = (Lh, h). \quad (5.14)$$

We recognize

$$Lh = -\frac{d}{dx}(f_{y_x y_x}h_x) + (f_{yy} - \frac{d}{dx}f_{yy_x})h \quad (5.15)$$

as a Sturm-Liouville operator. In fact, if $f(y, y_x, x) = \frac{1}{2}y_x^2 + g(y, x)$, then

$$Lh = -h_{xx} + g_{yy}h \quad (5.16)$$

is a Schrödinger operator. The coefficient g_{yy} has the solution of the Euler-Lagrange inserted, so it is regarded as a function of x . The operator has Dirichlet boundary conditions at the end points of the interval from x_1 to x_2 .

5.4 Interlude: The Legendre transform

The Legendre transform of a function L of v is defined as follows. Let $p = dL/dv$. Find the inverse function defining v as a function of p . Then the Legendre transform is a function H of p satisfying $dH/dp = v$. The constant of integration is chosen so that $H(0) + L(0) = 0$.

Example: Let $L = e^v - v - 1$. Then the derivative is $p = e^v - 1$. The inverse function is $v = \log(1 + p)$. The integral is $H = (1 + p)\log(1 + p) - p$.

There is a remarkable formula relating the Legendre transform of a function to the original function. Let L be a function of v . Define

$$p = \frac{dL}{dv}. \quad (5.17)$$

Then

$$H = pv - L. \quad (5.18)$$

is the Legendre transform of L . In the following we want to think of L as a function of the v , while H is a function of the dual variable p . The variable p is covariant and is dual to the variable v , which is contravariant. Thus in this formula v is defined in terms of p by solving the equation $p = dL/dv$ for v .

The fundamental theorem about the Legendre transform is

$$v = \frac{dH}{dp}. \quad (5.19)$$

Proof: Let $H = pv - L$ as above. By the product rule and the chain rule and the definition of p we get

$$\frac{dH}{dp} = v + p\frac{dv}{dp} - \frac{dL}{dv}\frac{dv}{dp} = v + p\frac{dv}{dp} - p\frac{dv}{dp} = v. \quad (5.20)$$

Thus the situation is symmetric, and one can go back from H to L in the same way as one got from L to H . Conclusion: Two functions are Legendre transforms of each other when their derivatives are inverse functions to each other.

One would like a condition that guarantees that the equations that define the Legendre transform actually have solutions. In order for $p = dL/dv$ to have a unique solution v , it would be useful to have dL/dv to be strictly increasing. This is the same as saying that $dp/dv = d^2L/dv^2 > 0$. The inverse function then has derivative given by the inverse $dv/dp = d^2H/dp^2 > 0$.

Example: Let $L = e^v - v - 1$. Then we have seen that $H = (p+1) \log(p+1) - p$. This also follows from the general formula $H = pv - L$. In fact, since $p = e^v - 1$ has inverse $v = \log(1+p)$, we have $H = pv - L = p \log(1+p) - [(1+p) - \log(1+p) - 1] = (1+p) \log(1+p) - p$. In this example the second derivatives are e^v and $1/(p+1)$. These are both positive, and they are reciprocals of each other.

Example: Let $L = v^a/a$ with $1 < a < \infty$ defined for $v \geq 0$. Then $H = p^b/b$ with $1 < b < \infty$ defined for $p \geq 0$. In this case $p = v^{a-1}$ and $v = p^{b-1}$. The relation between a and b is $(a-1)(b-1) = 1$. This may also be written as $1/a + 1/b = 1$. Thus a and b are conjugate exponents. The second derivatives are $(a-1)v^{a-2}$ and $(b-1)p^{b-2}$. Again they are reciprocals.

The Legendre transform plays a fundamental role in thermodynamics and in mechanics. Here are examples from mechanics involving kinetic energy.

Example: Start with $L = (1/2)mv^2$. Then $p = mv$. So $v = p/m$ and $H = pv - L = p^2/(2m)$. This H is the Legendre transform of L .

Example: If we start instead with $H = p^2/(2m)$, then $v = p/m$. So $p = mv$ and $L = pv - H = (1/2)mv^2$. The Legendre transform of L brings us back to the original H .

Example: A famous relativistic expression for energy is $H = \sqrt{m^2c^4 + p^2c^2}$. It is an amusing exercise to compute that $L = -mc^2\sqrt{1 - v^2/c^2}$. The key is the relation $p = mv/\sqrt{1 - v^2/c^2}$ between the derivatives.

The Legendre transform has a generalization to several dimensions. Let L be a function of v_1, \dots, v_n . Define

$$p_j = \frac{\partial L}{\partial v_j}, \quad (5.21)$$

so $dL = \sum_j p_j dv_j$. (This is the differential of L , so it is a one-form.) Then

$$H = \sum_k p_k v_k - L. \quad (5.22)$$

is the Legendre transform of L . In the following we want to think of L as a function of the v_j , while H is a function of the dual variables p_k . The variables p_k are covariant and dual to the variable v_j , which are contravariant. Thus in this formula the v_j are defined in terms of p_k by solving the equation $p_k = \partial L / \partial v_k$ for the v_j .

The fundamental theorem about the Legendre transform in this context is

$$v_k = \frac{\partial H}{\partial p_k}. \quad (5.23)$$

Proof: Let $H = \sum_k p_k v_k - L$ as above. By the product rule and the chain rule and the definition of p we get

$$\frac{\partial H}{\partial p_j} = v_j + \sum_k p_k \frac{\partial v_k}{\partial p_j} - \sum_k \frac{\partial L}{\partial v_k} \frac{\partial v_k}{\partial p_j} = v_j + \sum_k p_k \frac{\partial v_k}{\partial p_j} - \sum_k p_k \frac{\partial v_k}{\partial p_j} = v_j. \quad (5.24)$$

Thus the situation is symmetric, and one can go back from H to L in the same way as one got from L to H . Conclusion: Two functions are Legendre

transforms of each other when their derivatives are inverse functions to each other.

One would like a condition that guarantees that the equations that define the Legendre transform actually have solutions. In order for the equation to have a solution locally, the matrix

$$\frac{\partial p_k}{\partial v_j} = \frac{\partial^2 L}{\partial v_k \partial v_j} \quad (5.25)$$

should be invertible. A particularly nice condition is that this matrix is strictly positive definite. The inverse matrix is then

$$\frac{\partial v_j}{\partial p_k} = \frac{\partial^2 H}{\partial p_j \partial p_k}. \quad (5.26)$$

It is then also strictly positive definite.

5.5 Lagrangian mechanics

One is given a Lagrangian function L that is a function of position q and velocity \dot{q} and possibly time t . The problem is to find the critical points of

$$S(q) = \int_{t_1}^{t_2} L(q, \dot{q}, t) dt. \quad (5.27)$$

The function q represents position as a function of time. It has fixed values q_1 and q_2 at the end points t_1 and t_2 . The differential of S is

$$dS(q)h = \int_{t_1}^{t_2} (L_q h + L_{\dot{q}} h_t) dt = \int_{t_1}^{t_2} (L_q - \frac{d}{dt} L_{\dot{q}}) h dt. \quad (5.28)$$

Here h is a function with values 0 at the end points. Thus for the differential to be zero we must have the Euler-Lagrange equation

$$\frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}} = 0. \quad (5.29)$$

Say that one has a solution of the Euler-Lagrange equation. Then

$$\frac{d}{dt} L = L_q \dot{q} + L_{\dot{q}} \ddot{q} + L_t = \frac{d}{dt} (\dot{q} L_{\dot{q}}) + L_t. \quad (5.30)$$

Thus along a solution

$$\frac{d}{dt} H + \frac{\partial L}{\partial t} = 0, \quad (5.31)$$

where

$$H = \dot{q} L_{\dot{q}} - L. \quad (5.32)$$

If $L_t = 0$, then this says that H is constant. This is the energy conservation law.

Define

$$p = \frac{\partial L}{\partial \dot{q}}. \quad (5.33)$$

This is the momentum variable. Then the Euler-Lagrange equation says that along a solution

$$\frac{dp}{dt} = \frac{\partial L}{\partial q}. \quad (5.34)$$

5.6 Hamiltonian mechanics

Let

$$H = p\dot{q} - L. \quad (5.35)$$

be the Legendre transform of L , where \dot{q} is defined in terms of p implicitly by $p = \partial L / \partial \dot{q}$.

In the following we want to think of L as a function of q and \dot{q} and possibly t , while H is a function of q and p and possibly t . The momentum variable p is covariant and is dual to the velocity variable \dot{q} , which is contravariant. However p is expressed in terms of q and \dot{q} by solving the Legendre transform equation $p = \partial L / \partial \dot{q}$. According to the properties of the Legendre transform we have

$$\dot{q} = \frac{\partial H}{\partial p}. \quad (5.36)$$

In fact, the proof is simple if we remember that q is fixed:

$$\frac{\partial H}{\partial p} = \dot{q} + p \frac{\partial \dot{q}}{\partial p} - \frac{\partial L}{\partial \dot{q}} \frac{\partial \dot{q}}{\partial p} = \dot{q}. \quad (5.37)$$

It also follows that

$$\frac{\partial H}{\partial q} = -\frac{\partial L}{\partial q}. \quad (5.38)$$

This also has an easy proof if we remember that p is fixed and so \dot{q} depends on q :

$$\frac{\partial H}{\partial q} = p \frac{\partial \dot{q}}{\partial q} - \frac{\partial L}{\partial q} - \frac{\partial L}{\partial \dot{q}} \frac{\partial \dot{q}}{\partial q} = -\frac{\partial L}{\partial q}. \quad (5.39)$$

The Euler-Lagrange equation says that along a solution we have

$$\frac{dp}{dt} = -\frac{\partial H}{\partial q}. \quad (5.40)$$

Since $p = \partial L / \partial \dot{q}$ determines the function p in terms of q , dq/dt , and t , it follows that along a solution

$$\frac{dq}{dt} = \frac{\partial H}{\partial p} \quad (5.41)$$

determines the function dq/dt in terms of q , p , and t . The last two equations are Hamilton's equations. They are a first order system, linear in dq/dt and dp/dt , but nonlinear in q, p, t .

It is easy to check from Hamilton's equations that along a solution

$$\frac{dH}{dt} = \frac{\partial H}{\partial t}. \quad (5.42)$$

When H does not depend explicitly on t , then this has an integral in which H is constant. This integral is conservation of energy. It gives a situation in which p is related to q . This allows an equation in which dq/dt is related to q . This relation may be highly nonlinear.

5.7 Kinetic and potential energy

The most classical situation is when

$$L = \frac{1}{2}m\dot{q}^2 - V(q). \quad (5.43)$$

The Euler-Lagrange equation is just.

$$-V'(q) - \frac{dq_t}{dt} = 0. \quad (5.44)$$

This is Newton's law of motion.

In this problem the momentum is $p = m\dot{q}$. The Hamiltonian is

$$H = \frac{1}{2m}p^2 + V(q). \quad (5.45)$$

Let us look at this from the point of view of maximum and minimum. The second differential is determined by the Schrödinger operator

$$L = -m \frac{d^2}{dt^2} - V''(q(t)). \quad (5.46)$$

Here $q(t)$ is a function of t satisfying the Euler-Lagrange equation and the boundary conditions. So for instance if this operator with Dirichlet boundary conditions at t_1 and t_2 has strictly positive eigenvalues, then the solution will be a minimum. This will happen, for instance, if $V''(q) \leq 0$. In many other cases, the solution will not be a minimum of the action, but only a stationary point.

Example: Take an example with $V''(q) < 0$. Then the problem is to find $q(t)$ that minimizes the integral from t_1 to t_1 of $(1/2)m\dot{q}^2 - V(q)$. Here $q(t_1) = q_1$ and $q(t_2) = q_2$ are fixed. Clearly there is tradeoff. One would like the solution to linger as long as possible in the region where $V(q)$ is large. On the other hand, to satisfy the boundary conditions the solution should not move too fast. The solution will start at the point y_1 at time t_1 . Then it go reasonably, but not excessively, rapidly to the region where $V(q)$ is large. There it will linger. Then it will fall back to the point y_2 , arriving there at time t_2 . This minimization problem seems to have nothing to do with Newton's laws. But it gives the same answer.

5.8 Problems

Let L be a function of q and \dot{q} given by

$$L = \frac{1}{2}m\dot{q}^2 - V(q).$$

Let

$$S(q) = \int_{t_1}^{t_2} L dt,$$

with functions q of t satisfying $q = q_1$ at $t = t_1$ and $q = q_2$ at $t = t_2$.

1. Show that

$$dS(q)h = \left(-m \frac{d}{dt} \dot{q} - V'(q), h\right),$$

where h satisfies Dirichlet boundary conditions at t_1 and t_2 .

2. Consider a $q(t)$ for which $dS(q) = 0$. Show that

$$d^2S(q)(h, h) = \int_{t_1}^{t_2} \left[m \left(\frac{dh}{dt} \right)^2 - V''(q)h^2 \right] dt.$$

where the functions h satisfy Dirichlet boundary conditions at t_1 and t_2 .

3. Consider a $q(t)$ for which $dS(q) = 0$. Show that

$$d^2S(q)(h, h) = \left(h, \left[-m \frac{d^2}{dt^2} - V''(q) \right] h \right),$$

where the operator satisfies Dirichlet boundary conditions at t_1 and t_2 .

4. Show that if $V(q)$ is concave down, then the solution $q(t)$ of the variational problem is actually a minimum.
5. Let $H = m\dot{q}^2 - L = (1/2)m\dot{q}^2 + V(q)$. Show that $H = E$ along a solution, where E is a constant.
6. From now on take the example $V(q) = -(1/2)kq^2$. Here $k > 0$. Note the sign. We are interested in solutions with $E > 0$. Let $\omega = \sqrt{k/m}$. Show that $q = C \sinh(\omega t)$ is a solution, and find the constant C in terms of E .
7. Take $t_1 = -T$ and $t_2 = T$. Take $q_1 = -a$ and $q_2 = a$. Fix a . Write the boundary condition $a = C \sinh(\omega T)$ as a relation between T and E . Show that $T \rightarrow 0$ implies $E \rightarrow \infty$, while $T \rightarrow \infty$ implies $E \rightarrow 0$.
8. Interpret the result of the last problem intuitively in terms of particle motion satisfying conservation of energy.
9. Interpret the result of the same problem intuitively in terms of a minimization problem.

5.9 The path integral

Let us return to the problem of evaluating the quantum mechanical time evolution $\exp(-itH/\hbar)$. Here the total energy operator $H = H_0 + V$ is the sum of two non-commuting operators, corresponding to potential energy and kinetic energy.

The problem is easy when we just have kinetic energy. Then we must evaluate $\exp(-itH_0/\hbar)$, where $H_0 = p^2/(2m) = -\hbar^2/(2m)d^2/dx^2$. In the Fourier transform representation this is multiplication by $\hbar^2/(2m)k^2$. So in the Fourier transform representation the unitary time evolution is multiplication by $\exp(-it(\hbar/m)k^2/2)$. This is like the heat equation with complex diffusion coefficient $i\sigma^2 = i(\hbar/m)$. So the solution is convolution by the inverse Fourier transform, which is $1/\sqrt{2\pi i\sigma^2 t} \exp(-x^2/(2i\sigma^2 t)) = 1/\sqrt{2\pi i(\hbar/m)t} \exp(ix^2/(2(\hbar/m)t)$. In other words,

$$(\exp(-\frac{itH_0}{\hbar})u)(x) = \int_{-\infty}^{\infty} \exp(\frac{im(x-x')^2}{t\hbar})u(x') \frac{dx'}{\sqrt{2\pi i(\hbar/m)t}}. \quad (5.47)$$

The calculation of the exponential is also easy when we just have potential energy. In fact,

$$(\exp(-\frac{itV}{\hbar})u)(x) = \exp(-\frac{itv(x)}{\hbar})u(x). \quad (5.48)$$

The Trotter product formula is a remarkable formula that expresses the result for the sum $H = H_0 + V$ in terms of the separate results for H_0 and for V . It may be proved under various circumstances, for example when V is a bounded operator. The formula says that

$$\exp(-\frac{itH}{\hbar})u = \lim_{n \rightarrow \infty} \left(\exp(-\frac{i(t/n)H_0}{\hbar}) \exp(-\frac{i(t/n)V}{\hbar}) \right)^n u. \quad (5.49)$$

We can write out the Trotter product formula in detail using the results obtained before for H_0 and for V separately. Write $\Delta t = t/n$. Then the quantity of interest is $(\exp(-\frac{itH}{\hbar})u)(x) = u(x, t)$ given by the Trotter formula. This works out to be

$$u(x, t) = \lim_{n \rightarrow \infty} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \exp \left(\frac{i \sum_{j=0}^{n-1} \left[\left(\frac{x_{j+1} - x_j}{\Delta t} \right)^2 - V(x_j) \right] \Delta t}{\hbar} \right) u(x_0) \frac{dx_{n-1} \cdots dx_0}{(\sqrt{2\pi i(\hbar/m)\Delta t})^n}, \quad (5.50)$$

where $x_n = x$.

So far this has been rigorous. However now we follow Feynman and take the formal limit as $n \rightarrow \infty$ and $\Delta t \rightarrow 0$ with $n\Delta t = t$ fixed. This gives

$$(\exp(-\frac{itH}{\hbar})u)(x) = \int \exp \left(\frac{i \int_0^t \left[\left(\frac{dx(t')}{dt} \right)^2 - V(x(t')) \right] dt'}{\hbar} \right) u(x_0) \mathcal{D}x, \quad (5.51)$$

where the integral is over all paths x from some x_0 at time 0 to fixed final x at time t . This can also be written

$$\left(\exp\left(-\frac{itH}{\hbar}\right)u\right)(x) = \int \exp\left(\frac{i \int_0^t L(x(t'), \frac{dx(t')}{dt}) dt'}{\hbar}\right) u(x_0) \mathcal{D}x, \quad (5.52)$$

where

$$L(x(t'), \frac{dx(t')}{dt}) = \left(\frac{dx(t')}{dt}\right)^2 - V(x(t')) \quad (5.53)$$

is the Lagrangian. Thus the integrand in the path integral is the exponential of i times the action divided by \hbar .

This expression goes some way toward an explanation of the principle of stationary action. Consider a part of the integral near a path that is not a critical point of the action. Then nearby paths will have considerably different values of the action, and after division by \hbar the phase will be very different. So there will be a lot of cancelation. On the other hand, the part of the integral near a path that is a critical point will contribute more or less the same value of the action. So there will be no cancelation.

5.10 Appendix: Lagrange multipliers

In this section y will represent the coordinates y_1, \dots, y_n . We are interested in finding the critical points of a scalar function $F(y) = F(y_1, \dots, y_n)$.

Consider first the problem of finding a critical point of $F(y)$ with no constraints. The usual method is to compute the differential and set it equal to zero, so that the equation to be solved is $dF(y) = 0$.

Consider the problem of finding a critical point of $F(y)$ subject to a constraint $G(y) = \kappa$.

The method of Lagrange multipliers is to say that the differential of $F(y)$ without the constraint must satisfy

$$dF(y) = \lambda dG(y) \quad (5.54)$$

for some λ . That is, the function can vary only by relaxing the constraint. The change in the function must be proportional to the change in the constraint. This fundamental equation is an equation for differential forms, and it takes the same form in every coordinate system.

Example: Maximize $y_1 y_2$ subject to $y_1^2 + y_2^2 = \kappa$. The equation is $y_2 dy_1 + y_1 dy_2 = \lambda(2y_1 dy_1 + 2y_2 dy_2)$. This gives $y_2 = 2\lambda y_1$ and $y_1 = 2\lambda y_2$. Eliminating λ we get $y_1^2 = y_2^2$. Combine this with $y_1^2 + y_2^2 = \kappa$. We get $y_1 = \pm\sqrt{\kappa/2}$ and $y_2 = \pm\sqrt{\kappa/2}$. The value of $y_1 y_2$ at these points are $\pm\kappa/2$. If you care, you can also compute that $\lambda = \pm 1/2$.

Say that \bar{y} is a critical point, so in particular it satisfies the constraint $G(\bar{y}) = \kappa$. Then $dG(\bar{y}) = d\kappa$. It follows that $dF(\bar{y}) = \lambda dG(\bar{y}) = \lambda d\kappa$. The conclusion is that the Lagrange multiplier λ is the rate of change of the value of the function at the critical point as a function of the constraint parameter, that is, $\lambda = \frac{dF(\bar{y})}{d\kappa}$.

Example: In the last example the derivative of the maximum value $\pm\kappa/2$ with respect to κ is $\lambda = \pm 1/2$.

In calculations, one wants to maximize $F(y)$ subject to the constraint $G(y) = \kappa$. The idea is to take λ arbitrary. Set $\bar{F}(y) = F(y) - \lambda G(y)$ and require $d\bar{F}(y) = 0$. This gives the above equation for y and λ . However one also has the constraint equation for y and λ . These two are then solved simultaneously.

The method of Lagrange multipliers extends to the case when there are several constraints.

Consider the problem of finding a critical point of $F(y)$ subject to constraint $G_1(y) = \kappa_1, \dots, G_m(y) = \kappa_m$.

The method of Lagrange multipliers is to say that the differential of $F(y)$ without the constraint must satisfy

$$dF(y) = \sum_{j=1}^m \lambda_j dG_j(y) \quad (5.55)$$

for some λ . That is, the function can vary only by relaxing the constraint. The change in the function must be proportional to the change in the constraints. This fundamental equation is an equation for differential forms, and it takes the same form in every coordinate system.

Say that \bar{y} is a critical point, so in particular it satisfies the constraints $G_j(\bar{y}) = \kappa_j$. Then $dG_j(\bar{y}) = d\kappa_j$. It follows that $dF(\bar{y}) = \sum_j \lambda_j dG_j(\bar{y}) = \sum_j \lambda_j d\kappa_j$. The conclusion is that the Lagrange multiplier λ_j is the rate of change of the value of the function at the critical point as a function of the constraint parameter κ_j , with the other constraint parameters fixed. Thus $\lambda_j = \frac{\partial F(\bar{y})}{\partial \kappa_j}$.

Chapter 6

Perturbation theory

6.1 The implicit function theorem: scalar case

Let $f(u, \epsilon)$ be a function of u and ϵ . Let $f_u(u, \epsilon)$ be the partial derivative of F as a function of u . The assumption for the implicit function theorem is that $f(u_0, 0) = 0$ and the partial derivative $f_u(u_0, 0) \neq 0$. There are some other technical assumptions. The conclusion is that the equation

$$f(u, \epsilon) = 0 \tag{6.1}$$

defines u as a function of ϵ for ϵ sufficiently near 0, in such a way that $\epsilon = 0$ is mapped into u_0 .

The intuition behind the theorem is that one can expand

$$f(u, \epsilon) \approx f_u(u_0, 0)(u - u_0) + f_\epsilon(u_0, 0)\epsilon. \tag{6.2}$$

Then we can solve

$$u \approx u_0 - f_u(u_0, 0)^{-1} f_\epsilon(u_0, 0)\epsilon. \tag{6.3}$$

For ϵ small this is a very good approximation. In fact, this says that at the point $\epsilon = 0$ we have

$$\frac{du}{d\epsilon} = -f_u(u_0, 0)^{-1} f_\epsilon(u_0, 0). \tag{6.4}$$

This result is called first order perturbation theory.

For practical calculations one wants to compute higher derivatives $u^{(n)}$ as a function of ϵ . These are used in the Taylor expansion

$$u = \sum_{n=0}^{\infty} \frac{u^{(n)}}{n!} \epsilon^n. \tag{6.5}$$

In first order we have

$$f_u u^{(1)} + f_\epsilon = 0 \tag{6.6}$$

which gives the first order result

$$u^{(1)} = -f_u^{-1}f_\epsilon. \quad (6.7)$$

Usually we are interested in evaluating this expression at $u = u_0$ and $\epsilon = 0$.

The second order result is obtained by differentiating the first order equation. This gives

$$f_u u^{(2)} + f_{uu}(u^{(1)})^2 + 2f_{u\epsilon}u^{(1)} + f_{\epsilon\epsilon} = 0. \quad (6.8)$$

Thus the result of second order perturbation theory is that

$$u^{(2)} = -f_u^{-1}[f_{uu}(u^{(1)})^2 + 2f_{u\epsilon}u^{(1)} + f_{\epsilon\epsilon}]. \quad (6.9)$$

Again we are interested in evaluating this expression at $u = u_0$ and $\epsilon = 0$.

The general pattern is this. The n th derivative is given by

$$f_u u^{(n)} + r_n = 0 \quad (6.10)$$

Differentiate again. This gives

$$f_u u^{(n+1)} + [f_{uu}u^{(1)} + f_{u\epsilon}]u^{(n)} + [r_{nu}u^{(1)} + r_{n\epsilon}] = 0. \quad (6.11)$$

Thus at every stage one only has to invert the first derivative f_u .

Example: Here is a simple example. Fix a . We want to solve $u = a + \epsilon g(u)$ to find u as a function of ϵ . The first derivative is $u_\epsilon = g(u) + \epsilon g'(u)u_\epsilon$. The second derivative is $u_{\epsilon\epsilon} = 2g'(u)u_\epsilon + \epsilon g''(u)u_\epsilon^2 + \epsilon g''(u)u_\epsilon^2$. The third derivative is $u_{\epsilon\epsilon\epsilon} = 3g''(u)u_\epsilon^2 + 3g''(u)u_{\epsilon\epsilon} + \epsilon$ times other junk. Evaluate these at $\epsilon = 0$, that is, at $u = a$. This gives $u_\epsilon = g(a)$, $u_{\epsilon\epsilon} = 2g(a)g'(a)$, and $u_{\epsilon\epsilon\epsilon} = 3g(a)^2g''(a) + 6g(a)g'(a)^2$. It is difficult to see the pattern. However in the next section we will see that there is one, at least in this case.

Example: Another popular way of solving the problem is to substitute $u = a + u'\epsilon + \frac{1}{2}u''\epsilon^2 + \frac{1}{6}u'''\epsilon^3 + \dots$ in the equation. This gives

$$a + u'\epsilon + \frac{1}{2}u''\epsilon^2 + \frac{1}{6}u'''\epsilon^3 + \dots = a + \epsilon[g(a) + g'(a)(u'\epsilon + \frac{1}{2}u''\epsilon^2 + \dots) + \frac{1}{2}\epsilon^2 g''(a)(u'\epsilon + \dots)^2 + \dots]. \quad (6.12)$$

Equating powers of ϵ gives the same equations, and these are solved recursively the same way.

6.2 Problems

There is at least one case when one can evaluate Taylor coefficients to all orders. Say that x is a fixed parameter and one wants to solve

$$u = x + tg(u) \quad (6.13)$$

for u as a function of t . When $t = 0$ the solution is $u = x$. The Lagrange expansion says that for arbitrary $n \geq 1$ we have

$$\frac{d^n u}{dt^n} \Big|_{t=0} = \frac{d^{n-1}}{dx^{n-1}} g(x)^n. \quad (6.14)$$

So the expansion to all orders is

$$u = x + tg(x) + \frac{t^2}{2} \frac{d}{dx} g(x)^2 + \frac{t^3}{6} \frac{d^2}{dx^2} g(x)^3 + \cdots + \frac{t^n}{n!} \frac{d^{n-1}}{dx^{n-1}} g(x)^n + \cdots \quad (6.15)$$

1. Let $w = g(u)$. Show that the equation becomes

$$w = g(x + wt). \quad (6.16)$$

2. Show that the function w is the solution of the partial differential equation

$$\frac{\partial w}{\partial t} = w \frac{\partial w}{\partial x} \quad (6.17)$$

with initial condition $w = g(x)$ at $t = 0$.

3. Prove the conservation laws

$$\frac{\partial w^n}{\partial t} = \frac{n}{n+1} \frac{\partial w^{n+1}}{\partial x}. \quad (6.18)$$

4. Prove that for $n \geq 1$ we have

$$\frac{\partial^{n-1} w}{\partial t^{n-1}} = \frac{1}{n} \frac{\partial^{n-1}}{\partial x^{n-1}} w^n. \quad (6.19)$$

5. Prove that at $t = 0$

$$w = g(x) + \frac{t}{2} \frac{d}{dx} g(x)^2 + \frac{t^2}{6} \frac{d^2}{dx^2} g(x)^3 + \cdots + \frac{t^{n-1}}{n!} \frac{d^{n-1}}{dx^{n-1}} g(x)^n + \cdots \quad (6.20)$$

6. Use $u = x + tw$ to prove the Lagrange expansion.

Note: The partial differential equation has a physical interpretation. If $v = -w$, then the equation is

$$\frac{\partial v}{\partial t} + v \frac{\partial v}{\partial x} = 0 \quad (6.21)$$

with initial condition $v = h(x)$ at time zero. This is the equation for the velocity of a gas of particles moving freely in one dimension. The motion of a particle moving with the gas is given by

$$\frac{dx}{dt} = v. \quad (6.22)$$

The acceleration of a particle is

$$\frac{dv}{dt} = \frac{\partial v}{\partial t} + \frac{dx}{dt} \frac{\partial v}{\partial x} = 0 \quad (6.23)$$

That is why it is considered free particle motion. Since v is constant along particle paths $x = x_0 + vt$, we have that v at arbitrary x, t is given in terms of the initial $v = h(x_0)$ at time zero by $v = h(x_0) = h(x - vt)$. So the solution of the equation is

$$v = h(x - vt). \quad (6.24)$$

6.3 The implicit function theorem: systems

We can take $F(u, \epsilon) = 0$ to be a system of n equations for n variables u and an extra variable ϵ . Or more generally we can take $F(u, \epsilon)$ to be a function of u and ϵ , where u ranges over a region U in some Banach space E . Then $F(u, \epsilon)$ takes values in a Banach space E' . The parameter ϵ is still real.

We shall often abbreviate the function by F . Let F_u be the partial derivative of F as a function of u . In the case of a system this is the n by n matrix of partial derivatives of the components of F with respect to the components of u . More generally, it is a function on U with values that are linear transformations from E to E' .

The assumption for the implicit function theorem is that $F(u_0, 0) = 0$ and the partial derivative $F_u(u_0, 0)$ has a bounded inverse. There are some other technical assumptions. The conclusion is that the equation

$$F(u, \epsilon) = 0 \tag{6.25}$$

defines u as a function of ϵ for ϵ sufficiently near 0, in such a way that $\epsilon = 0$ is mapped into u_0 .

For practical calculations one wants to compute higher derivatives $u^{(n)}$ as a function of ϵ . These are used in the Taylor expansion

$$u = \sum_{n=0}^{\infty} \frac{u^{(n)}}{n!} \epsilon^n. \tag{6.26}$$

In first order we have

$$F_u u^{(1)} + F_\epsilon = 0 \tag{6.27}$$

which gives the first order result

$$u^{(1)} = -F_u^{-1} F_\epsilon. \tag{6.28}$$

Note that F_ϵ has values in E' . On the other hand, F_u^{-1} has values that are linear transformation from E' to E . So the right hand side has values in E , as it must. Usually we are interested in evaluating this expression at $u = u_0$ and $\epsilon = 0$.

The second order result is obtained by differentiating the first order equation. This gives

$$F_u u^{(2)} + F_{uu}(u^{(1)}, u^{(1)}) + 2F_{u\epsilon} u^{(1)} + F_{\epsilon\epsilon} = 0. \tag{6.29}$$

Thus the result of second order perturbation theory is that

$$u^{(2)} = -F_u^{-1} [F_{uu}(u^{(1)}, u^{(1)}) + 2F_{u\epsilon} u^{(1)} + F_{\epsilon\epsilon}]. \tag{6.30}$$

Note that F_{uu} has values that are bilinear functions from $E \times E$ to E' . On the other hand, $F_{u\epsilon}$ has values that are linear functions from E to E' . Again the right hand side has values in E . Again we are interested in evaluating this expression at $u = u_0$ and $\epsilon = 0$.

The general pattern is this. The n th derivative is given by

$$F_u u^{(n)} + R_n = 0 \quad (6.31)$$

Differentiate again. This gives

$$F_u u^{(n+1)} + [F_{uu} u^{(1)} + F_{u\epsilon}] u^{(n)} + [R_{nu} u^{(1)} + R_{n\epsilon}] = 0. \quad (6.32)$$

Thus at every stage one only has to invert the first derivative F_u .

6.4 Nonlinear differential equations

Let us look at the simplest non-linear differential equation

$$Lu = \epsilon g(u). \quad (6.33)$$

Here L is the restriction of a linear operator to satisfy an inhomogeneous boundary condition. For example, we could have $L = \partial/\partial t + 1$ acting on functions with $u(0) = 1$. When $\epsilon = 0$ there is a solution $u = u_0$. In the example $u_0(t) = e^{-t}$.

The derivative of L is L_0 , where L_0 is a linear operator with zero boundary conditions. For instance, we could have $L_0 = \partial/\partial t + 1$ acting on functions h with $h(0) = 0$. Thus the first order perturbation equation is $L_0 u_\epsilon = g(u) + \epsilon g'(u) u_\epsilon$. The second order equation is $L_0 u_{\epsilon\epsilon} = 2g'(u) u_\epsilon + \text{terms with } \epsilon$. One can continue in this way. Thus the hierarchy of equations begins $Lu_0 = 0$, $L_0 u_\epsilon = g(u_0)$, $L_0 u_{\epsilon\epsilon} = 2g'(u_0) u_\epsilon$.

It is possible to think of this in another way. Write $u = u_0 + w$. Then the equation $Lu = \epsilon g(u)$ becomes $L_0 w = \epsilon g(u)$, where L_0 is the operator defined with zero boundary conditions. Suppose that L_0 has an inverse. Then

$$u = u_0 + \epsilon L_0^{-1} g(u). \quad (6.34)$$

This is very close in form to the scalar equation examined in the first two sections.

This equation may be solved by the usual procedure. The first derivative is $u_\epsilon = L_0^{-1} g(u) + \epsilon L_0^{-1} g'(u) u_\epsilon$. The second derivative is $u_{\epsilon\epsilon} = 2L_0^{-1} g'(u) u_\epsilon + \epsilon L_0^{-1} g''(u) u_\epsilon^2$. The third derivative is $u_{\epsilon\epsilon\epsilon} = 3L_0^{-1} [g''(u) u_\epsilon^2] + 3L_0^{-1} [g'(u) u_{\epsilon\epsilon}] + \epsilon$ times other junk. Evaluate these at $\epsilon = 0$, that is, at $u = u_0$. This gives

$$u_\epsilon = L_0^{-1} g(u_0), \quad (6.35)$$

and

$$u_{\epsilon\epsilon} = 2L_0^{-1} [g'(u_0) L_0^{-1} g(u_0)], \quad (6.36)$$

and

$$u_{\epsilon\epsilon\epsilon} = 3L_0^{-1} [g''(u_0) (L_0^{-1} g(u_0))^2] + 6L_0^{-1} [g'(u_0) L_0^{-1} (g'(u_0) L_0^{-1} g(u_0))]. \quad (6.37)$$

Notice how the linear operators occur in various places, making it impossible to do the simplification that we found in the scalar case.

6.5 A singular perturbation example

In quantum mechanics one often wants to solve a problem of the form

$$-\frac{\hbar^2}{2m}u'' + v(x)u = \lambda u. \quad (6.38)$$

Here $v(x)$ is a given real function. In many contexts λ plays the role of an eigenvalue parameter. The constant $m > 0$ is fixed. The constant $\hbar > 0$ is the perturbation parameter.

The characteristic feature of this problem is that the zero order perturbation is not apparent. Thus setting $\hbar = 0$ it is not clear that there will be a solution at all.

The resolution of this problem is to write the equation in new variables. For simplicity think of the case when $\lambda > v(x)$. Thus

$$u = Ae^{i\frac{S}{\hbar}}. \quad (6.39)$$

The equation becomes

$$A \left[\frac{1}{2m}(S')^2 + v(x) - \lambda \right] = \frac{\hbar^2}{2m}A'' \quad (6.40)$$

together with

$$2A'S' + AS'' = 0. \quad (6.41)$$

Now we can recognize the $\hbar = 0$ limit as

$$\frac{1}{2m}(S')^2 + v(x) = \lambda. \quad (6.42)$$

The other equation may be written

$$(A^2S')' = 0. \quad (6.43)$$

These equations have a physical interpretation. The function S' is a velocity that depends on position. The first equation is the classical equation for conservation of energy. The function A^2 is a density. The second equation is the conservation law that says that the current A^2S' is constant in space. Of course these equations are only valid in the zero order approximation, which in this case is called the WKB approximation.

This approximation is relevant to a heuristic understanding of Schrödinger operators of the form

$$H = -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + v(x). \quad (6.44)$$

The Hilbert space is the space of L^2 functions on the line. If $v(x)$ is bounded below, then one expects this to be a well-defined self-adjoint operator. In fact, for $\lambda < 0$ sufficiently negative, one expects $(H - \lambda I)^{-1}$ to be a well-defined bounded self-adjoint operator. The reason is that the solutions of the differential

equation either decay rapidly at infinity or grow rapidly at infinity. The Green's function is defined by the solutions that decay rapidly at infinity. There is a solution of $Hu = \lambda u$ that is in L^2 at $-\infty$ and by another solution of the same equation that is in L^2 at $+\infty$. The two solutions are matched by a jump condition to form the kernel of the integral operator.

When $v(x)$ is very badly unbounded below, then something quite different happens. Let us look at the example of $v(x) \sim \epsilon x^{2n}$ with $\epsilon < 0$. According to the WKB approximation, we have $S'(x) \sim \pm|x|^n$. It follows that $A^2 \sim |x|^{-n}$. Thus it is plausible that if $n > 1$ then every solution is in L^2 . These solutions are highly oscillatory and decrease at infinity at a moderate rate.

If this is indeed the case, then the operator H is not self-adjoint. The problem is that when one wants to define a Green's function, there are two linearly independent solutions, and they are both in L^2 at $\pm\infty$. So there is no boundary condition at infinity that determines the Green's function in a natural way.

6.6 Eigenvalues and eigenvectors

We have an operator $H_0 + \epsilon V$. We are interested in an isolated eigenvalue λ of multiplicity 1. The corresponding eigenvector is p . It is only determined up to a constant multiple. The adjoint operator has eigenvalue $\bar{\lambda}$ with eigenvector q . It is convenient to choose p and q so that the inner product $(q, p) = 1$. The associated projection onto the eigenspace is P satisfying $P^2 = P$. It is given by $Pu = (q, u)p$. Thus p is an eigenvector with

$$(H_0 + \epsilon V - \lambda)p = 0. \quad (6.45)$$

Furthermore, $(q, (H_0 + \epsilon V - \lambda)u) = 0$ for all u . We also have the reduced resolvent S with $SP = PS = 0$ and $S(H_0 + \epsilon V - \lambda I) = 1 - P$.

If we indicate the dependence on ϵ explicitly, then the equation we want to solve is

$$(H_0 + \epsilon V - \lambda(\epsilon))p(\epsilon) = 0. \quad (6.46)$$

We would like to find the perturbation expansion in terms of the λ and p and q and S associated with $\epsilon = 0$.

The following discussion is meant to establish two formulas, one for the eigenvalue and one for the eigenvector (not normalized). The first is the formula for the eigenvalue to second order

$$\lambda(\epsilon) = \lambda + (q, Vp)\epsilon - (q, VSVp)\epsilon^2 + \dots \quad (6.47)$$

The other is the formula for the (unnormalized) eigenvector to second order

$$p(\epsilon) = p - SVp\epsilon + [SVSVp - S^2Vp(q, Vp)]\epsilon^2 + \dots \quad (6.48)$$

These formulas are so important that they should be memorized, at least the one for the eigenvalues.

We now proceed to the derivations. Differentiate the eigenvector equation with respect to ϵ . This gives

$$(H_0 + \epsilon V - \lambda)\dot{p} - \dot{\lambda}p + Vp = 0. \quad (6.49)$$

Now take the inner product with q on the left. This gives

$$\dot{\lambda} = (q, Vp). \quad (6.50)$$

This gives the first order change of the eigenvalue. That is,

$$\lambda(\epsilon) = \lambda + (q, Vp)\epsilon + \dots \quad (6.51)$$

Next apply $I - P$ on the left. This gives

$$(I - P)(H_0 + \epsilon V - \lambda)\dot{p} + (I - P)Vp = 0, \quad (6.52)$$

or

$$(I - P)\dot{p} = -SVp. \quad (6.53)$$

Thus

$$\dot{p} = -SVp + c_1p. \quad (6.54)$$

Notice that the derivative of the eigenvector is not uniquely determined. It is ordinarily harmless to take $c_1 = 0$. This gives the first order change of the eigenvector

$$p(\epsilon) = p + [-SVp + c_1p]\epsilon. \quad (6.55)$$

Now differentiate again. This gives

$$(H_0 + \epsilon V - \lambda)\ddot{p} + \ddot{\lambda}p + 2V\dot{p} - 2\dot{\lambda}\dot{p} = 0. \quad (6.56)$$

Take the inner product with q on the left. This gives

$$\ddot{\lambda} + 2(q, V\dot{p}) - 2\dot{\lambda}(q, \dot{p}), \quad (6.57)$$

or

$$\frac{1}{2}\ddot{\lambda} = -(q, VSVp). \quad (6.58)$$

This gives the second order change of the eigenvalue. That is,

$$\lambda(\epsilon) = \lambda + (q, Vp)\epsilon - (q, VSVp)\epsilon^2 + \dots \quad (6.59)$$

Now apply $1 - P$ on the left. This gives

$$(1 - P)(H_0 + \epsilon V - \lambda)\ddot{p} + 2(1 - P)V\dot{p} - 2\dot{\lambda}(1 - P)\dot{p} = 0. \quad (6.60)$$

Thus

$$\frac{1}{2}(1 - P)\ddot{p} = SVSVp - \frac{1}{2}c_1SVp - S^2Vp(q, Vp). \quad (6.61)$$

Hence

$$\frac{1}{2}\ddot{p} = SVSVp - S^2Vp(q, Vp) - \frac{1}{2}c_1SVp + \frac{1}{2}c_2p. \quad (6.62)$$

This gives the second order change of the eigenvector

$$p(\epsilon) = p + [-SVp + c_1p]\epsilon + [SVSVp - S^2Vp(q, Vp) - \frac{1}{2}c_1SVp + \frac{1}{2}c_2p]\epsilon^2 + \dots \quad (6.63)$$

Now let us change notation. We will think of a basis p_n for the Hilbert space and a dual basis q_m , with $(q_m, p_n) = \delta_{mn}$. We write q_m^* for the functional that takes u to (q_m, u) . Then the various operators have special forms. We take the p_n to be the eigenvectors of H_0 . Thus $H_0p_n = \lambda_n p_n$. The operator $P_n = p_n q_n^*$ is the spectral projection associated with the n th eigenvalue λ_n of H_0 . The operator $S_n = \sum_{j \neq n} p_j q_j^* / (\lambda_k - \lambda_n)$.

Then the second order expression for the eigenvalue can be written as

$$\lambda_n(\epsilon) = \lambda_n + (q_n, Vp_n)\epsilon - (q_n, VS_nVp_n)\epsilon^2 + \dots \quad (6.64)$$

Even more explicitly, this is

$$\lambda_n(\epsilon) = \lambda_n + (q_n, Vp_n)\epsilon - \sum_{j \neq n} (q_n, Vp_j) \frac{1}{\lambda_j - \lambda_n} (q_j, Vp_n)\epsilon^2 + \dots \quad (6.65)$$

The second order expression for the eigenvector $p_n(\epsilon)$ is (taking $c_1 = 0$)

$$p_n(\epsilon) = p_n - S_nVp_n\epsilon + [S_nVS_nVp_n - S_n^2Vp_n(q_n, Vp_n) + \frac{1}{2}c_2p_n]\epsilon^2 + \dots \quad (6.66)$$

If H_0 has discrete spectrum, then (leaving out the last term) this may be written

$$\begin{aligned} p_n(\epsilon) &= p_n - \sum_{j \neq n} \frac{1}{\lambda_j - \lambda_n} (q_j, Vp_n) p_j \epsilon \\ &+ \sum_{j \neq n} \frac{1}{\lambda_j - \lambda_n} \left[\sum_{k \neq n} (q_j, Vp_k) \frac{1}{\lambda_k - \lambda_n} (q_k, Vp_n) - \frac{1}{\lambda_j - \lambda_n} (q_j, Vp_n) (q_n, Vp_n) \right] p_j \epsilon^2 + \dots \end{aligned} \quad (6.67)$$

6.7 The self-adjoint case

In the self-adjoint case $q = p$ and it is natural to normalize so that $(p(\epsilon), p(\epsilon)) = 1$ to second order. This gives $c_1 = 0$ and

$$c_2 = -(p, VS^2Vp). \quad (6.68)$$

So in this case

$$\lambda(\epsilon) = \lambda + (p, Vp)\epsilon - (p, VSVp)\epsilon^2 + \dots \quad (6.69)$$

and

$$p(\epsilon) = p - SVp\epsilon + [SVSVp - S^2Vp(p, Vp) - \frac{1}{2}(p, VS^2Vp)p]\epsilon^2 + \dots \quad (6.70)$$

If $\lambda = \lambda_n$ and $p = p_n$, we can write the coefficient c_2 even more explicitly as

$$c_2 = - \sum_{j \neq n} (p_n, V p_j) \frac{1}{(\lambda_j - \lambda_n)^2} (p_j, V p_n). \quad (6.71)$$

Example: Take

$$H_0 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad (6.72)$$

and

$$V = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}. \quad (6.73)$$

The eigenvalues are $\lambda(\epsilon) = \pm\sqrt{1 + \epsilon^2}$. However it is interesting to compare this exact result with the result of second order perturbation theory.

Let us look at the perturbation of the eigenvalue -1 of H_0 . For this eigenvalue the spectral projection is

$$P = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \quad (6.74)$$

and the eigenvector p is the second column of P . The reduced resolvent is

$$S = \begin{bmatrix} \frac{1}{1 - (-1)} & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & 0 \end{bmatrix}. \quad (6.75)$$

So the coefficient in second order perturbation theory is

$$-(p, V S V p) = -\frac{1}{2}. \quad (6.76)$$

Thus

$$\lambda(\epsilon) = -1 - \frac{1}{2}\epsilon^2 + \dots \quad (6.77)$$

6.8 The anharmonic oscillator

In this section we let

$$H_0 = \frac{1}{2} \left[\frac{d^2}{dx^2} + x^2 - 1 \right]. \quad (6.78)$$

The perturbation V is multiplication by

$$v(x) = \epsilon x^4. \quad (6.79)$$

Let

$$A = \frac{1}{\sqrt{2}} \left(x + \frac{d}{dx} \right) \quad (6.80)$$

and

$$A^* = \frac{1}{\sqrt{2}} \left(x - \frac{d}{dx} \right). \quad (6.81)$$

Then $AA^* = A^*A + I$ and $H_0 = A^*A$. Furthermore, the solution of $Au_0 = 0$ is

$$u_0(x) = \frac{1}{\sqrt[4]{\pi}} e^{-\frac{x^2}{2}}. \quad (6.82)$$

Define for each $n = 0, 1, 2, 3, \dots$ the vector

$$u_n = \frac{1}{n!} A^{*n} u_0. \quad (6.83)$$

We have $A^*u_m = \sqrt{m+1}u_{m+1}$ and $Au_m = \sqrt{m}u_{m-1}$. Then the u_n form an orthonormal basis consisting of eigenvectors of H_0 . Furthermore,

$$H_0 u_n = n u_n. \quad (6.84)$$

Say that we are interested in a particular n . The corresponding reduced resolvent S_n is given by

$$S_n u_m = \frac{1}{m-n} u_m \quad (6.85)$$

for $m \neq n$, and by $S_n u_n = 0$.

The operator multiplication by x may be expressed in terms of A and A^* by

$$x = \frac{1}{\sqrt{2}}(A + A^*). \quad (6.86)$$

This allows us to calculate the matrix elements of x . These are given by

$$(u_{m+1}, x u_m) = \frac{1}{\sqrt{2}}(u_{m+1}, A^* u_m) = \frac{1}{\sqrt{2}} \sqrt{m+1} \quad (6.87)$$

and

$$(u_{m-1}, x u_m) = \frac{1}{\sqrt{2}}(u_{m-1}, A u_m) = \frac{1}{\sqrt{2}} \sqrt{m}. \quad (6.88)$$

So each transition is to a neighboring natural number.

As a warmup, let us compute the matrix element $(u_n, x^2 u_n)$. This is the sum of two possible transition paths, one up-down and one down-up. The result is $(n+1)/2 + (n/2) = (2n+1)/2$.

To compute the first order perturbation coefficient $(u_n, x^4 u_n)$, we need to look at all transitions that go from n to n in 4 steps. After two steps one is either at u_{n-2} , at u_n , or at u_{n+2} . So

$$(u_n, x^4 u_n) = (u_n, x^2 u_{n-2})^2 + (u_n, x^2 u_n)^2 + (u_n, x^2 u_{n+2})^2. \quad (6.89)$$

This is

$$(u_n, x^4 u_n) = \frac{n(n-1)}{4} + \left(\frac{2n+1}{2}\right)^2 + \frac{(n+1)(n+2)}{4} = \frac{6n^2 + 6n + 3}{4}. \quad (6.90)$$

Higher order perturbations may be computed with more labor. It turns out that the perturbation series in ϵ does not converge. Some intuition for this is provided by the observation that $H_0 + \epsilon V$ is not a self-adjoint operator for $\epsilon < 0$. So for $\epsilon < 0$ the eigenvalue problem is meaningless. On the other hand, if the Taylor series had a positive radius of convergence, then it would converge also for some values of $\epsilon < 0$.