

# The Central Limit Theorem

November 19, 2009

Convergence in distribution  $X_n \rightarrow^{\mathcal{D}} X$  is defined to be

$$\lim_{n \rightarrow \infty} Eh(X_n) = Eh(X).$$

or every bounded continuous function  $h : \mathbb{R} \rightarrow \mathbb{R}$ .

However, it is not necessary to verify this for each choice of  $h$ . We can limit ourselves to a smaller so-called **convergence determining** family of functions.

- For random variables taking values in the natural numbers,  $\{h_z(x) = z^x; |z| < 1\}$  is convergence determining. In this case, we are looking at convergence of the probability generating function.
- For real valued random variables,  $\{h_t(x) = \exp tx; -h < t < h\}$  is convergence determining provided the necessary expected values exist. Note that  $\exp tx$  is not bounded and so we need to make an additional argument to include these functions. In this case, we are looking at convergence of the moment generating function.

**Example 1.** For the binomial distribution with parameters  $n$  and  $p$ , the probability generating function is

$$\rho_{X_n}(z) = ((1-p) + pz)^n = (1 - p(1-z))^n$$

If we take the success probability  $p = \lambda/n$  to depend on  $n$ , then

$$\rho_{X_n}(z) = ((1-p) + pz)^n = \left(1 - \frac{\lambda}{n}(1-z)\right)^n \rightarrow \exp \lambda(1-z) = \rho_X(z),$$

the probability generating function for a Poisson random variable  $X$  with parameter  $\lambda$ . Thus, we have that the given binomial random variables converge in distribution to a Poisson random variable.

To use this, assume that  $n$  is large, but  $\lambda = np$  is moderate the binomial random variable is well approximated by a Poisson random variable. In particular,  $Eh(X_n) \approx Eh(X)$  for any bounded continuous  $h$

## 1 Central Limit Theorem

If we look at distributions for the sum  $T_n = X_1 + X_2 + \dots + X_n$ , what do we see. Let's look first to the simplest case,  $X_i$  Bernoulli random variables.

This is looking like the bell curve. To make the comparisons fair, let's look at standardized versions of the random variables with mean  $\mu$  and variance  $\sigma^2$ ,




Figure 1: a. Successes in 100 Bernoulli trials with  $p = 0.2, 0.4, 0.6$  and  $0.8$ . b. Successes in Bernoulli trials with  $p = 1/2$  and  $n = 20, 40$  and  $80$ .

$$Z_n = \frac{T_n - n\mu}{\sigma\sqrt{n}} \quad (1)$$

and look at the density of the sum of standardized exponential random variables.

Again, we see the densities approaching that of a bell curve. The **classical central limit theorem** states that if  $\{X_i; i \geq 1\}$  are independent and identically distributed with common mean  $\mu$  and common variance  $\sigma^2$ , then  $Z_n$  as defined by equation (1) converges to  $Z$  a standard normal random variable. In terms of the cumulative distribution function

$$\lim_{n \rightarrow \infty} P\{Z_n \leq z\} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^z e^{-x^2/2} dx = \Phi(z)$$

where  $\Phi$  is the cumulative distribution function of the standard normal.

We will prove this in the case that the  $X_i$  have a moment generating function  $M_X(t)$  for the interval  $t \in (-h, h)$  by showing that

$$\lim_{n \rightarrow \infty} M_{Z_n}(t) = \exp \frac{t^2}{2}$$

or equivalently, show that the cumulant generating functions  $K_{Z_n}(t) = \log M_{Z_n}(t)$  satisfy

$$\lim_{n \rightarrow \infty} K_{Z_n}(t) = \frac{t^2}{2}$$

Write

$$Y_i = \frac{X_i - \mu}{\sigma}$$




Figure 2: Density of the standardized version of the sum of  $n$  independent exponential random variables for  $n = 2, 4, 8, 16$  and  $32$ .

then

$$Z_n = \frac{1}{\sqrt{n}} \sum_{i=1}^n Y_i.$$

For  $M_Y(t)$  the moment generating function for the  $Y_i$  and  $M_{T_n}(t)$ , the moment generating function for  $T_n$ ,

$$M_{Z_n}(t) = E \exp(tZ_n) = E \exp\left(\frac{t}{\sqrt{n}} \sum_{i=1}^n Y_i\right) = \prod_{i=1}^n E \exp\left(\frac{t}{\sqrt{n}} Y_i\right) = \left(M_Y\left(\frac{t}{\sqrt{n}}\right)\right)^n$$

and

$$K_{Z_n}(t) = n \log M_Y\left(\frac{t}{\sqrt{n}}\right) = nK_Y\left(\frac{t}{\sqrt{n}}\right).$$

Recall that for the cumulant generating function  $K_Y$ ,

$$K'_Y(0) = EY_1 = 0, \quad K''_Y(0) = \text{Var}(Y) = 1.$$

Finally, from two applications of L'Hôpital's rule,

$$\lim_{n \rightarrow \infty} K_{Z_n}(t) = \lim_{n \rightarrow \infty} nK_Y\left(\frac{t}{\sqrt{n}}\right) = \lim_{\epsilon \rightarrow 0} \frac{K_Y(\epsilon t)}{\epsilon^2} = \lim_{\epsilon \rightarrow 0} \frac{tK'_Y(\epsilon t)}{2\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{t^2 K''_Y(\epsilon t)}{2} = \frac{t^2 K''_Y(0)}{2} = \frac{t^2}{2}.$$

**Example 2.** For Bernoulli trials,  $\mu = p$  and  $\sigma^2 = p(1-p)$ . Thus, for large enough  $n$

$$Z_n = \frac{T_n - np}{\sqrt{np(1-p)}},$$

has approximately the distribution of a standard normal random variable. For 100 tosses of a fair coin,

$$Z_n = \frac{T_n - 50}{5},$$

and

$$\{T_n \leq 40\} = \{Z_n \leq -2\}$$

So,

$$P\{T_n \leq 40\} \approx P\{Z \leq -2\} = 0.054.$$

**Example 3.** For an exponential sample with mean 1. Then, the standard deviation is also 1 and for 64 observations

$$Z_n = \frac{T_n - 64}{8},$$

$$\{T_n \geq 78\} = \{Z_n \geq 1.75\}$$

So,

$$P\{T_n \geq 78\} \approx P\{Z \geq -1.75\} = 0.086.$$

## 2 Slutsky's Theorem

Some useful extensions of the central limit theorem are based on Slutsky's theorem.

**Theorem 4.** Let  $X_n \rightarrow^{\mathcal{D}} X$  and  $Y_n \rightarrow^P a$ , a constant as  $n \rightarrow \infty$ . Then

1.  $Y_n X_n \rightarrow^{\mathcal{D}} aX$ , and
2.  $X_n + Y_n \rightarrow^{\mathcal{D}} X + a$ .

For example, by the law of large numbers, the sample variance

$$S_n^2 \xrightarrow{a.s.} \sigma^2,$$

the distribution variance as  $n \rightarrow \infty$ . Thus,

$$\frac{S_n}{\sigma} \xrightarrow{a.s.} 1.$$

Thus, it also converges in probability. So, by Slutsky's theorem, the  $t$ -statistic

$$\frac{T_n - n\mu}{S_n \sqrt{n}} = \frac{S_n}{\sigma} \frac{T_n - n\mu}{\sigma \sqrt{n}} = \frac{S_n}{\sigma} Z_n \rightarrow^{\mathcal{D}} 1 \cdot Z,$$

a standard normal as  $n \rightarrow \infty$

### 3 Delta Method

For a random sample  $\{X_n \geq 1\}$  with common mean  $\mu$  and common variance  $\sigma^2$ , we can write the central limit theorem using the sample mean.

$$\sqrt{n}(\bar{X}_n - \mu) \rightarrow^{\mathcal{D}} \sigma Z$$

where  $Z$  is a standard normal.

To generalize this, assume that  $\{Y_n \geq 1\}$  is a sequence of random variables satisfying

$$\sqrt{n}(Y_n - \theta) \rightarrow^{\mathcal{D}} \sigma Z$$

for some value  $\theta$

Then the **delta method** states that if a function  $g$  has a continuous derivative and  $g'(\theta) \neq 0$ , then

$$\sqrt{n}(g(Y_n) - g(\theta)) \rightarrow^{\mathcal{D}} \sigma g'(\theta) \tilde{Z}$$

where  $\tilde{Z}$  is also a standard normal.

To prove this, expand  $g$  as a Taylor's series about the value  $\theta$

$$g(Y_n) = g(\theta) + g'(\tilde{\theta})(Y_n - \theta),$$

or

$$\sqrt{n}(g(Y_n) - g(\theta)) = g'(\tilde{\theta})\sqrt{n}(Y_n - \theta).$$

where  $\tilde{\theta}$  lies between  $Y_n$  and  $\theta$ . Note that since  $Y_n \rightarrow^P \theta$  implies  $\tilde{\theta} \rightarrow^P \theta$  and  $g'(\theta)$  is continuous,

$$g'(\tilde{\theta}) \rightarrow^P g'(\theta).$$

and the theorem follows from applying Slutsky's theorem.

**Example 5.** For Bernoulli trials, write  $\bar{X} = \hat{p}$ , then

$$\sqrt{n}(\hat{p} - p) \rightarrow^{\mathcal{D}} \sqrt{p(1-p)}Z.$$

If we could find  $g$  so that

$$g'(p) = \frac{1}{\sqrt{p(1-p)}},$$

then

$$\sqrt{n}(g(\hat{p}) - g(p)) \rightarrow^{\mathcal{D}} Z.$$

Such a choice, which here is  $g(p) = 2 \arcsin(\sqrt{p})$  is called a **variance stabilizing transformation**.