

Calculus and Differential Equations I

MATH 250 A

Differential equations of the form $y' = g(x, y)$

Separable equations

- A first order differential equation is **separable** if it is of the form

$$\frac{dy}{dx} = f(y)g(x).$$

- To solve such an equation, we write

$$\frac{1}{f(y)} dy = g(x) dx,$$

and integrate both sides.

- Of course, when we divide by $f(y)$ we may lose **equilibrium solutions**, which are then **singular solutions**.
- **Example:** Solve $x^2y' = y(y - 1)$. Then, check your answer.

Equations with homogeneous coefficients

- A differential equation of the form

$$y' = g\left(\frac{y}{x}\right)$$

is said to have **homogeneous coefficients**.

- To solve such an equation, **set $z = \frac{y}{x}$** , and find an equation for z :

$$g(z) = \frac{dy}{dx} = x \frac{dz}{dx} + z.$$

- The above equation is **separable** and can then be solved by **separation of variables**. As usual, one has to be careful to keep track of **singular solutions**.
- **Example:** Solve $y' = \frac{x^2 + xy + y^2}{xy}$. Then, check your answer.

Linear equations

- We now consider **Linear first order** differential equations of the form

$$y' + p(x)y = q(x).$$

- To solve this equation, realize that **the left-hand side can be written as**

$$y' + p(x)y = \exp\left(-\int p(x) dx\right) \frac{d}{dx} \left(y \exp\left(\int p(x) dx\right)\right).$$

- As a consequence, we have

$$y(x) = \frac{1}{K(x)} \int_a^x K(t)q(t) dt + \frac{C}{K(x)}, \quad K(x) = \exp\left(\int_a^x p(t) dt\right).$$

Linear equations (continued)

- The function $K(x)$ is called an **integrating factor**.
- **Existence and uniqueness theorem** for first-order linear equations:
If p and q are continuous functions of x on the interval (a, b) , and if $x_0 \in (a, b)$, then there is a unique solution of

$$y' + p(x)y = q(x)$$

for $x \in (a, b)$.

- **Example:** Solve the equation for an RL electrical circuit

$$L \frac{dI}{dt} + RI = E(t),$$

where I is the current intensity, $E(t)$ is the voltage, R is a resistance and L an inductance.

Bernoulli equations

- A first-order differential equation of the form

$$y' + p(x)y = q(x)y^n,$$

where n is a (positive or negative) integer with $n \neq 1$, is called a **Bernoulli equation**.

- To solve this equation, **make the change of variable**

$$u = y^{1-n},$$

to obtain a **linear equation for u** . Then, solve the equation for u and express the answer in terms of y .

- As usual when manipulating differential equations, one should be careful not to lose or introduce solutions.
- **Example:** Solve $y' = ay - y^3$, which is the equation describing a **Pitchfork bifurcation** as a goes through 0.

Existence

$$\frac{dy}{dx} = g(x, y)$$

- **Existence:** If g is a **continuous** function of x and y on the rectangle

$$\mathcal{R} = \{(x, y), |x - x_0| \leq a, |y - y_0| \leq b\}$$

where $a > 0$ and $b > 0$, then there exists a continuously differentiable solution y of the above differential equation on $|x - x_0| \leq \alpha$ for which $y(x_0) = y_0$, where

$$\alpha = \min \left(a, \frac{b}{M} \right), \quad M = \max_{(x, y) \in \mathcal{R}} |g(x, y)|.$$

This is the **Cauchy-Peano theorem**.

Uniqueness

$$\frac{dy}{dx} = g(x, y)$$

- **Uniqueness:** If g is a continuous function of x and if there exists a constant $k > 0$ such that for all x, y_1 and y_2 in \mathcal{R} , we have

$$|g(x, y_1) - g(x, y_2)| \leq k |y_1 - y_2|,$$

then there exists a unique solution y to the above differential equation on $|x - x_0| \leq \alpha$, such that $y(x_0) = y_0$. The rectangle \mathcal{R} and the number α are defined as in the previous theorem.

This is the **Picard-Lindelöf Theorem**.