

Example

$$\frac{dy}{dx} = \frac{2}{\sqrt{\pi}} e^{-x^2}$$

9/7
1
(*)

$$F(x) = \int_0^x \frac{2}{\sqrt{\pi}} e^{-u^2} du$$

$$F'(x) = \frac{2}{\sqrt{\pi}} e^{-x^2}$$

$F(x)$ is a solution to (*)

$F(x)$ is called erf(x)

I. c. $y(1) = 0$

$$F(1) = \int_0^1 \frac{2}{\sqrt{\pi}} e^{-u^2} du$$


want $y(x) = F(x) - F(1)$

Example 1.2

9/7
2

$$\frac{dy}{dx} = \frac{1}{x^2}$$

$$y(x) = \frac{-1}{x} + C$$



Monotonicity

$$x^2 > 0$$

\Rightarrow always increasing

Concavity

$$\frac{d^2y}{dx^2} = \frac{-2}{x^3}$$

$x > 0 \Rightarrow \frac{d^2y}{dx^2} < 0 \Rightarrow$ concave down


$x < 0 \Rightarrow \frac{d^2y}{dx^2} > 0 \Rightarrow$ concave up

Suppose

$g(x)$ is even

9/7
3

$$g(-x) = g(x)$$



$$\frac{dy}{dx} = g(x)$$

Suppose $y(x)$ is a solution.

Look at $y(-x)$

$$\frac{d}{dx} y(-x) = y'(-x) (-1)$$

$$= -y'(-x)$$

$$= -g(x)$$

$$\frac{d}{dx} -y(-x) = g(x)$$

Conclusion: $y(x)$ is a solution

$\Rightarrow -y(-x)$ is a solution