

Families of Random Variables

Homework 7

Problems

- Let $X \sim Negbin(4, 1/3)$.
 - Find $P\{X > 12\}$
 - Find the smallest value for x so that $P\{X > x\} = 0.05, 0.01, 0.001$
 - Simulate 10,000 independent observations of $Negbin(4, 1/3)$ random variable and check your answers to (a) and (b) from the simulation.
- For some natural numbered-valued random variables, the case $\{X = 0\}$ cannot be observed. Thus, we can define a second random variable Y so that

$$P\{Y = y\} = P\{X = y | X \neq 0\}.$$

- Find the mass function for Y .
 - Give this mass function for $X \sim Bin(n, p)$
 - Give this mass function for $X \sim Geom(p)$
 - Find the mean for these random variables,
- On the other hand, the mass function for a zero inflated random variable Y is a mixture of a degenerate random variable taking only the value 0. Thus, the mass function

$$f_Y(y) = \alpha I_0(y) + (1 - \alpha)f_X(y)$$

for some natural numbered-valued random variable X (mean μ_X and variance σ_X^2) and some $\alpha \in (0, 1)$

- Verify that f_Y is a mass function.
- Find the mean of Y in terms of μ_X and σ_X^2 .
- Find the variance of Y in terms of μ_X and σ_X^2 .

Challenging Problems

- Let $X \sim Beta(\alpha, \beta)$
 - Write the density of X as an exponential family with natural parameters η_1, η_2 .
 - Find $E \ln X$
 - Find $EX \ln X$

- (d) What do these values tell you in the case of $X \sim \text{Beta}(1, 1) = \text{Unif}(0, 1)$?
2. Let X_1, \dots, X_n be independent and identically distributed with kurtosis α_4 .
- (a) Find the kurtosis $\alpha_{4,n}$ of \bar{X}_n , the average of X_1, \dots, X_n .
- (b) Find

$$\lim_{n \rightarrow \infty} \alpha_{4,n}.$$