

Central Limit Theorem*

Worksheet 14

1. The focal length f of an optical instrument is needed. This is determined by using the thin lens formula,

$$\frac{1}{r_1} + \frac{1}{r_2} = \frac{1}{f}.$$

where r_1 is the distance from the lens to the object and r_2 is the distance from the lens to the real image of the object. The distance r_1 is independently measured 12 times and r_2 is independently measured 40 times. The mean of the measurements is the actual distances, 9 centimeters and 18 centimeters, respectively. The standard deviation of the measurement is 0.1 centimeters for r_1 and 0.5 centimeter for r_2 .

- (a) Let \bar{R}_1 be the sample mean of the 12 measurements to the object. Estimate, using the central limit theorem, $P\{\bar{R}_1 < 8.9\text{cm}\}$.
- (b) Let \bar{R}_2 be the sample mean of the 40 measurements to the image. Estimate, using the central limit theorem, $P\{\bar{R}_2 < 17.9\text{cm}\}$.
- (c) How many measurements are needed so that $P\{|\bar{R}_2 - 18\text{cm}| > 0.1\text{cm}\} \leq 0.02$.
- (d) Would more or fewer measurements be needed so that
- $P\{|\bar{R}_2 - 18\text{cm}| > 0.1\text{cm}\} \leq 0.01$?
 - $P\{|\bar{R}_2 - 18\text{cm}| > 0.2\text{cm}\} \leq 0.02$?
- Explain your answer.
- (e) For measurements $r_{1,1}, \dots, r_{1,12}$ and $r_{2,1}, \dots, r_{2,40}$, estimate the focal length using

$$\frac{1}{\bar{r}_1} + \frac{1}{\bar{r}_2} = \frac{1}{\hat{f}}.$$

Use the delta method to give an estimate of the standard deviation of \hat{f} .

- (f) Simulate this protocol 10000 times using the `rnorm` command and compare the results from your simulation to the results from the delta method.

*©2019 Joseph C Watkins